

Phytotoxic effect of silver nanoparticles in *Triticum aestivum*: Improper regulation of photosystem I activity as the reason for oxidative damage in the chloroplast

A. RASTOGI^{*,+}, M. ZIVCAK^{**}, D.K. TRIPATHI^{***}, S. YADAV^{****}, H.M. KALAJI^{#,###}, and M. BRESTIC^{**,+}

*Department of Meteorology, Poznan University of Life Sciences, Piatkowska 94, 60-649 Poznan, Poland**

*Department of Plant Physiology, Slovak University of Agriculture, A. Hlinku 2, 94976 Nitra, Slovak Republic***

*Amity Institute of Organic Agriculture (AIOA) Amity University, Noida Sector 125, Noida, Uttar Pradesh- 201313, India****

*Department of Biotechnology, Hemvati Nandan Bahuguna Garhwal (Central) University, Srinagar Garhwal, Uttarakhand, 246174, India*****

Institute of Technology and Life Sciences (ITP), Falenty, Al. Hrabaska 3, 05-090 Raszyn, Poland#

White Hill Company, Żurawia 71/3, 15-540 Białystok, Poland###

Abstract

Silver nanoparticles (AgNPs) are used in several industries and their continuous release into environment can damage the ecosystem. Our study observed the impact of two concentrations of AgNPs (1 and 5 mM) on wheat seedlings. In this study, the treatment of AgNPs was found to have a significant impact on growth parameters, such as root and shoot length of wheat seedlings, in addition to that biochemical parameters, such as activity of catalase, glutathione, flavonoid, and chlorophyll concentrations, were also affected. AgNPs apparently suppressed the photosynthetic activity of the seedlings, with a clear destruction of photosystems at 5 mM concentration. The result also indicated an improper regulation of PSI electron transport, resulting in a damage to chloroplast structure, including photosystem itself. Our study presents a mechanism of the AgNPs action of on photosynthetic processes in plants.

Additional key words: antioxidants; chlorophyll fluorescence; JIP test.

Introduction

Nanoparticles cover a heterogeneous range of materials (Santos *et al.* 2015), but only a few of them are extensively used. The present environment is at risk to be exposed by extensively used nanoparticles. Silver nanoparticles (AgNPs) are among one of the most used nanoparticles, which can be found in various products. AgNPs can be used as antimicrobial agents, waste-water treatment agent, shampoo, toothpaste, soap, food-packaging materials, food-storage containers, room sprays, detergents, fabrics, paint, *etc.* (Boxall *et al.* 2007, Rai *et al.* 2009, Wijnhoven *et al.* 2009). Recent reports have shown that about 6% of

AgNPs produced in the United States ends up in sludge, whereas around 3.5% in surface water; thus, AgNPs may contaminate agricultural lands (Khaydarov *et al.* 2009, El-Temseh and Joner 2012).

Previous studies have shown that AgNPs may interfere with plant metabolism and may have a positive or negative impact, which depends on its concentration, size, and properties (Gruyer *et al.* 2014, Pallavi *et al.* 2016, Jasim *et al.* 2017, Rastogi *et al.* 2017). Dimkpa *et al.* (2013), have shown that AgNPs of 10-nm size, when hydroponically applied at a concentration of 2.5 mg kg⁻¹, are nontoxic for wheat, whereas, when present in excess they causes a toxic response in the form of morphological variation and

Received 4 March 2018, *accepted* 19 July 2018.

⁺Corresponding authors; phone: 0048-53 51 49922, e-mail: anshu.rastogi@up.poznan.pl (A. Rastogi); phone: 00421- 37 61 44448, e-mail: marian.brestic@uniag.sk (M. Brestic)

Abbreviations: ABS – absorption; AgNPs – silver nanoparticles; UV – ultra-violet radiation; FLAV – flavonoids; GSH – glutathione; CAT – catalase; OEC – oxygen-evolving complex; OEC_{act} – active oxygen-evolving complex; SSA – 5-sulfosalicylic acid; F_m – maximum fluorescence intensity; NPQ – nonphotochemical quenching; F_v/F_m – photosynthetic efficiency of PSII under dark adaptation; P_m – maximum amplitude of P700 signal; Φ_{PSII} – maximum quantum yield of PSII photochemistry; Φ_{NPQ} – quantum yield of regulated nonphotochemical energy loss in PSII; Φ_{NO} – quantum yield of nonregulated nonphotochemical energy loss in PSII; Φ_{PSI} – maximum quantum yield of PSI; Φ_{ND} – quantum yield of regulated nonphotochemical energy loss in PSI; Φ_{NA} – quantum yield of nonregulated nonphotochemical energy loss in PSI; Φ_{po} – photochemical efficiency; PI_{abs} – overall photochemical performance; ROS – reactive oxygen species; Ψ_{ET0} – electron transport at PSII acceptor side; RC/ABS – fraction of active PSII reaction centers.

Acknowledgement: This work was supported by the research project of the Slovak Research and Development Agency under the project APVV-15-0721, VEGA 1/0923/16, and National Science Centre Poland, under the project UMO-2016/21/B/ST10/02271. AR thanks to Slovak Academic Information Agency (SAIA) for providing National Scholarship for research in the year 2017.

accumulation of oxidized glutathione. The AgNP of 10 nm was observed to enhance a root length at a concentration of 2.5 mg L⁻¹ in barley, whereas it reduced the root length at a higher concentration (Gruyer *et al.* 2014). All the studied concentrations of AgNP were observed to reduce the root length when applied on lettuce, whereas for radish, the studied concentrations of AgNP were not able to cause any significant impact (Gruyer *et al.* 2014). The biosynthesized silver nanoparticles of 8–21 nm diameter at 1 µm mL⁻¹ concentration were observed to significantly impact the growth parameters such as root length, shoot length, and fresh mass of fenugreek seedlings (Jasim *et al.* 2017). The foliar treatment of wheat seedlings by 35-nm AgNPs was observed to be ineffective in causing positive or negative impact at studied concentrations (Pallavi *et al.* 2016). Several other studies have indicated that AgNPs of different sizes in variable concentrations can lead to morphological changes in plants, in addition, it can also influence the reduction in root/shoot and overall biomass (Hawthorne *et al.* 2012, Yin *et al.* 2012, Mirzajani *et al.* 2013, Pokhrel and Dubey 2013, Qian *et al.* 2013, Zuverza-Mena *et al.* 2016, Tripathi *et al.* 2017, Vinković *et al.* 2017). Different studies have shown that the plants exposed to AgNPs produce a higher amount of reactive oxygen species (ROS) and antioxidant molecules (Dimkpa *et al.* 2013, Jiang *et al.* 2014, Belava *et al.* 2017, Tripathi *et al.* 2017). Few studies have shown a significant decline in photosynthetic pigment contents as a response to AgNPs exposure (Qian *et al.* 2013, Tripathi *et al.* 2017). Whereas, according to our knowledge, only a few studies have been focused on the impact of AgNPs on photosynthetic efficiency of plants.

Photosynthesis is an important, and one of the most complex phenomena in plants. Chlorophyll (Chl) fluorescence emission is a process which is going in parallel with photosynthesis in plants, therefore, photosynthetic process and Chl fluorescence are closely related. It has been now established, that changes in photosynthetic activity due to different environmental factors can be detected through the observation of the changes in Chl fluorescence (Kalaji *et al.* 2014). Analysis of Chl fluorescence parameters can be used as a precise tool for testing a direct impact of adverse environmental conditions on photosynthesis and, therefore, the indirect assessment of their impact on plants (Kalaji *et al.* 2014). Therefore, the study of photosynthetic efficiency is needed to understand the plant–nanoparticle interaction. CuO nanoparticles were observed to influence photosynthesis in *Lemna gibba* L. (Perreault *et al.* 2014). The authors observed a significant difference in Chl *a* fluorescence, which indicated a decrease in the number of active PSII reaction centers, as a result to CuO nanoparticles exposure.

Because of its economic importance, wheat was chosen as an experimental crop in this study. The phytotoxicity of AgNPs on wheat seedlings was analyzed by observing variations in morphology, physiology, and contents of biochemical compounds.

Materials and methods

Silver nanoparticle preparation: The nano-powder

of AgNP of particle size <100 nm was purchased from SIGMA–ALDRICH®. The nanoparticles were containing polyvinylpyrrolidone (PVP) as a dispersant. The nanoparticles were sterilized for 20 min under ultraviolet radiation (UV), in laminar flow. The stock of 100 mM AgNP suspension was prepared in sterile distilled water using sterile glassware. The nanoparticle suspension was prepared using the ultrasonic bath for 3 h; the water in the ultrasonic bath was exchanged with the ice-cold water, every 20 min. A homogeneously dispersed nanoparticle suspension was obtained. Different dilutions (1 and 5 mM) of AgNP were prepared with sterile water. Nanoparticle suspension of 20 ml was added to sterile petri plates containing UV-treated filter paper.

Seed preparation and growth conditions: The seeds of spring wheat (*Triticum aestivum*) cv. “Corso” were obtained from phenotyping laboratory of Slovak Agriculture University in Nitra. Spring genotype was selected because of its fast development and higher initial vitality. The seeds were washed for 1 h in running tap water; surface sterilized with 0.1% HgCl₂ for 2.5 min with 80% ethanol, then the seeds were washed three times with sterile distilled water. The seeds were distributed in Petri plates with 0, 1, and 5 mM AgNPs (15 seeds each). The seeds were grown in a plant chamber with 250 µmol(photon) m⁻² s⁻¹, photoperiod of 16-h light/8-h dark, relative air humidity 60–70%, and day temperature of 22°C and 18°C at night for three weeks. Three replica plates were kept for each condition.

Plant morphological analysis: The growth of seedlings was observed at regular intervals for their morphological variations. Root and shoot sizes were measured using graph paper and scale. The final measurement for root and shoot sizes were performed after three weeks of the AgNPs treatment. The statistical analysis was done by random measurements of ten plants from three different plates under the same conditions. The root morphology was observed using Nikon Optiphot (Japan) light microscope fitted with Canon (Japan) digital camera.

Catalase and glutathione: Catalase activity (EC 1.11.1.6) and reduced glutathione (GSH) from the wheat root was measured using catalase assay kit (CAT100, SIGMA–ALDRICH) and glutathione assay kit (CS0260, SIGMA–ALDRICH), respectively. For catalase, 0.5 g of fresh roots were collected and homogenized with 5 mL of 50 mM potassium phosphate buffer (PBS, pH 7.0) containing 0.4% PVP in a mortar and pestle on the ice. The homogenate was further centrifuged at 10,000 × g for 20 min at 4°C. The supernatant was collected and examined for catalase activity following the protocol provided with the kit. The reaction time was kept for 1 min for 10 µl of the supernatant. The spectroscopic measurement was taken at 240 nm using Cary 60 UV-Vis spectrophotometer (Agilent, Santa Clara, USA). Catalase was calculated according to the kit and expressed as enzyme Unit per gram of fresh mass, where enzyme Unit is amount required for transformation of 1 µmol of the hydrogen peroxide per min at standard

conditions. For the measurement of GSH, the root tissues were frozen dry with liquid nitrogen immediately after excision. The tissues were grounded in a mortar and pestle with liquid nitrogen to obtain a fine powder. Powdered sample of 0.1 g was taken and mixed with 300 μ l of 5% 5-sulfosalicylic acid (SSA) and vortexed vigorously for 1 min, then 2,100 μ l of 5% SSA was added and homogenized with a 3-ml PTFE pestle in a glass tube for 5 min. The samples were adapted to 4°C and centrifuged at 10,000 \times g for 10 min at 4°C. The further analyses were performed according to the protocol provided with the kit and expressed as nmol per gram of dry mass. The spectroscopic measurement was done at 412 nm using *Cary 60 UV-Vis* spectrophotometer (*Agilent*, Santa Clara, USA). Five repetitions were performed for both antioxidants to calculate standard deviation.

Chl and flavonoid: The Chl content was measured non-invasively by using *SPAD-502* (*Minolta Camera Co.*, Osaka, Japan). The Chl meter readings were obtained from the middle of the grown leaves. The flavonoid content was measured noninvasively by using *Multiplex-3fluorimetric* sensor (*Force-A*, Orsay, France). The *Multiplex* measures fluorescence emitted by Chl, in the red and far-red spectral regions, under excitation by light-emitting diodes of 375, 450, 515, and 630 nm. The synchronized filtered-photodiode detectors recorded fluorescence in yellow, red, and far-red (Zivcak *et al.* 2017). The relative amount of flavonoids (FLAV) was calculated observing Chl fluorescence signals under UV (FRF-UV) and red excitation (FRF-R) (Syta *et al.* 2015).

The measurements were performed on nine randomly selected plants for each treatment, by taking the average value of three measurements from the same seedling. The measurements were converted to the percentage assuming the mean value of the control to be 100%.

P700 redox state and Chl fluorescence: P700 redox state and Chl *a* fluorescence were simultaneously measured through *Dual PAM-100* (*Heinz Walz GmbH*, Germany). The plants were transferred to laboratory bench and dark-adapted for 20 min in a dark box, and around 2 min in the measuring head. To activate the photosynthetic processes, a light intensity of 134 μ mol(photon) $m^{-2} s^{-1}$ (similar to ambient light) was used. Once the steady state was reached, a rapid light curve was triggered [at light intensities of 14, 30, 61, 103, 134, 174, 224, 347, 539, 833, 1,036; 1,295; 1,602; and 1,930 μ mol(photon) $m^{-2} s^{-1}$; 30 s at each light intensity] with saturation pulse and far-red pulse. The measurements were performed on three plants for each experimental conditions, and different parameters were calculated as described previously in Brestic *et al.* (2016).

Chl fluorescence induction curve was measured using *Handy PEA* fluorometer (*Hansatech Instruments Ltd.*, Norfolk, UK). The plants were transferred to laboratory bench for the measurement, the plants were dark-adapted for 20 min using the clip provided by the producer. Measurements were performed following the standard protocol with a 3,500 μ mol(photon) $m^{-2} s^{-1}$ of light intensity (as max) and a 1-s duration of the pulse light

(saturation light). Analysis of measured Chl *a* fluorescence signal was based on a JIP test (Strasser and Strasser 1995, Srivastava *et al.* 1999, Strasser *et al.* 2000, 2004, 2010). The measurement was performed in the middle of the oldest leaf from nine plants for a treatment. The Chl induction curve was drawn from the mean of the obtained data points.

Results

Plant morphology: The wheat seedlings were analyzed for the root and shoot growth each week. The toxic concentration of AgNPs for this study was determined by observing the morphological changes in the seedlings after a week of germination (Fig. 1S, *supplement*). More than 50% decrease in root and shoot growth was observed after three weeks of germination (Table 1, Fig. 1A). At 5 mM AgNPs concentration the seedling roots were observed to avoid its contact with filter paper (Fig. 2S, *supplement*). The microscopic study showed that the roots at 5 mM concentration were darker in comparison to control and a large number of root hairs was observed at 5 mM concentration (Fig. 1B). The results clearly indicated that the concentrations studied were phytotoxic for wheat.

Table 1. Impact of AgNPs on root and shoot length of wheat after 3-week exposure. The data are presented as mean of at least three measurements with its standard deviation.

AgNPs [mM]	Root [cm]	Shoot [cm]
0	11.39 \pm 1.22	12.25 \pm 2.06
1	5.54 \pm 0.89	5.2 \pm 0.6
5	4.2 \pm 0.56	4.94 \pm 0.73

Catalase and glutathione: The catalase activity increased by 50% in plant roots treated with 1 mM AgNPs, whereas it was 100% higher in roots treated with 5 mM AgNPs when compared with control plants (Fig. 2A). The GSH content also increased in plant roots by approximately 100 and 300% at 1 mM and 5 mM AgNPs, respectively (Fig. 2B). The results indicate the increase in catalase activity and in the amount of GSH as the protective molecules in plant roots in response to AgNPs exposure.

Total Chl concentration and flavonoids: The total Chl in wheat decreased by 20% at both studied concentrations of AgNPs (Fig. 3A). As a response to 1 mM and 5 mM AgNPs exposure, the flavonoid content of the wheat leaves increased by 100 and 200%, respectively (Fig. 3B). The results indicate the decrease in total Chl content, whereas a significant increase in flavonoid content as a response to AgNPs exposure of wheat plants.

P700 redox state and Chl fluorescence: Fig. 4 indicates the response of different fluorescence parameters to 1 mM and 5 mM AgNPs. P_m indicates the maximum amplitude of P700 signal, which decreased by around 15 and 50% in 1 mM and 5 mM AgNPs-treated plants, respectively, when compared with control (Fig. 4A).




Fig. 1. Impact of AgNPs on wheat morphology: (A) A photograph of wheat plants after 3-week exposure to 0, 1, and 5 mM AgNPs. (B) A microscopic image of root from 0 and 5 mM AgNPs-exposed plants.




Fig. 2. Impact of AgNPs on antioxidants: (A) The enzymatic activity of catalase when exposed to AgNPs. The measurement was taken from plant root in enzyme Unit per gram fresh mass [U g⁻¹(FM)]. (B) The GSH content of the wheat root under different exposure to AgNPs. The amount of GSH was expressed in nmol mg⁻¹(DM).

F_v/F_m indicates the estimated maximum quantum efficiency of PSII photochemistry, which decreased by 12 and 80%, whereas, nonphotochemical quenching (NPQ) decreased by 10 and 85% in 1 mM and 5 mM AgNPs-treated plants, respectively, when compared with control (Fig. 4B,C). ϕ_{PSII} indicates the estimated effective quantum yield of PSII at given PAR, which decreased by 25 and 95% at 833 PAR for 1 mM and 5 mM AgNPs-treated plants, respectively, when compared with control (Fig. 4D). ϕ_{NPQ} indicates the quantum yield of pH-dependent energy dissipation in PSII, whereas ϕ_{NO} indicates quantum




Fig. 3. Relative content of chlorophyll and flavonoids in wheat leaves under different AgNPs exposure conditions.

yield of nonregulated energy dissipation in PSII. At 1 mM AgNPs concentration, only 4% decrease in ϕ_{NPQ} , and 5% increase in ϕ_{NO} was observed, whereas, the differences were observed to be very significant for these parameters, in case of 5 mM AgNPs, when compared with control (Fig. 4E,F). Parameter ϕ_{PSI} represent the photochemical quantum yield of PSI; it slightly increased in case of 1 mM AgNPs concentration, whereas a significant decrease in ϕ_{PSI} was observed in the plant sample treated with 5 mM AgNPs when compared with control. Parameter ϕ_{ND} , indicating the nonphotochemical quantum yield of PSI, which is oxidized at a given state, decreased by 70 and 40% for 1 mM and 5 mM samples, respectively (Fig. 4H). Parameter ϕ_{NA} , which indicates nonphotochemical quantum yield of PSI that cannot be oxidized by a saturation pulse at a given




Fig. 4. Impact of AgNPs on photochemistry of wheat plants: P_m – maximum amplitude of P700 signal (A); F_v/F_m – photosynthetic efficiency of PSII under dark adaptation (B); NPQ – nonphotochemical quenching (C); Φ_{PSII} – maximum quantum yield of PSII photochemistry (D); Φ_{NPQ} – quantum yield of regulated nonphotochemical energy loss in PSII (E); Φ_{NO} – quantum yield of nonregulated nonphotochemical energy loss in PSII (F); Φ_{PSI} – maximum quantum yield of PSI (G); Φ_{ND} – quantum yield of regulated nonphotochemical energy loss in PSI (H); Φ_{NA} – quantum yield of nonregulated nonphotochemical energy loss in PSI (I).

state due to lack of acceptors, was 125% higher in case of 1 mM AgNPs-treated samples, whereas it was 100% higher for 5 mM AgNPs-treated samples in comparison with controls (Fig. 4I).

The Chl *a* fluorescence induction curve with OJIP steps obtained from wheat plants showed a significant change under AgNPs treatment; it was almost linear under 5 mM concentration (Fig. 5A). Some of the important photosynthetic parameters, such as photochemical efficiency (Φ_{P_0}), overall photochemical performance index (PI_{abs}), electron transport at PSII acceptor side (Ψ_{ET_0}), fraction of active PSII reaction centers (RC/ABS), and maximum fluorescence intensity (F_m), were calculated according to Strasser (1997), whereas a fraction of active oxygen evolving complex (OEC_{act}) was calculated according to Chen and Cheng (2009) (Fig. 5B). The results indicate a clear impact of AgNPs on P700 redox state and Chl fluorescence in wheat plants.

Discussion

Our research confirmed previous reports on phytotoxic properties of AgNPs (Rastogi *et al.* 2017) by showing the reduction in root and shoot growth (Table 1, Fig. 1A). The

studied AgNPs were <100 nm in diameter, whereas size exclusion limit for plant cell wall has been reported to be in between 5–20 nm (Dietz and Herth 2011), which may restrict its absorption by root cells. It has been previously observed that at high concentrations, the nanoparticles may destroy the cell structure and enter the root cells (Mirzajani *et al.* 2013), which is also supported by the observation that some of the nanoparticles may induce larger pores in the cell wall (Navarro *et al.* 2008, Kurepa *et al.* 2010). Therefore, it can be expected that at high concentrations the studied AgNPs may create an additional pore in the cell wall and may enter the plant. The microscope pictures of roots showed a clear dark spot in roots of the 5 mM AgNPs-treated plant (Fig. 1B), which indicates the presence of AgNPs in root cells. At 5 mM AgNPs concentration, the plants were observed to avoid its contact with filter paper, and therefore absorbed the water from the air. Due to plant dependence on atmospheric water, the root hairs were observed to be significantly developed in plants treated with 5 mM AgNPs in comparison with control.

Several studies have shown that the phytotoxic effect on plants is mediated through oxidative burst (Hossain *et al.* 2015, Rastogi *et al.* 2017), which may initiate a protective mechanism in plant cells by increasing the production of




Fig. 5. OJIP test analysis. (A) The fluorescence induction curve and (B) different photosynthetic parameters (Φ_{po} – photochemical efficiency; F_m – maximum fluorescence intensity; OEC_{act} – active oxygen evolving complex; RC/ABS – fraction of active PSII reaction centres; Ψ_{ETo} – electron transport at PSII acceptor side; PI_{abs} – overall photochemical performance index) after exposure of wheat to different concentrations of AgNPs.

antioxidants. Catalase is the enzyme known to catalyze the decomposition of hydrogen peroxide to water and oxygen. An increase in catalase activity indicates an increased amount of hydrogen peroxide, whereas, an increased amount of GSH in AgNPs-treated samples indicates a higher production of glutathione (Fig. 2).

Chl concentration is an important indicator for accessing the photosynthesis and growth conditions, whereas flavonoids are molecules which have been detected to be increased under different abiotic stress conditions (Syta *et al.* 2015, Schulz *et al.* 2016, Zivcak *et al.* 2017). The decrease in the Chl concentration and the increase in flavonoids concentration (Fig. 3) confirmed that studied AgNPs were phytotoxic to wheat.

In addition to other physiological responses, we observed the severe effect of AgNPs on PSI and PSII photochemistry. Whereas the effect of 5 mM AgNPs was evidently lethal, suppressing almost fully the photochemical activities in chloroplasts (Fig. 4), the effect of 1 mM AgNPs was much more interesting. With a decrease in PSII quantum efficiency, a moderate impact of nanoparticles on PSII photochemistry under high light indicated a decrease in electron transport rate by 30%. Typically, under stress conditions, the decrease of CO₂ assimilation limits the demand for NADPH and ATP, resulting in a decrease of the rate of linear electron

transport, but also in the accumulation of H⁺ in thylakoid lumen. The phenomena result in a significant increase of NPQ and the downregulation of linear electron flow at cytochrome *b₆f*. This is one of the most important photoprotective responses in the chloroplast, which protect the electron transport chain against over-reduction (Joliot and Johnson 2011). However, we observed a decrease in NPQ instead of its increase, which indicates that the regulatory mechanism is not working properly under AgNPs treatment (Fig. 4C–F). This effect is even more evident when looking at the redox poise of PSI donor (Φ_{ND}) and acceptor (Φ_{NA}) side (Fig. 4H,I). It is obvious that PSI acceptor side is highly over-reduced. This state of PSI acceptor side was shown to be responsible for the excessive production of ROS, especially hydroxyl radicals, which may attack sensitive structures in chloroplasts and the whole cell (Takagi *et al.* 2016). It was clearly shown that, under high light conditions, high level of P700 oxidation (high Φ_{ND}) is crucial for the protection of the PSI acceptor side against over-reduction. However, under conditions of 1 mM AgNPs, the Φ_{ND} was observed to be very low. The observed phenomena may be caused by insufficient downregulation of linear electron transport at cytochrome *b₆f*, probably due to low trans-thylakoid proton gradient. A similar effect was observed in mutants with a low PSI content (Brestic *et al.* 2015) or with a mutation limiting the ability to drive the cyclic electron flow around PSI (Grieco *et al.* 2012). However, increase in Φ_{PSII} , in parallel with a decrease in Φ_{PSII} suggest that there was an increase of cyclic electron flow due to 1 mM AgNPs treatment. It has been previously reported that, under some specific conditions such as high temperature, excessive leakage of H⁺ through thylakoid membrane can occur (Bukhov *et al.* 1999). The excessive leakage of H⁺ may lead to low ΔpH , which results in to over-reduction of PSI acceptor side despite a high cyclic electron flow (Brestic *et al.* 2016). Thus, we can hypothesize that the AgNPs may cause the leak of H⁺ through thylakoid membrane or proton channels and result in a decrease in ΔpH and, hence, the proton motive force, which may negatively influence the ATP synthesis in chloroplasts (Zhang *et al.* 2009).

One of the direct effects of improper regulation of linear electron transport is the PSI photoinhibition. Our measurements indicate an evident decrease in the content of active PSI unit, indicating PSI photoinhibition. Whereas the PSII photoinhibition is common, PSI in wild type plants is thought to be very resistant to photodamage (Powles 1984), except chilling stress (Sonoike 1996) or fluctuating light (Grieco *et al.* 2012) conditions. Recently, it was assumed that the photoinhibition of PSI may represent a protective mechanism against oxidative damage of key cell structures under conditions where the linear electron transport cannot be well regulated (Brestic *et al.* 2016, Tiwari *et al.* 2016). The nonfunctional PSI centres turn into PSI-NPQs of excitation energy and dissipate thermally the surplus of excitation energy. Photodamage of PSI increases the antenna cross-section of PSI, and conversely decreases that of PSII, through thylakoid protein phosphorylation-dependent mechanisms, thus maximizing the amount of excess energy dissipation by PSI. It can represent the

mechanism of high physiological significance for the survival of plants under specific conditions, in which the flexible mechanisms are not working properly (Tiwari *et al.* 2016). However, it was shown that the photodamage of PSI may have a high physiological significance, and it indicates a decrease in photosynthetic efficiency and CO₂ assimilation capacity, as well as photoprotection of PSII (Zivcak *et al.* 2015).

Thus, it is obvious that the photochemistry, especially the processes at the level of PSI, is severely affected by the AgNPs treatment. The acceptor site of PSI can be a major source of free radicals (ROS), which are responsible for the oxidative damage observed in this study as well as previously reported by the other authors (Oukarroum *et al.* 2013, Tripathi *et al.* 2017, Rastogi *et al.* 2017). ROS produced at PSI acceptor side can cause damage of other chloroplast structures, such as PSII; this may explain the effects of AgNPs on PSII photochemistry well demonstrated by the analyses of OJIP transient (Fig. 5). In addition to a decrease in Φ_{Po} and PI_{abs} , which was evident also from the PAM measurements, we observed several important phenomena related to the status of PSII photochemistry. First, a significant decrease of RC/ABS ratio can be attributed mostly to a decrease in the fraction of active PSII RCs. It means that presence of AgNPs leads to a severe inactivation of PSII even at the lower concentration. This can be partly explained by the presence of enhanced K-step in OJIP transient, which indicates a decrease in PSII donor side activity (Strasser 1997). The oxygen evolving complex (OEC) represents one of the sensitive sites, frequently damaged by ROS produced in chloroplasts (Vass 2012). Although the direct effect of AgNPs on OEC cannot be excluded, the indirect effect through ROS produced at the PSI acceptor side is more probable. Interestingly, we observed only a moderate limitation of electron transport at PSII acceptor side (Ψ_{ET_0}), which is a typical response to many stress factors (Kalaji *et al.* 2018). The maximum fluorescence signal was still relatively high, which indicates that the decrease in photochemical activity was not associated with a severe molecular degradation of PSII complexes. As the photoinhibition of PSII resulting from a high excitation pressure at PSII acceptor side is typically associated with a decrease in F_m (Baker and Rosenquist 2004), the observed changes in PSII photochemistry cannot be attributed dominantly to the light-induced PSII damage with a typical turnover of PSII proteins. At least, the repair of damaged PSII is probably not working and inactive PSII are accumulated without recovery. In this respect, we can hypothesize that AgNPs may inhibit the repair of PSII, which may have more serious consequences such as inhibition of PSII activity. Such an effect was also observed under other environmental stresses (Nishiyama *et al.* 2006). The effect can be caused by the possible insufficient production of ATP needed for PSII repair due to impaired activity of PSI discussed above. Thus, it can be concluded that the studied AgNPs is phytotoxic for wheat seedlings and cause a severe impact on the photosynthetic process, which occurs due to the damage of PSI in the chloroplasts.

References

- Baker N.R., Rosenqvist E.: Applications of chlorophyll fluorescence can improve crop production strategies: an examination of future possibilities. – *J. Exp. Bot.* **55**: 1607-1621, 2004.
- Belava V.N., Panyuta O.O., Yakovleva G.M. *et al.*: The effect of silver and copper nanoparticles on the wheat-*Pseudocercospora herpotrichoides* pathosystem. – *Nanoscale Res. Lett.* **12**: 250, 2017.
- Boxall P., Purcell J., Wright P.: Human resource management: scope, analysis and significance. – In: Boxall P., Purcell J., Wright P. (ed.): *The Oxford Handbook of Human Resource Management*. Pp. 1-18. Oxford University Press, Oxford 2007.
- Brestic M., Zivcak M., Kunderlikova K. *et al.*: Low PSI content limits the photoprotection of PSI and PSII in early growth stages of chlorophyll *b*-deficient wheat mutant lines. – *Photosynth. Res.* **125**: 151-166, 2015
- Brestic M., Zivcak M., Kunderlikova K., Allakhverdiev S.I.: High temperature specifically affects the photoprotective responses of chlorophyll *b*-deficient wheat mutant lines. – *Photosynth. Res.* **130**: 251-266, 2016.
- Bukhov N.G., Wiese C., Neimanis S., Heber U.: Heat sensitivity of chloroplasts and leaves: leakage of protons from thylakoids and reversible activation of cyclic electron transport. – *Photosynth. Res.* **59**: 81-93, 1999.
- Chen L.S., Cheng L.: Photosystem 2 is more tolerant to high temperature in apple (*Malus domestica* Borkh.) leaves than in fruit peel. – *Photosynthetica* **47**: 112-120, 2009.
- Dietz K.-J., Herth S.: Plant nanotoxicology. – *Trends Plant Sci.* **16**: 582-589, 2011.
- Dimkpa C.O., McLean J.E., Martineau N. *et al.*: Silver nanoparticles disrupt wheat (*Triticum aestivum* L.) growth in a sand matrix. – *Environ. Sci. Technol.* **47**: 1082-1090, 2013.
- El-Temseh Y.S., Joner E.J.: Ecotoxicological effects on earthworms of fresh and aged nano-sized zero-valent iron (nZVI) in soil. – *Chemosphere* **89**: 76-82, 2012.
- Grieco M., Tikkanen M., Paakkanen V. *et al.*: Steady-state phosphorylation of light-harvesting complex II proteins preserves photosystem I under fluctuating white light. – *Plant Physiol.* **160**: 1896-1910, 2012.
- Gruyer N., Dorais M., Bastien C. *et al.*: Interaction between silver nanoparticles and plant growth. – In: Son J.E., Lee I.B., Oh M.M. (ed.): *International Symposium on New Technologies for Environment Control, Energy-Saving and Crop Production in Greenhouse and Plant*. Pp. 795-800. ISHS Acta Horticulturae, Jeju, Korea 2014.
- Hawthorne J., Musante C., Sinha S.K., White J.C.: Accumulation and phytotoxicity of engineered nanoparticles to *Cucurbita pepo*. – *Int. J. Phytoremediat.* **14**: 429-442, 2012.
- Hossain Z., Mustafa G., Komatsu S.: Plant responses to nanoparticle stress. – *Int. J. Mol. Sci.* **16**: 26644-26653, 2015.
- Jasim B., Thomas R., Mathew J., Radhakrishnan E.K.: Plant growth and diosgenin enhancement effect of silver nanoparticles in Fenugreek (*Trigonella foenumgraecum* L.). – *Saudi Pharm. J.* **25**: 443-447, 2017.
- Jiang H.S., Qiu X.N., Li G.B. *et al.*: Silver nanoparticles induced accumulation of reactive oxygen species and alteration of antioxidant systems in the aquatic plant *Spirodela polyrhiza*. – *Environ. Toxicol. Chem.* **33**: 1398-1405, 2014.
- Joliot P., Johnson G.N.: Regulation of cyclic and linear electron flow in higher plants. – *P. Natl. Acad. Sci. USA* **108**: 13317-13322, 2011.
- Kalaji H.M., Rastogi A., Živčák M. *et al.*: Prompt chlorophyll fluorescence as a tool for crop phenotyping: an example of barley landraces exposed to various abiotic stress factors. –

- Photosynthetica **56**: 953-961, 2018.
- Kalaji H.M., Schansker G., Ladle R.J. *et al.*: Frequently asked questions about *in vivo* chlorophyll fluorescence: practical issues. – *Photosynth. Res.* **122**: 121-158, 2014.
- Khaydarov R.R., Khaydarov R.A., Gapurova O. *et al.*: Antimicrobial effects of silver nanoparticles synthesized by an electrochemical method. – In Reithmaier J.P., Petkov P., Kulisch W., Popov C. (ed.): *Nanostructured Material for Advanced Technological Applications*. NATO Science for Peace and Security Series B: Physics and Biophysics. Pp. 215-218. Springer, Dordrecht 2009.
- Kurepa J., Paunesku T., Vogt S. *et al.*: Uptake and distribution of ultrasmall anatase TiO₂ Alizarin red S nanoconjugates in *Arabidopsis thaliana*. – *Nano Lett.* **10**: 2296-2302, 2010.
- Mirzajani F., Askari H., Hamzelou S. *et al.*: Effect of silver nanoparticles on *Oryza sativa* L. and its rhizosphere bacteria. – *Ecotoxicol. Environ. Safe.* **88**: 48-54, 2013.
- Navarro E., Piccapietra F., Wagner B. *et al.*: Toxicity of silver nanoparticles to *Chlamydomonas reinhardtii*. – *Environ. Sci. Technol.* **42**: 8959-8964, 2008.
- Nishiyama Y., Allakhverdiev S.I., Murata N.: A new paradigm for the action of reactive oxygen species in the photoinhibition of photosystem II. – *BBA- Bioenergetics.* **1757**: 742-749, 2006.
- Oukarroum A., Barhoumi L., Pirastru L., Dewez D.: Silver nanoparticle toxicity effect on growth and cellular viability of the aquatic plant *Lemna gibba*. – *Environ. Toxicol. Chem.* **32**: 902-907, 2013.
- Pallavi, Mehta C.M., Srivastava R. *et al.*: Impact assessment of silver nanoparticles on plant growth and soil bacterial diversity. – *3 Biotech.* **6**: 254, 2016.
- Perreault F., Samadani M., Dewez D.: Effect of soluble copper released from copper oxide nanoparticles solubilisation on growth and photosynthetic processes of *Lemna gibba* L. – *Nanotoxicology.* **8**: 374-382, 2014.
- Pokhrel L.R., Dubey B.: Evaluation of developmental responses of two crop plants exposed to silver and zinc oxide nanoparticles. – *Sci. Total Environ.* **452-453**: 321-332, 2013.
- Powles S.B.: Photoinhibition of photosynthesis induced by visible light. – *Annu. Rev. Plant Physiol.* **35**: 15-44, 1984.
- Qian H., Peng X., Han X. *et al.*: Comparison of the toxicity of silver nanoparticles and silver ions on the growth of terrestrial plant model *Arabidopsis thaliana*. – *J. Environ. Sci.-China* **25**: 1947-1955, 2013.
- Rai M., Yadav A., Gade A.: Silver nanoparticles as a new generation of antimicrobials. – *Biotechnol. Adv.* **27**: 76-83, 2009.
- Rastogi A., Zivcak M., Sytar O. *et al.*: Impact of metal and metal oxide nanoparticles on plant: A critical review. – *Front. Chem.* **5**: 78, 2017.
- Santos C.S.C., Gabriel B., Blanchy M. *et al.*: Industrial applications of nanoparticles – A prospective overview. – *Mater. Today-Proc.* **2**: 456-465, 2015.
- Schulz E., Tohge T., Zuther E. *et al.*: Flavonoids are determinants of freezing tolerance and cold acclimation in *Arabidopsis thaliana*. – *Sci. Rep.* **6**: 34027, 2016.
- Sonoike K.: Photoinhibition of photosystem I: Its physiological significance in the chilling sensitivity of plants. – *Plant Cell Physiol.* **37**: 239-247, 1996.
- Srivastava A., Strasser R.J., Govindjee: Greening of peas: parallel measurements of 77K emission spectra, OJIP chlorophyll *a* fluorescence transient, period four oscillation of the initial fluorescence level, delayed light emission and P700*. – *Photosynthetica* **37**: 365-392, 1999.
- Strasser B.J., Strasser R.J.: Measuring fast fluorescence transients to address environmental questions: The JIP-test. – In: Mathi P. (ed.): *Photosynthesis: from Light to Biosphere*. Pp. 977-980. Kluwer Academic Publishers, Dordrecht 1995.
- Strasser R.J., Tsimilli-Michael M., Qiang S., Goltsev V.: Simultaneous *in vivo* recording of prompt and delayed fluorescence and 820-nm reflection changes during drying and after rehydration of the resurrection plant *Haberlea rhodopensis*. – *Biochim. Biophys. Acta* **1797**: 1313-1326, 2010.
- Strasser R.J., Tsimilli-Michael M., Dangre D., Rai M.: Biophysical phenomics reveals functional building blocks of plants system biology: a case study for evaluation of the impact of Mycorrhization with *Piriformospora indica*. – In: Varma A., Oelmüller R. (ed.): *Advanced Techniques in Soil Biology*. Pp. 319-338. Springer, Dordrecht 2004.
- Strasser R.J., Srivastava A., Tsimilli-Michael M.: The fluorescence transient as a tool to characterize and screen photosynthetic samples. – In: Yunus M., Pathre U., Mohanty P. (ed.): *Probing Photosynthesis: Mechanisms, Regulation and Adaptation*. Pp. 445-483. Taylor & Francis, London 2000.
- Strasser B.J.: Donor side capacity of photosystem II probed by chlorophyll *a* fluorescence transient. – *Photosynth. Res.* **52**: 147-155, 1997.
- Sytar O., Bruckova K., Hunkova E. *et al.*: The application of multiplex fluorimetric sensor for the analysis of flavonoids content in the medicinal herbs family Asteraceae, Lamiaceae, Rosaceae. – *Biol. Res.* **48**: 5, 2015.
- Takagi D., Takumi S., Hashiguchi M. *et al.*: Superoxide and singlet oxygen produced within the thylakoid membranes both cause photosystem I photoinhibition. – *Plant Physiol.* **171**: 1626-1634, 2016.
- Tiwari A., Mamedov F., Grieco M. *et al.*: Photodamage of iron-sulphur clusters in photosystem I induces non-photochemical energy dissipation. – *Nat. Plants* **2**: 16035, 2016.
- Tripathi D.K., Singh S., Singh S. *et al.*: Nitric oxide alleviates silver nanoparticles (AgNps)-induced phytotoxicity in *Pisum sativum* seedlings. – *Plant Physiol. Biochem.* **110**: 167-177, 2017.
- Vass I.: Molecular mechanisms of photodamage in the Photosystem II complex. – *BBA-Bioenergetics* **1817**: 209-217, 2012.
- Vinković T., Novák O., Strnad M. *et al.*: Cytokinin response in pepper plants (*Capsicum annuum* L.) exposed to silver nanoparticles. – *Environ. Res.* **156**: 10-18, 2017.
- Wijnhoven S.W.P., Peijnenburg W.J.G.M., Herberts C.A. *et al.*: Nano-silver - A review of available data and knowledge gaps in human and environmental risk assessment. – *Nanotoxicology* **3**: 109-138, 2009.
- Yin L., Colman B.P., McGill B.M. *et al.*: Effects of silver nanoparticle exposure on germination and early growth of eleven wetland plants. – *PLoS ONE* **7**: e47674, 2012.
- Zhang R., Cruz J.A., Kramer D.M. *et al.*: Moderate heat stress reduces the pH component of the transthylakoid proton motive force in light-adapted, intact tobacco leaves. – *Plant Cell Environ.* **32**: 1538-1547, 2009.
- Zivcak M., Brestic M., Kunderlikova K. *et al.*: Repetitive light pulse-induced photoinhibition of photosystem I severely affects CO₂ assimilation and photoprotection in wheat leaves. – *Photosynth. Res.* **126**: 449-463, 2015.
- Zivcak M., Brückova K., Sytar O. *et al.*: Lettuce flavonoids screening and phenotyping by chlorophyll fluorescence excitation ratio. – *Planta* **245**: 1215-1229, 2017.
- Zuverza-Mena N., Armendariz R., Peralta-Videa J.R., Gardea-Torresdey J.L.: Effects of silver nanoparticles on radish sprouts: Root growth reduction and modifications in the nutritional value. – *Front Plant Sci.* **7**: 90, 2016.