

Characterization of Calcium Signals Provoked by Lysophosphatidylinositol in Human Microvascular Endothelial Cells

Y. M. AL SULEIMANI¹, C. R. HILEY²

¹Department of Pharmacology and Clinical Pharmacy, College of Medicine and Health Sciences, Sultan Qaboos University, Sultanate of Oman, ²Department of Pharmacology, University of Cambridge, Cambridge, United Kingdom

Received November 17, 2014

Accepted July 17, 2015

On-line November 24, 2015

Summary

The lipid molecule, lysophosphatidylinositol (LPI), is hypothesised to form part of a novel lipid signalling system that involves the G protein-coupled receptor GPR55 and distinct intracellular signalling cascades in endothelial cells. This work aimed to study the possible mechanisms involved in LPI-evoked cytosolic Ca^{2+} mobilization in human brain microvascular endothelial cells. Changes in intracellular Ca^{2+} concentrations were measured using cell population Ca^{2+} assay. LPI evoked biphasic elevation of intracellular calcium concentration, a rapid phase and a sustained phase. The rapid phase was attenuated by the inhibitor of PLC (U 73122), inhibitor of IP_3 receptors, 2-APB and the depletor of endoplasmic reticulum Ca^{2+} store, thapsigargin. The sustained phase, on the other hand, was enhanced by U 73122 and abolished by the RhoA kinase inhibitor, Y-27632. In conclusion, the Ca^{2+} signal evoked by LPI is characterised by a rapid phase of Ca^{2+} release from the endoplasmic reticulum, and requires activation of the PLC- IP_3 signalling pathway. The sustained phase mainly depends on RhoA kinase activation. LPI acts as novel lipid signalling molecule in endothelial cells, and elevation of cytosolic Ca^{2+} triggered by it may present an important intracellular message required in gene expression and controlling of vascular tone.

Key words

Lysophosphatidylinositol • GPR55 • Calcium • PLC • RhoA kinase

Corresponding author

Y. M. Al Suleimani, Department of Pharmacology and Clinical Pharmacy, College of Medicine and Health Sciences, Sultan Qaboos University, Sultanate of Oman. Fax: +96824141107. E-mail: yousufm@squ.edu.om

Introduction

There is increasing evidence that long-chain lipids have a wide variety of actions that are exerted through G protein-coupled receptors (Makide *et al.* 2014). The bioactive lipid, lysophosphatidylinositol (LPI) is now accepted as an endogenous ligand of G protein-coupled receptor GPR55. When LPI tested in systems expressing GPR55, it initiated responses, which were either absent in untransfected cells, diminished by siRNA knockdown of GPR55 or enhanced by GPR55 overexpression (Bondarenko *et al.* 2010, Ford *et al.* 2010, Henstridge *et al.* 2009, 2010, Oka *et al.* 2007, 2010, Waldeck-Weiermair *et al.* 2008). For instance, in HEK293 cells expressing a human GPR55 receptor, LPI induced Ca^{2+} mobilization (Henstridge *et al.* 2009), caused phosphorylation of extracellular signal-regulated kinases (Oka *et al.* 2007, 2009), and activated RhoA and nuclear factor of activated-T cells (Henstridge *et al.* 2009). Additionally, LPI initiated Ca^{2+} mobilization in mice dorsal root ganglion neurons (Lauckner *et al.* 2008) and human endothelial cells naturally expressing GPR55 (Bondarenko *et al.* 2010, Waldeck-Weiermair *et al.* 2008). In human umbilical vein endothelial cells (HUVEC), LPI induced both GPR55-dependent and -independent signals (Bondarenko *et al.* 2010, Waldeck-Weiermair *et al.* 2008), and acted as an intracellular messenger modulating Ca^{2+} -activated K^+ channels (Bondarenko *et al.* 2011a). The lipid also induced wound healing in primary human lung microvascular endothelial cells and causes platelets aggregation (Kargl *et al.* 2013).

With increasing evidence of vascular role of GPR55 and with lack of clarity of LPI action in vasculature, this study aims to further characterize the cellular action of LPI in endothelial cells. The intracellular Ca^{2+} signals in response to LPI in human brain microvascular endothelial cells were investigated.

Methods

Cell culture

The human immortalised brain endothelial cell line (hCMEC/D3) was provided by Dr Margery Barrand, Department of Pharmacology, University of Cambridge. The cell line was originally donated by Dr Pierre Couraud (INSERM, rue de Tolbiac, Paris, France) and line was established from isolated human microvascular brain endothelial cells (Weksler *et al.* 2005). The cells were grown on surfaces coated with 0.5 % bovine gelatine (Sigma), and were maintained in endothelial cell basal medium-2 (Lonza, Cologne, Germany) supplemented with 10 % (v/v) fetal bovine serum (FBS; Invitrogen) together with 100 U ml^{-1} penicillin, 100 $\mu\text{g ml}^{-1}$ streptomycin, 2 mM L-glutamine, 80 $\mu\text{g ml}^{-1}$ heparin, 5 $\mu\text{g ml}^{-1}$ ascorbic acid (all from Sigma), and 75 $\mu\text{g ml}^{-1}$ endothelial growth supplement (First Link, Birmingham). The cells were incubated at 37 °C in a humidified 5 % CO_2 atmosphere, and passaged when they reached ~90–95 % confluence. Experiments were conducted on cultures between passages 36 and 40.

Real-time PCR

Total RNA was extracted from the hCMEC/D3 cells using TRIZOL® (Invitrogen) as explained in the manufacturer's protocol. Briefly, the cells at ~90 % confluence in T75 flasks were homogenised in 1 ml TRIZOL® and total RNA was separated and precipitated and used to produce cDNA. Briefly, total RNA was subjected to DNase digestion before proceeding to cDNA synthesis using SuperScript™ II reverse transcriptase (Invitrogen) to synthesize the first strand cDNA. A quantitative polymerase chain reaction (qPCR) was carried out using the SensiMix™ SYBR & Fluorescein Kit (Bioline, London).

Population Ca^{2+} imaging

HCMEC/D3 cells were seeded onto 0.5 % gelatine-coated wells of 96-well black plates (Greiner Bio-One Ltd, Stonehouse, Gloucestershire), and used for

experiments 24 h later. Confluent cultures were first washed twice with phosphate-buffered saline (PBS, Invitrogen) and then loaded with 2 μM fluo-4 acetoxyethyl ester (Invitrogen). This was achieved by incubating the cells in HEPES-buffered saline (HBS; containing in mM: NaCl 135, KCl 5.9, MgCl_2 1.2, HEPES 11.6, glucose 11.5, CaCl_2 1.5, pH 7.3) supplemented with 2.5 mM probenecid (Sigma) and fluo-4 acetoxyethyl ester for 1 h at 20 °C. The cells were then washed with HBS and further incubated for 30 min to allow de-esterification of the dye before being used for experiments. All experiments were performed in HBS (or nominally Ca^{2+} -free HBS) at 20 °C. The plate containing fluo-4-loaded cells was mounted in a FlexStation III (MDS Analytical Technologies, Wokingham, Surrey, UK) to measure changes in fluorescence (excitation at 485 nm; emission at 525 nm). Drugs were prepared in another 96-well plate (Greiner Bio-One Ltd), and were automatically added to the loaded cells. The machine was programmed to allow for six simultaneous readings from each well at intervals of 1.52 s. At the end of each experiment, a solution containing 0.05 % Triton X-100 (Sigma) and 10 mM CaCl_2 was added to the cells in order to determine the fluorescence of Ca^{2+} -saturated indicator (F_{\max}). Background fluorescence (F_{\min}), where cells containing only Ca^{2+} -free indicator, was measured in parallel wells treated with a combination of 0.05 % Triton X-100 and 10 mM BAPTA (Molekula, Gillingham, Dorset, UK). This background was subtracted from all measurements. Data were recorded on SoftMax Pro version 5.4 (Govindan *et al.* 2010). Calcium Responses to LPI were recorded for a period of 5 min. Antagonists were incubated with cells for 10 min before being exposed to LPI.

Data and statistical analysis

In the population Ca^{2+} assay, both basal and peak cytosolic free Ca^{2+} concentrations were calculated using the formula:

$$[\text{Ca}^{2+}]_i = K_D F / (F_{\max} - F)$$

where K_D is the Ca^{2+} dissociation constant of fluo-4 (345 nM; Gee *et al.* 2000), F is the corrected fluorescence (for basal and peak Ca^{2+} levels) and F_{\max} is the fluorescence of the Ca^{2+} -saturated indicator. Data are expressed as $\Delta[\text{Ca}^{2+}]_i$, which indicates the difference in cytosolic Ca^{2+} concentration between basal and maximum release in response to a drug. In single cell Ca^{2+} imaging

experiments, data are expressed as Δ ratio (F_{340}/F_{380}) which indicates the difference in 340 to 380 nm fluorescence ratios between basal and peak Ca^{2+} . n represents number of cells from at least 2 independent isolations.

In experiments where mRNA expression was examined, the authenticity of each product was assessed by melting-curve analysis and the results were analysed using the “comparative quantitation” feature of the Rotor-Gene software. Concentration-response curves were analysed using 2-way ANOVA, and in those experiments where more than two groups were compared, statistical analysis was performed using one-way ANOVA followed by Bonferroni’s *post-hoc* test. A P value of less than 0.05 was taken as statistically significant. Data were analysed using GraphPad Prism (GraphPad, San Diego, CA, USA).

Drugs

L- α -lysophosphatidylinositol, histamine (all from the Sigma Chemical Company, Poole, Dorset, UK) and Y-27632 (Tocris Cookson, Bristol, UK) were dissolved in distilled water. 2-aminoethoxydiphenyl borate (2-APB; from Tocris) was dissolved in 100 % ethanol. U 73122 (Tocris), U 73343 (Sigma), and thapsigargin (Sigma) were dissolved in 100 % dimethyl sulfoxide (DMSO). All solutions were prepared on the day of the experiment.

Results

Expression profile of GPR55, CB₁ and CB₂ receptors in hCMEC/D3 cell line

Expression of GPR55 and the cannabinoid receptors CB₁ and CB₂ in the hCMEC/D3 cell line was examined using qPCR. Analyses of the melting curves revealed single specific peaks beyond 80 °C. Low levels of mRNA transcripts for GPR55, CB₁ and CB₂ receptors were detected in this cell line and all the receptors were approximately equally expressed when compared to the reference gene β -actin (Fig. 1).

Effect of histamine on [Ca²⁺]_i mobilization in hCMEC/D3 cells

In this study, the effect of histamine on cytosolic Ca^{2+} mobilization was first tested before proceeding to examine the actions of LPI. In endothelial cells, histamine acts on H₁-histamine receptors that are linked

to the G_{aq}-PLC-IP₃ pathway, resulting in elevation of $[\text{Ca}^{2+}]_i$ (Smit *et al.* 1999). In the population Ca^{2+} assay when extracellular Ca^{2+} was present, histamine (10 μM) provoked a rapid rise in $[\text{Ca}^{2+}]_i$ which was characterised by a peak (218±30 nM) followed by a sustained phase (Fig. 2A). This latter phase disappeared when Ca^{2+} -free buffer was used, indicating the need for Ca^{2+} entry (Fig. 2B). The downstream signalling pathway involved in the elevation of $[\text{Ca}^{2+}]_i$ was then investigated using enzyme inhibitors and receptor antagonists. As shown in Figure 2B and C, in nominally Ca^{2+} -free HBS, the rise in $[\text{Ca}^{2+}]_i$ in response to histamine was abolished by depletion of intracellular Ca^{2+} stores by thapsigargin (1 μM), inhibition of PLC by U 73122 (10 μM) and antagonism of IP₃ receptors by 2-APB (100 μM). The selective inhibitor of the RhoA-specific kinase p160ROCK, Y-27632 (50 μM) had no effect on the histamine response.

Characterization of LPI-evoked Ca^{2+} transients in hCMEC/D3 cells

Figure 3 shows that, using the population Ca^{2+} assay, LPI elicited a biphasic, concentration-dependent, elevation of $[\text{Ca}^{2+}]_i$ in hCMEC/D3 cells which was independent of extracellular Ca^{2+} . The initial phase was rapid (~8 s following LPI application) and transient, whereas the late phase was characterised by a slowly developing, sustained, elevation of $[\text{Ca}^{2+}]_i$ following the rapid phase, which was only observed at higher concentrations (above 10 μM).

Effects of thapsigargin, 2-APB, U 73122, U 73343 and Y-27632 on the early phase of the Ca^{2+} response

Since LPI induced a rise in $[\text{Ca}^{2+}]_i$ in which there were two peaks, the signalling pathway involved in each was studied independently. Pretreatment with thapsigargin (1 μM , 15 min incubation) attenuated the response to LPI (Fig. 4A) and the IP₃ receptor antagonist, 2-APB (100 μM , 10-min incubation) also abolished the response (Fig. 4B). The PLC inhibitor, U 73122 (10 μM , 10-min incubation) also abolished the rapid rise in $[\text{Ca}^{2+}]_i$ in response to LPI, whereas its inactive analogue, U 73343 (10 μM) had no effect (Fig. 5). Treating the cells with 50 μM Y-27632 (20-min incubation) resulted in a significant reduction in the elevation of $[\text{Ca}^{2+}]_i$ by LPI (Fig. 6).




Fig. 1. Expression of mRNA transcripts for GPR55, CB₁ and CB₂ receptors in the hCMEC/D3 endothelial cell line. **(A)** and **(B)** Representative real-time PCR measurements of the mRNA levels for the reference gene β -actin, GPR55, and the CB₁ and CB₂ receptors. Each curve is an average of duplicates from samples obtained from three different cell passages. The dashed lines represent samples where reverse-transcriptase (RT) was omitted, showing no specific amplification. **(C)** mRNA expression levels. Overexpression of human GPR55 in HEK293 cells (oxpGPR55) is shown as a positive control. Values are presented as mean expression relative to β -actin and vertical lines indicate standard error of mean (SEM).




Fig. 2. Ca^{2+} signals evoked by histamine in hCMEC/D3 endothelial cells. Typical results from populations of the cells stimulated with histamine either in Ca^{2+} -containing buffer **(A)** or in Ca^{2+} -free buffer in the presence of thapsigargin, U 73122, 2-APB or Y-27632 **(B)**. The results shown are means from three wells on a single plate and are typical of results from three independent plates from at least 3 different cell passages. **(C)** Histogram representing the peak $[Ca^{2+}]_i$ rise to histamine alone or in the presence of the enzyme inhibitors or the IP₃ receptor antagonist. *** represents statistical significance compared to the control group ($P < 0.001$). Values are means \pm SEM of triplicates obtained from 3 different cell passages.




Fig. 3. Ca^{2+} signals evoked by LPI in hCMEC/D3 endothelial cells. **(A)** Typical results from cell populations stimulated with 30 μM LPI in Ca^{2+} -free buffer, showing the biphasic $[\text{Ca}^{2+}]_i$ response. Results are means \pm SEM from three wells on a single plate. **(B)** Concentration-response relationship for phase 1 of the LPI-stimulated elevation of $[\text{Ca}^{2+}]_i$ in the presence or absence of extracellular Ca^{2+} . The effect of the vehicle (negative control) is also shown. **(C)** Effect of two concentrations of LPI on phase 2 of the elevation of the $[\text{Ca}^{2+}]_i$. Data from B and C are means \pm SEM of triplicates obtained from 3 different cell passages. **(D)** Recordings from populations of the cells each obtained from a single well showing the concentration-dependent effect of LPI on phase 1 of the elevation of $[\text{Ca}^{2+}]_i$. Each well of the 96-well plate was exposed to only a single concentration of LPI.





Fig. 4. Ca^{2+} signals evoked by LPI in hCMEC/D3 endothelial cells. Responses were initiated by LPI either alone or in the presence of thapsigargin **(A)** or 2-APB **(B)**. Data are means of triplicates obtained from 3 different cell passages and vertical lines represent SEM. $[\text{Ca}^{2+}]_i$ responses were obtained in nominally Ca^{2+} -free buffer. Each well of the 96-well plate was exposed to only a single concentration of LPI.



Effect of thapsigargin, 2-APB, U 73122 and Y-27632 on the sustained Ca^{2+} response to LPI

As mentioned above, the slowly developing, sustained, phase of the cytosolic Ca^{2+} response to LPI was observed at higher concentrations of the lysolipid. Interestingly, neither thapsigargin nor 2-APB, used at the same concentrations as above, had any noticeable effect on this phase in responses to 30 μM LPI (Fig. 7). Even more interestingly, exposing the cells to 10 μM U 73122 for 10 min produced a prominent concentration-dependent potentiation of the slow phase of Ca^{2+} signals evoked by LPI (Fig. 8). On the other hand, Y-27632 (50 μM) abolished the sustained phase (Fig. 9).





Fig. 7. LPI-evoked sustained elevation of $[\text{Ca}^{2+}]_i$ in hCMEC/D3 endothelial cells. Responses were initiated by LPI either alone or in the presence of thapsigargin (**A**) or 2-APB (**B**). Data are means of triplicates obtained from 3 different cell passages and vertical lines represent SEM. $[\text{Ca}^{2+}]_i$ responses were obtained in nominally Ca^{2+} -free buffer. Each well of the 96-well plate was exposed to only a single concentration of LPI.





Fig. 8. LPI-evoked sustained elevation of $[Ca^{2+}]_i$ in hCMEC/D3 endothelial cells. Responses were initiated by LPI alone or in the presence of U 73122 (**A**). Data are means of triplicates obtained from 3 different cell passages and vertical lines represent SEM. (**B**) Typical results from cell populations stimulated with LPI showing enhancement of the sustained elevation of $[Ca^{2+}]_i$ by U 73122. Results are means \pm SEM from three wells on a single plate. Ca^{2+} signals were recorded in nominally Ca^{2+} -free buffer. Each well of the 96-well plate was exposed to only a single concentration of LPI.



Discussion

In endothelial cells, recent studies have shown that LPI induces intracellular Ca^{2+} mobilization by activation of GPR55 (Bondarenko *et al.* 2010, Waldeck-Weiermair *et al.* 2008). The present study extends these observations to demonstrate that LPI initiates concentration-dependent cytosolic Ca^{2+} mobilization

in human brain microvascular endothelial cells (hCMEC/D3). The Ca^{2+} signal evoked by LPI is characterised by a rapid phase of Ca^{2+} release from the endoplasmic reticulum, and requires activation of the PLC-IP₃ signalling pathway and RhoA-dependent kinases. LPI also mediates a sustained phase of elevation of $[Ca^{2+}]_i$ which entirely depends on RhoA kinase activation.

In hCMEC/D3 endothelial cells, LPI, concentration-dependently, stimulated biphasic elevation of cytosolic Ca^{2+} . The rapid phase was characterised by a modest elevation of $[\text{Ca}^{2+}]_i$ (~40 nM at 30 μM LPI). Higher concentrations of LPI (>30 μM) were not tested due to the possibility of micelle formation beyond this concentration (Bondarenko *et al.* 2011b). Since there was no obvious maximal effect at the highest concentration used, the EC₅₀ of LPI could not be determined in this population assay. The LPI-initiated rapid elevation of $[\text{Ca}^{2+}]_i$ was due to release from intracellular stores, as thapsigargin, the inhibitor of SERCA, attenuated the response. This was further confirmed by abolition of the signal by antagonism of the IP₃ receptor using 2-APB, suggesting release from endoplasmic reticulum. Moreover, to test if PLC was involved in this process, the cells were treated with U 73122, an inhibitor of PLC, at a concentration previously reported to inhibit the enzyme (Bleasdale *et al.* 1990). Indeed, exposing the cells to U 73122 for 15 min diminished the Ca^{2+} signals to LPI, indicating a role for PLC. Thus, the rapid elevation of $[\text{Ca}^{2+}]_i$ is a receptor-initiated response, and involves activation of PLC-IP₃ pathway. In human endothelial cells, LPI, in addition to reported receptor-independent effects (including activation of nonselective cation channels, inhibition of Na⁺-K⁺ ATPase, and bidirectional modulation of large- and intermediate-conductance Ca^{2+} -activated K⁺ channels) (Bondarenko *et al.* 2010, 2011a,b), activates signalling through GPR55 (Bondarenko *et al.* 2010, Waldeck-Weiermair *et al.* 2008). There is an increasing body of evidence suggesting that GPR55 signals through G_{aq} linked to PLC-IP₃ and G_{α12/13} linked to RhoA. GPR55 is possibly involved in the rapid phase of $[\text{Ca}^{2+}]_i$ stimulated by LPI in hCMEC/D3 cells, as its mRNA transcript was detected in this cell line. In order to further investigate this, the involvement of RhoA-dependent kinases was examined. Indeed, a selective inhibitor of RhoA-specific kinase (p160ROCK), Y-27632 significantly reduced the response, pointing to a likely role of this kinase. RhoA-dependent Ca^{2+} signalling through GPR55 has recently been suggested (Henstridge *et al.* 2009). Overall, these findings suggest that LPI, by activation of GPR55, initiates Ca^{2+} transients through both PLC-IP₃ and RhoA pathways. Further studies will be required to determine if the response is mediated by dual signalling through both G_{aq} and G_{α13}.

As noted previously, LPI, in addition to inducing a rapid rise in $[\text{Ca}^{2+}]_i$, stimulated a sustained

elevation of cytosolic Ca^{2+} , although this was seen only at higher concentrations. Study of this phase was interesting. Since the response was observed in Ca^{2+} -free extracellular buffer, this may suggest an intracellular mechanism of Ca^{2+} release. However, depletion of Ca^{2+} stores by thapsigargin and antagonism of the IP₃ receptor by 2-APB did not inhibit the response, indicating Ca^{2+} stores other than those in the endoplasmic reticulum might be involved in the sustained elevation of $[\text{Ca}^{2+}]_i$ caused by LPI. The contribution of PLC to this response was also investigated with the commonly used inhibitor U 73122. Surprisingly, when the cells were pretreated with the inhibitor, the LPI-induced sustained elevation of cytosolic Ca^{2+} was greatly enhanced in a concentration-dependent manner, suggesting involvement of PLC. In fact, a very recent study reported that U 73122, in addition to inhibition of some isoforms of PLC, possesses stimulatory activity on others (Klein *et al.* 2011). It was concluded that, at 10 μM , it increased the activity of human PLC isoforms $\beta 2$, $\beta 3$ and $\gamma 1$, whereas the $\delta 1$ isoform was not affected. It was proposed that U 73122 binds covalently to PLC via cysteine residues leading to an increase in the affinity of the protein for cell membranes. One or more molecules of U 73122 may serve as lipid anchors for the modified PLC, allowing it to dock within the cell membrane in close proximity to substrate, thus leading to increased catalytic activity. Therefore, it is possible that LPI binding to a receptor (or by itself within the cell) activates one of the PLC isoforms that are activated by U 73122 leading to the observed enhanced elevation of cytosolic Ca^{2+} . As the LPI-initiated sustained elevation in $[\text{Ca}^{2+}]_i$ did not require activation of the IP₃ receptors, and was not sensitive to depletion of thapsigargin-sensitive stores, the role of PLC in cytosolic release of Ca^{2+} in this case remains unclear. It may suggest activation of other Ca^{2+} stores sensitive to PLC but not IP₃. One possible suggestion would be the mitochondria (De Marchi *et al.* 2014, Park *et al.* 2014), however, further studies will be required to confirm this. In the current study, inhibition of RhoA-dependent kinase by Y-27632 attenuated the sustained phase of the LPI-induced $[\text{Ca}^{2+}]_i$ release, suggesting a possible role for RhoA kinases. Whether this effect is a receptor-mediated or not will require further studies to determine. In a recent work on LPI and endothelial cells (Al Suleimani and Hiley 2015), similar findings were noticed in mesenteric endothelial cells. It was suggested that PLC ϵ (which is regulated by RhoA) is the possible target in this phase. Further studies by using specific inhibitors of this PLC

isoform will confirm this. Moreover, whether or not p160ROCK has direct regulatory activity on PLC, or indirectly through RhoA, is open to investigation. At least, this late phase of the Ca^{2+} response seems to require activation of both PLC and RhoA.

In the human umbilical vein endothelial cell line, EA.hy926, LPI induces GPR55-dependent and -independent elevation of cytosolic Ca^{2+} (Bondarenko *et al.* 2010). The receptor-dependent effect is sensitive to PLC inhibition while the receptor-independent effects involve activation of non-selective cation channels and inhibition of the Na^+/K^+ -ATPase. The current study agrees with these findings in that LPI produced a biphasic effect on cytosolic Ca^{2+} elevation that was characterised by an initial rapid peak, most probably mediated by GPR55, and a sustained phase which could be a GPR55-independent effect. In HEK293 cells expressing GPR55, LPI stimulates elevation of $[\text{Ca}^{2+}]_i$ that is oscillatory in nature (Henstridge *et al.* 2009). In our very recent published study (Al Suleimani and Hiley 2015) we have shown that LPI induced endothelium-dependent

vasorelaxation of rat resistance mesenteric arteries and that the effect was independent of activation of guanylyl cyclase and cyclooxygenase metabolites, but largely involved activation of Ca^{2+} -activated K^+ channels.

Overall, the experiments reported here on human brain microvascular endothelial cells show that LPI act as novel lipid signalling molecule, and elevation of cytosolic Ca^{2+} triggered by it may present an important intracellular message required in gene expression and controlling of vascular tone.

Conflict of Interest

There is no conflict of interest.

Acknowledgements

The authors would like to thank Professor Colin Taylor (Department of Pharmacology, University of Cambridge) and his laboratory members for their great help with calcium imaging. Great thanks also to the Sultan Qaboos University for the financial support.

References

- AL SULEIMANI YM, HILEY CR: The GPR55 agonist lysophosphatidylinositol relaxes rat mesenteric resistance artery and induces Ca^{2+} release in rat mesenteric artery endothelial cells. *Br J Pharmacol* **172**: 3043-3057, 2015.
- BLEASDALE JE, THAKUR NR, GREMBAN RS, BUNDY GL, FITZPATRICK FA, SMITH RJ, BUNTING S: Selective inhibition of receptor-coupled phospholipase C-dependent processes in human platelets and polymorphonuclear neutrophils. *J Pharmacol Exp Ther* **255**: 756-768, 1990.
- BONDARENKO A, WALDECK-WEIERMAIR M, NAGHDI S, POTESER M, MALLI R, GRAIER WF: GPR55-dependent and -independent ion signalling in response to lysophosphatidylinositol in endothelial cells. *Br J Pharmacol* **161**: 308-320, 2010.
- BONDARENKO AI, MALLI R, GRAIER WF: The GPR55 agonist lysophosphatidylinositol acts as an intracellular messenger and bidirectionally modulates Ca^{2+} -activated large-conductance K^+ channels in endothelial cells. *Pflugers Arch* **461**: 177-189, 2011a.
- BONDARENKO AI, MALLI R, GRAIER WF: The GPR55 agonist lysophosphatidylinositol directly activates intermediate-conductance Ca^{2+} -activated K^+ channels. *Pflugers Arch* **462**: 245-255, 2011b.
- DE MARCHI U, SANTO-DOMINGO J, CASTELBOU C, SEKLER I, WIEDERKEHR A, DEMAUREX N: NCLX protein, but not LETM1, mediates mitochondrial Ca^{2+} extrusion, thereby limiting Ca^{2+} -induced NAD(P)H production and modulating matrix redox state. *J Biol Chem* **289**: 20377-20385, 2014.
- FORD LA, ROELOFS AJ, ANAVI-GOFFER S, MOWAT L, SIMPSON DG, IRVING AJ, ROGERS MJ, RAJNICEK AM, ROSS RA: A role for L-alpha-lysophosphatidylinositol and GPR55 in the modulation of migration, orientation and polarization of human breast cancer cells. *Br J Pharmacol* **160**: 762-771, 2010.
- GEE KR, BROWN KA, CHEN WN, BISHOP-STEWART J, GRAY D, JOHNSON I: Chemical and physiological characterization of fluo-4 Ca^{2+} -indicator dyes. *Cell Calcium* **27**: 97-106, 2000.
- GOVINDAN S, TAYLOR EJ, TAYLOR CW: Ca^{2+} signalling by P2Y receptors in cultured rat aortic smooth muscle cells. *Br J Pharmacol* **160**: 1953-1962, 2010.

- HENSTRIDGE CM, BALENGA NA, FORD LA, ROSS RA, WALDHOER M, IRVING AJ: The GPR55 ligand L-alpha-lysophosphatidylinositol promotes RhoA-dependent Ca^{2+} signaling and NFAT activation. *FASEB J* **23**: 183-193, 2009.
- HENSTRIDGE CM, BALENGA NA, SCHRODER R, KARGL JK, PLATZER W, MARTINI L, ARTHUR S, PENMAN J, WHISTLER JL, KOSTENIS E, WALDHOER M, IRVING AJ: GPR55 ligands promote receptor coupling to multiple signalling pathways. *Br J Pharmacol* **160**: 604-614, 2010.
- KARGL J, BROWN AJ, ANDERSEN L, DORN G, SCHICHO R, WALDHOER M, HEINEMANN A: A selective antagonist reveals a potential role of G protein-coupled receptor 55 in platelet and endothelial cell function. *J Pharmacol Exp Ther* **346**: 54-66, 2013.
- KLEIN RR, BOURDON DM, COSTALES CL, WAGNER CD, WHITE WL, WILLIAMS JD, HICKS SN, SONDEK J, THAKKER DR: Direct activation of human phospholipase C by its well known inhibitor u73122. *J Biol Chem* **286**: 12407-12416, 2011.
- LAUCKNER JE, JENSEN JB, CHEN HY, LU HC, HILLE B, MACKIE K: GPR55 is a cannabinoid receptor that increases intracellular calcium and inhibits M current. *Proc Natl Acad Sci USA* **105**: 2699-2704, 2008.
- MAKIDE K, UWAMIZU A, SHINJO Y, ISHIGURO J, OKUTANI M, INOUE A, AOKI J: Novel lysophospholipid receptors; their structure and function. *J Lipid Res* **55**: 1986-1995, 2014.
- OKA S, NAKAJIMA K, YAMASHITA A, KISHIMOTO S, SUGIURA T: Identification of GPR55 as a lysophosphatidylinositol receptor. *Biochem Biophys Res Commun* **362**: 928-934, 2007.
- OKA S, TOSHIDA T, MARUYAMA K, NAKAJIMA K, YAMASHITA A, SUGIURA T: 2-Arachidonoyl-sn-glycero-3-phosphoinositol: a possible natural ligand for GPR55. *J Biochem* **145**: 13-20, 2009.
- OKA S, KIMURA S, TOSHIDA T, OTA R, YAMASHITA A, SUGIURA T: Lysophosphatidylinositol induces rapid phosphorylation of p38 mitogen-activated protein kinase and activating transcription factor 2 in HEK293 cells expressing GPR55 and IM-9 lymphoblastoid cells. *J Biochem* **147**: 671-678, 2010.
- PARK HK, LEE JE, LIM J, KANG BH: Mitochondrial Hsp90s suppress calcium-mediated stress signals propagating from mitochondria to the ER in cancer cells. *Mol Cancer* **13**: 148, 2014.
- SMIT MJ, HOFFMANN M, TIMMERMAN H, LEURS R: Molecular properties and signalling pathways of the histamine H1 receptor. *Clin Exp Allergy* **29** (Suppl 3): 19-28, 1999.
- WALDECK-WEIERMAIR M, ZORATTI C, OSIBOW K, BALENGA N, GOESSNITZER E, WALDHOER M, MALLI R, GRAIER WF: Integrin clustering enables anandamide-induced Ca^{2+} signaling in endothelial cells via GPR55 by protection against CB1-receptor-triggered repression. *J Cell Sci* **121**: 1704-1717, 2008.
- WEKSLER BB, SUBILEAU EA, PERRIERE N, CHARNEAU P, HOLLOWAY K, LEVEQUE M, TRICOIRE-LEIGNEL H, NICOTRA A, BOURDOULOUS S, TUROWSKI P, MALE DK, ROUX F, GREENWOOD J, ROMERO IA, COURAUD PO: Blood-brain barrier-specific properties of a human adult brain endothelial cell line. *FASEB J* **19**: 1872-1874, 2005.