

Adenosine A₁, A_{2a}, A_{2b}, and A₃ Receptors in Hematopoiesis. 2. Expression of Receptor mRNA in Resting and Lipopolysaccharide-Activated Mouse RAW 264.7 Macrophages

D. ŠTREITOVÁ¹, M. HOFER¹, J. HOLÁ¹, A. VACEK¹, M. POSPÍŠIL¹

¹Institute of Biophysics, v.v.i., Academy of Sciences of the Czech Republic, Brno, Czech Republic

Received November 24, 2008

Accepted January 22, 2009

On-line February 27, 2009

Summary

Expression of mRNA for adenosine receptor subtypes A₁, A_{2a}, A_{2b}, and A₃ in normal and lipopolysaccharide (LPS)-activated murine RAW 264.7 macrophages has been investigated using the method of quantitative real-time polymerase chain reaction. The results have shown a very low, unquantifiable expression of adenosine A₁ receptor mRNA in both normal and LPS-activated macrophages. The other three adenosine receptor mRNAs have been found to be expressed at various but always quantifiable levels. Activation of the macrophages by LPS induced upregulation of the expression of adenosine receptor A_{2a} and A_{2b} mRNA, whereas the expression of adenosine receptor A₃ mRNA was downregulated. Unstimulated macrophages exhibited a high expression of the A_{2b} adenosine receptor mRNA. The findings are discussed from the point of view of the antiinflammatory and hematopoiesis-stimulating roles of the adenosine receptor signaling.

Key words

Adenosine receptors • Macrophage • mRNA expression

Corresponding author

M. Hofer, Institute of Biophysics, v.v.i., Královopolská 135, CZ-612 65 Brno, Czech Republic. E-mail: hofer@ibp.cz

Adenosine, an ubiquitous purine nucleoside released into the extracellular environment from metabolically active or stressed cells, has been recognized to act as a paracrine regulator of a number of cellular functions (Poulsen and Quinn 1998, Schulte and Fredholm 2003). The regulatory role of extracellular

adenosine is based on the activation of specific receptors located on the cell surface. Functional and molecular studies made it possible to classify adenosine receptors as A₁, A_{2a}, A_{2b}, and A₃ subtypes. These G-protein-coupled receptors differ in their abilities to modulate various functions and their roles have been widely investigated in various areas of medicine (Olah and Stiles 1995).

An important regulatory system, which is believed to involve both macrophages and adenosine receptor signaling, is the modulation of hematopoietic processes. Hematopoiesis-modulating effects of adenosine receptor agonists have been reported (for review see Hofer and Pospíšil 2006). Activated macrophages have been found to produce hematopoiesis-stimulating factors, e.g. interleukin-6 (IL-6) or granulocyte colony-stimulating factor (G-CSF) (Hofer *et al.* 2007, Kamio *et al.* 2008). Administration of drugs increasing extracellular adenosine and strengthening its receptor-mediated action has been found to enhance the ability of serum of the treated mice to support the proliferation of granulocyte/macrophage progenitor cells (Weiterová *et al.* 2007). It may be hypothesized from these observations that the actions of adenosine receptor agonists on hematopoietic cells are indirect, mediated by other cells and that macrophages might play a role in this phenomenon.

A considerable attention has also been paid to the participation of adenosine receptor signaling in immunological and inflammatory processes. Here, the interest was focused on the functions of macrophages and to their modulation by the adenosine receptor system, i.e.

on the expression of individual adenosine receptors by these cells, as well as on their role in mechanisms of inflammation. The accumulated evidence indicates that it is especially the activation of adenosine A_{2a} receptor subtype which downregulates the inflammation and mediates protection from tissue damage (Ohta and Sitkovsky 2001, Haskó and Pacher 2008). An increased expression of mRNA for adenosine A_{2a} receptor in macrophages under inflammatory states has also been reported (Khoa *et al.* 2001, Bshesh *et al.* 2002) and confirmed recently in experiments of Murphree *et al.* (2005) on mouse peritoneal macrophages, human macrophages, and Wehi-3 cells.

Experiments, the results of which are presented here, were focused on the investigations of levels of adenosine receptor mRNAs of all four adenosine receptor subtypes in resting and LPS-activated mouse RAW 264.7 macrophages. They comprise determinations of mRNA expression in three time intervals up to 24 h following the initiation of macrophage activation by LPS as well as statistical processing of the data obtained.

Murine macrophage cell line RAW 264.7, purchased from American Type Culture Collection (ATCC, Manassas, VA, USA) was maintained in D-MEM medium supplemented with 10 % fetal heat-inactivated bovine serum. For experiments 1×10^6 cell/well (1.0 ml) were seeded onto 24-well plates. The cells were allowed to adhere for one hour at 37 °C, non-adherent cells were removed by aspiration and wells were filled with 1.0 ml of D-MEM medium with 10 % fetal bovine serum. The cells were cultivated in a thermostat (Forma Scientific, U.S.A.) at 37 °C in a fully humidified atmosphere containing 5 % CO_2 in air. The density of the cells in the cultures during the experiments was approximately 1×10^6 cells per ml.

Stimulation of RAW 264.7 cells with LPS: One μ g LPS (Sigma, St. Louis, MO, USA) in a volume of 100 μ l was added to the cell culture to obtain the final concentration of 0.1 μ g/ml. Corresponding volume of medium (saline) was added to control cell cultures.

RNA isolation from the RAW 264.7 murine macrophages was done using a kit with Dnase treatment (RNeasy Mini Kit, QIAGEN, Hilden, Germany). RNA concentration and its purity were quantified by UV spectrophotometry. The 260/280 ratio was not less than 1.8 for each RNA sample.

Quantitative real-time RT-PCR: cDNA was prepared using iScript cDNA Synthesis Kit (Bio-Rad, CA, USA). 1 μ g of RNA template was used per each




Fig. 1. Adenosine A_{2a} , A_{2b} , and A_3 receptor mRNA expression in control untreated RAW 264.7 macrophages. Relative values of mRNA expression are given as means \pm SD.

reverse transcription. The mRNA levels of selected genes in RNA from mouse RAW 264.7 macrophages were measured by real-time reverse-transcription-polymerase chain reaction (RT-PCR) in triplicates on RotorGene 6000 (Corbett Research, Sydney, Australia) using SYBR Green Master Mix reagent (Roche Diagnostics, Mannheim, Germany). The final reaction volume (20 μ l) included 10 μ l SYBR Green Master Mix reagent, 5 μ l of diluted cDNA, and 1000 nM concentration of each primer. The first reaction cycle was carried out at 95 °C for 10 min and was followed by 40 cycles, each of which consisted in 15 s at 95 °C, 20 s at 60 °C and 20 s at 72 °C. Sequences of primers were taken over from Ashton *et al.* (2003) and Overbergh *et al.* (1999) (the sequences are shown in Štreitová *et al.* 2010). To calculate relative gene expression, we used delta-delta Ct method based on the difference of threshold cycles (Ct) of the target gene and the β -actin sequence. We assumed a twofold increase in PCR products per cycle. A receptor was taken for relative quantification if the threshold cycle number was less than 36. If the threshold cycle number was greater than 36, the receptor was considered to be present in minimal quantities and relative quantification was not performed. A gene was considered not to be expressed if no amplification was detected by cycle 40.

Statistics: all experiments were performed three times. The results are expressed as means \pm SD. For statistical evaluation of the differences between control and LPS-activated macrophages, ANOVA and Tukey-HSD test were used. The significance level was set at $P < 0.05$.

In initial experiments, relative mRNA expression for A_1 , A_{2a} , A_{2b} , and A_3 receptors was evaluated in control




Fig. 2. Effects of LPS-induced activation on the expression of adenosine A_{2a}, A_{2b}, and A₃ receptor mRNA in RAW 264.7 macrophages. Data are expressed as relative changes compared to controls (control untreated cells, value = 1) and given as means \pm SD. Time interval (hours) – time interval between the beginning of induction of activation and material sampling. *, ** - $P < 0.05$, $P < 0.01$, respectively, in comparison with control; + - $P < 0.05$ in comparison with 24 h.

untreated RAW 264.7 macrophages. The threshold cycle number for A₁ receptor mRNA occurred in the interval between 36 and 40. Therefore, the A₁ receptor mRNA was considered to be present in minimal quantities not suitable for quantification (see above). The values for the relative mRNA expression for A_{2a}, A_{2b}, and A₃ receptors are shown in Figure 1. The relative mRNA level for A_{2a} receptors was

rather low, though quantifiable ($1.06 \pm 0.47 \times 10^{-6}$), those for A_{2b} and A₃ receptors were markedly higher ($5.80 \pm 1.41 \times 10^{-4}$ and $1.73 \pm 0.28 \times 10^{-4}$, respectively).

The relative changes of the above mRNA levels induced by LPS-activation represented the subject of consequential studies. The level of mRNA expression for A₁ receptor did not attain values suitable for quantification of the experiments. Results on A_{2a}, A_{2b}, and A₃ mRNA expression are shown in Figure 2. It follows from the results that an up-regulation was induced in the expression of mRNA of adenosine A_{2a} and A_{2b} receptor, whereas a down-regulation occurred in the expression of mRNA of the adenosine A₃ receptor. A particularly expressive increase was found in the A_{2a} mRNA expression, namely to more than 120 times of that found in the controls at 3 and 6 h after the beginning of the stimulation with LPS. It was followed by a decrease to values not significantly different in comparison with control values. The elevation of the level of the A_{2b} receptor mRNA expression attained values approximately 8 times higher than those of the controls at 3 and 6 h and returned to nearly control levels after 24 h. The A₃ adenosine receptor mRNA levels significantly decreased to about one half of those in the controls at 3 h after the beginning of the stimulation with a subsequent gradual return to control levels after 24 h.

Our results on the induction of changes in the expression of mRNA for adenosine A_{2a}, A_{2b}, and A₃ receptors following stimulation of RAW 264.7 murine macrophages by LPS confirm that the action of these receptors can be included into the spectrum of the macrophage activities initiated by their LPS-induced activation. Such a definite statement cannot be expressed for adenosine A₁ receptors since their mRNA has been found to be present only in minimal levels not enabling quantification. Adenosine receptor agonists have been shown to modulate production of many cytokines and other regulatory factors by macrophages, e.g. interleukin-10 (Haskó *et al.* 1996, Le Moine *et al.* 1996), tumor necrosis factor- α (TNF- α) (Haskó *et al.* 1996, 2000), interleukin-12 (Haskó *et al.* 2000), or prostaglandin E₂ (Ezeamuzie and Khan 2007). As mentioned, earlier studies have shown that the activation of macrophages by LPS induces a production of proinflammatory (TNF- α) (Franco-Molina *et al.* 2005) and hematopoiesis-stimulating (IL-6, G-CSF) (Hofer *et al.* 2007, Kamio *et al.* 2008) cytokines by these cells. Thus, LPS-mediated macrophage activation has been shown to change the profile of active substances released from macrophages towards an up-regulation of immune reactions and production of blood cells.

The data from our experiments have shown that LPS induces an up-regulation of the expression of adenosine A_{2a} and A_{2b} receptor mRNA and a down-regulation of the expression of adenosine A_3 receptor mRNA in RAW 264.7 macrophages. These effects are similar to those observed by Murphree *et al.* (2005) who used other cell lines of macrophages. As stated by these authors, the up-regulation of the expression of adenosine A_{2a} and A_{2b} receptor mRNA can be understood as a feedback mechanism aimed at macrophage deactivation. This is logical in terms of the evidence on antiinflammatory action of both these receptors deduced from studies using mice deficient in A_{2a} (Ohta and Sitkovsky 2001) and A_{2b} receptors (Yang *et al.* 2006). The reason for down-regulated expression of adenosine A_3 receptor mRNA after LPS-induced activation of macrophages observed by Murphree *et al.* (2005), as well as in our study, is unknown and remains to be elucidated.

In context of the considered role of adenosine receptor signaling in regulation of hematopoietic processes by macrophages it should be mentioned that a positive regulatory role in stimulation of proliferation of hematopoietic progenitor cells has been reported for an agonist of the adenosine A_3 receptors whereas suppression of their proliferation for an agonist of the adenosine A_{2a} receptors (Pospíšil *et al.* 2004). Since activation of the RAW 264.7 macrophages has been described in the present communication to down-regulate the expression of the A_3 receptors and to up-regulate that of the A_{2a} receptors, it seems that activation of the macrophages does not increase the ability of these cells to respond to adenosine receptor agonists in a hematopoiesis-stimulating manner. Further studies, including those on primary mammalian macrophages, are needed to supply more data on this topic and to elucidate the biological meaning of the observed phenomena.

As shown, the changes in the expression of mRNA of adenosine receptors induced by LPS are temporary. Maximum of the changes can be observed at 3 and 6 h after activation of the macrophage by LPS, whereas the level of mRNA expression returns back to control values at 24 h (Fig. 2). However, maximal manifestations of functional changes in macrophages after their activation with LPS, like an increase in production of nitric oxide, prostaglandin E_2 or granulocyte colony-stimulating factor, are recorded at time interval between 18 to 24 h (Hulkower *et al.* 1996, Hofer *et al.* 2007). Our data document well that the changes in mRNA expression are primary and precede modulation of the functional state of

the macrophages.

Besides the changes in mRNA expression of the adenosine receptors induced by LPS, the differences in mRNA expression of individual subtypes of adenosine receptors in resting conditions of the cell system can also provide an interesting information. Assuming that the adenosine receptor signaling presents the retaliatory mechanism (Newby 1984), the higher expression of some of the adenosine receptor mRNAs under resting conditions can indicate a higher readiness of the cells to activate such regulatory mechanism under stress conditions. As shown by our experiments, such an emergency role in the activated macrophage system can be played by the adenosine A_{2b} receptor whose mRNA has been found to exhibit a relatively high level of expression. It is interesting that Németh *et al.* (2005) detected A_{2b} receptors in membrane protein fractions from RAW 264.7 cells.

The level of mRNA does not always correlate with levels of respective proteins. In our experiments protein levels could not be determined since antibodies against mouse adenosine receptor proteins, with the exception of those against A_{2a} , are not available. However, the convincing data on the mRNA expression for the adenosine A_{2a} , A_{2b} , and A_3 receptors in control as well as LPS-activated RAW 264.7 mouse macrophages enable us to express an opinion that adenosine receptor signaling plays a role in macrophage-mediated regulatory processes. Based on the results shown here taken together with those obtained in studies on mouse primary hematopoietic precursor cells (Štreitová *et al.* 2010), it is possible to conclude that cells of the hematopoietic system express, though in various levels, mRNA for all four subtypes of adenosine receptors and are, thus, able to interact with both non-selective and selective adenosine receptor agonists. Nevertheless, more data are needed for specifying the regulatory pathways through which the effects of adenosine receptor agonists on the hematopoietic system are exerted.

Conflict of Interest

There is no conflict of interest.

Acknowledgements

This work was supported by the Academy of Sciences of the Czech Republic (grants AV0Z50040507 and AV0Z50040702), and by the Grant Agency of the Czech Republic (grants 305/06/0015 and 305/08/0158). The authors thank dr. Vladimír Znojil for cooperation in preparation of the manuscript.

References

- ASHTON KJ, NILSSON U, WILLEMS L, HOLMGREN K, HEADRICK JP: Effects of aging and ischemia on adenosine receptor transcription in mouse myocardium. *Biochem Biophys Res Commun* **312**: 367-372, 2003.
- BSHESH K, ZHAO B, SPIGHT D, BIAGGIONI I, FEOKTISTOV I, DENENBERG A, WONG HR, SHANLEY TP: The A_{2A} receptor mediates an endogenous regulatory pathway of cytokine expression in THP-1 cells. *J Leukoc Biol* **72**: 1027-1036, 2002.
- EZEAMUZIE CI, KHAN I: The role of adenosine A₂ receptors in the regulation of TNF- α production and PGE₂ release in mouse peritoneal macrophages. *Int Immunopharmacol* **7**: 483-490, 2007.
- FRANCO-MOLINA MA, MENDOZA-GAMBOA E, CASTILLO-LEÓN I, TAMEZ-GUERRA RS, RODRÍGUEZ-PADILLA C: Bovine dialyzable leukocyte extract modulates the nitric oxide and pro-inflammatory cytokine production in lipopolysaccharide-stimulated murine peritoneal macrophages in vitro. *J Med Food* **8**: 20-26, 2005.
- HASKÓ G, KUHEL DG, CHEN JF, SCHWARZSCHILD MA, DEITCH EA, MABLEY JG, MARTON A, SZABÓ C: Adenosine inhibits IL-12 and TNF- α production via adenosine A_{2a} receptor-dependent and independent mechanisms. *FASEB J* **14**: 2065-2074, 2000.
- HASKÓ G, PACHER P: A(2A) receptors in inflammation and injury: lessons learned from transgenic animals. *J Leukoc Biol* **83**: 447-455, 2008.
- HASKÓ G, SZABÓ C, NÉMETH ZH, KVETAN V, PASTORES SM, VIZI ES: Adenosine receptor agonists differentially regulate IL-10, TNF-alpha, and nitric oxide production in RAW 264.7 macrophages and in endotoxemic mice. *J Immunol* **157**: 4634-4640, 1996.
- HOFER M, POSPÍŠIL M: Role of adenosine signaling in hematopoiesis – a short review. *Med Hypotheses Res* **3**: 629-635, 2006.
- HOFER M, VACEK A, LOJEK A, HOLÁ J, ŠTREITOVÁ D: Ultrafiltered pig leukocyte extract (IMUNOR®) decreases nitric oxide formation and hematopoiesis-stimulating cytokine production in lipopolysaccharide-stimulated RAW 264.7 macrophages. *Int Immunopharmacol* **7**: 1369-1374, 2007.
- HULKOWER KI, POLLOCK JS, WALSH RE, HUANG R, OTIS ER, BROOKS CDW, BELL RL: Leukotrienes do not regulate nitric oxide production in RAW 264.7 macrophages. *Prostaglandins Leukot Essent Fatty Acids* **55**: 145-149, 1996.
- KAMIO N, AKIFUSA S, YAMAGUCHI N, YAMASHITA Y: Induction of granulocyte colony-stimulating factor by globular adiponectin via the MEK-ERK pathway. *Mol Cell Endocrinol* **292**: 20-25, 2008.
- KHOA ND, MONTESINOS MC, REISS AB, DELANO D, AWADALAH N, CRONSTEIN BN: Inflammatory cytokines regulate function and expression of adenosine A_{2A} receptors in human monocytic THP-1 cells. *J Immunol* **167**: 4026-4032, 2001.
- LE MOINE O, STORDEUR P, SCHANDENE L, MARCHANT A, DE GROOTE D, GOLDMAN M, DEVIERE J: Adenosine enhances IL-10 secretion by human monocytes. *J Immunol* **156**: 4408-4414, 1996.
- MURPHREE LJ, SULLIVAN GW, MARSHALL MA, LINDEN J: Lipopolysaccharide rapidly modifies adenosine receptor transcripts in murine and human macrophages: role of NF- κ B in A_{2A} adenosine receptor induction. *Biochem J* **391**: 575-580, 2005.
- NÉMETH ZH, LUTZ CS, CSÓKA B, DEITCH EA, LEIBOVICH SJ, GAUSE WC, TONE M, PACHER P, VIZI ES, HASKÓ G: Adenosine augments IL-10 production by macrophages through an A_{2B} receptor-mediated posttranscriptional mechanism. *J Immunol* **175**: 8260-8270, 2005.
- NEWBY AC: Adenosine and the concept of “retaliatory” metabolite. *Trends Biochem Sci* **9**: 42-44, 1984.
- OHTA A, SITKOVSKY M: Role of G-protein-coupled adenosine receptors in downregulation of inflammation and protection from tissue damage. *Nature* **414**: 916-920, 2001.
- OLAH ME, STILES GL: Adenosine receptor subtypes: characterization and therapeutic regulation. *Annu Rev Pharmacol Toxicol* **35**: 581-606, 1995.
- OVERBERGH L, VALCKX D, WAER M, MATHIEU C: Quantification of murine cytokine mRNAs using real time quantitative reverse transcriptase PCR. *Cytokine* **11**: 305-312, 1999.

-
- POSPÍŠIL M, HOFER M, VACEK A, ZNOJIL V, PIPALOVÁ I: Effects of stable adenosine receptor agonists on bone marrow haematopoietic cells as inferred from the cytotoxic action of 5-fluorouracil. *Physiol Res* **53**: 549-556, 2004.
- POULSEN S-A, QUINN RJ: Adenosine receptors: new opportunities for future drugs. *Bioorg Med Chem* **6**: 619-641, 1998.
- SCHULTE G, FREDHOLM BB. Signalling from adenosine receptors to mitogen-activated protein kinases. *Cell Signal* **15**: 813-827, 2003.
- ŠTREITOVÁ D, ŠEFC L, SAVVULIDI F, POSPÍŠIL M, HOLÁ J, HOFER M: Adenosine A₁, A_{2a}, A_{2b}, and A₃ receptors in hematopoiesis. 1. Expression of receptor mRNA in four mouse hematopoietic precursor cells. *Physiol Res* **59**: 133-137, 2010.
- WEITEROVÁ L, HOFER M, POSPÍŠIL M, ZNOJIL V, ŠTREITOVÁ D: Drugs elevating extracellular adenosine administered in vivo induce serum colony-stimulating activity and interleukin-6 in mice. *Physiol Res* **56**: 463-473, 2007.
- YANG D, ZHANG Y, NGUYEN HG, KOUPENOVA M, CHAUHAN AK, MAKITALO M, JONES MR, HILAIRE CS, SELDIN DC, TOSELLI P, LARNPERTI E, SCHREIBER BM, GAVRAS H, WAGNER DD, RAVID K: The A_{2B} adenosine receptor protects against inflammation and excessive vascular adhesion. *J Clin Invest* **116**: 1913-1923, 2006.
-