

Hydrogen Sulfide Improves the Endothelial Dysfunction in Renovascular Hypertensive Rats

H. XUE^{1*}, S. ZHOU^{1*}, L. XIAO¹, Q. GUO¹, S. LIU¹, Y. WU¹

*These authors contributed equally to this work.

¹Department of Physiology, Hebei Medical University, Shijiazhuang, Hebei, China

Received June 12, 2014

Accepted January 16, 2015

On-line March 24, 2015

Summary

As a novel gasotransmitter, hydrogen sulfide (H₂S) has vasodilating and antihypertensive effects in cardiovascular system. Thus, we hypothesized that H₂S might have beneficial effects on thoracic endothelial function in two-kidney one-clip (2K1C) rats, a model of renovascular hypertension. Sodium hydrosulfide (NaHS, 56 μmol/kg/day) was administrated intraperitoneally from the third day after the 2K1C operation. Along with the development of hypertension, the systolic blood pressure (SBP) was measured before the operation and each week thereafter. The oxidative stress was determined by measurement of malondialdehyde (MDA) concentration, superoxide dismutase (SOD) activity and protein expression of oxidative stress-related proteins (AT₁R, NADPH oxidase subunits). Acetylcholine (ACh)-induced vasorelaxation and angiotensin II (Ang II)-induced vasoconstriction were performed on isolated thoracic aorta. The SBP was significantly increased from the first week after operation, and was lowered by NaHS. NaHS supplementation ameliorated endothelial dysfunction. The protein expression of oxidative stress-related proteins were downregulated, while SOD activity upregulated. In conclusion, improvement of endothelial function is involved in the antihypertensive mechanism of H₂S. The protective effect of H₂S is attributable to suppression of vascular oxidative stress that involves inhibition of Ang II-AT₁R action, downregulation of oxidases, as well as upregulation of antioxidant enzyme.

Key words

Hydrogen sulfide • Angiotensin II • Endothelial dysfunction • Oxidative stress

Corresponding author

Y. Wu, Department of Physiology, Hebei Medical University,

361 East Zhongshan Road, Shijiazhuang, Hebei 050017, China.

Fax: +86-311-86266407. E-mail: wuyum@yahoo.com

Introduction

The novel gasotransmitter hydrogen sulfide (H₂S) possessed various physiological and pharmacological functions in cardiovascular system (Wang 2002). Nowadays, H₂S has been found to be produced by cystathionine β-synthase (CBS), cystathionine γ-lyase (CSE) and newly determined 3-mercaptopyruvate sulfurtransferase (3-MST) in concert with cysteine aminotransferase (CAT) (Tanizawa 2011). In particular, CSE is the main generating enzyme to be present in vascular bed and additionally located in vascular endothelial cells, which contributes to endothelium-dependent relaxation (Zhao *et al.* 2001). The vasorelaxing effect of H₂S has been appeared in various vascular tissues, such as rat aorta and mesenteric arteries (Wang *et al.* 2009). Moreover, CSE knockdown increased blood pressure (Yang *et al.* 2008), which might be eliminated by *in vivo* administration of H₂S (Mancardi *et al.* 2009). In addition, plasma level of H₂S was significantly decreased in spontaneously hypertensive rats (SHRs) (Zhao *et al.* 2008) and renovascular hypertensive rats (Lu *et al.* 2010). Thus, we hypothesized that there might be a correlation between H₂S and vascular function in hypertensive animal model.

Renovascular hypertension caused by partial renal-artery stenosis always leads to persistent high blood pressure and increased angiotensin II (Ang II) activity (Higashi *et al.* 2002, Takahashi and Smithies 2004). Ang II, acting *via* the angiotensin II type 1 receptor

(AT₁R), increases the activity of nicotinamide adenine dinucleotide phosphate (NADPH) oxidase, which was proposed as a mechanism underlying the enhanced production of reactive oxygen species (ROS), especially superoxide anion (O₂⁻) in a rat model of renovascular hypertension (Oliveira-Sales *et al.* 2009). The association between increased production of O₂⁻ and endothelial dysfunction has been reported in both experimental and human renovascular hypertension (Cai and Harrison 2000, Dijkhorst-Oei *et al.* 1999, Taddei *et al.* 1993).

Vasoactive substances released by endothelium play an important role in regulating vascular tone and blood pressure (Bayraktutan 2002). The imbalance between endothelium-derived relaxing and contracting factors leads to endothelial dysfunction, which is related to cardiovascular mortality (Vanhouste *et al.* 2009). In a large amount of experimental hypertensive models, such as two-kidney, one-clip (2K1C) renovascular, aortic coarctation, Dahl salt-sensitive, deoxycorticosterone acetate-salt and SHR, the endothelium-dependent vasorelaxation is severely impaired (Choi *et al.* 2012). H₂S has been reported to be produced from vascular endothelial cells and be considered as a newly endothelial-dependent relaxing factor (EDRF) or endothelial-dependent hyperpolarized factor (EDHF), which plays a pivotal role in regulating endothelial function (Wang 2009). However, no studies show that H₂S could improve endothelial dysfunction in 2K1C model.

Sodium hydrosulfide (NaHS) is an H₂S donor. Although previous studies reported the antihypertensive effect of this compound (Ahmad *et al.* 2014, Ford *et al.* 2013), the effects of NaHS on vascular function in renovascular hypertension remain poorly understood. In this study, we investigated the effect of NaHS on endothelial function in chronic 2K1C hypertensive rats, which is an ideal animal model of renovascular hypertension, with further elucidation of the underlying mechanisms by focusing on vascular oxidative stress.

Methods

Animals

All the rats used in present study were obtained from our local certified animal facility (Animal Research Center of Hebei Medical University, Hebei, China). 7-week old male Sprague-Dawley rats were randomly divided into 3 groups (n=6): Sham, 2K1C, 2K1C+NaHS. The rats were anesthetized with intraperitoneal injections

of pentobarbital sodium (30 mg/kg). In the 2K1C and 2K1C+NaHS groups, a lumbar incision was made to provide access to the left renal artery. Cleared of the connective tissues, the left renal artery was clipped by a rigid U-shaped solid silver clip with an open slit of 0.25 mm, resulting in partial occlusion of renal perfusion. The contralateral kidney was left untouched. Sham group underwent the same procedure, except for the clip placement. The rats were kept in cages after surgery, maintained on 12 h light/dark cycle with free access to normal chow and tap water. 2K1C+NaHS group received NaHS 56 μmol/kg/day intraperitoneally. The treatment was started from the third day after the surgery and maintained for 4 weeks. Sham and 2K1C groups were received saline as vehicle.

Blood pressure measurement

Systolic blood pressure (SBP) was measured non-invasively by tail-cuff plethysmography (Chengdu Instrument Factory, Sichuan, China) in calm, conscious rats. Briefly, SBP was measured before and further at each week after surgery for 4 weeks. SBP measurement was always conducted between 9:00 and 12:00 AM and was calculated as the average value of 3 to 4 successive measurement.

Isometric force studies

The thoracic aorta was dissected, cleaned of connective tissue and cut into ring segments (-3 mm length). The segments were then mounted between stainless steel triangles in a 10 ml organ bath filled with oxygenated (95 % O₂ and 5 % CO₂) Krebs-Henseleit solution (pH 7.3-7.5) (composition in mmol/l: NaCl 119, KCl 4.7, CaCl₂ 2.5, MgCl₂ 1, NaHCO₃ 25, KH₂PO₄ 1.2, glucose 11) at 37 °C. An optional baseline tone of 2 g was applied to all rings as a resting tension. After a 60 min equilibration, arteries were stimulated by 60 mmol/l KCl. After washing out, rings were pre-contracted by phenylephrine (Phe, 1 μmol/l). When the contraction was steady, acetylcholine (ACh, 10⁻⁸~10⁻⁴ mol/l) or sodium nitroprusside (SNP, 10⁻⁹~10⁻⁶ mol/l) were added respectively in a cumulative manner to record the endothelial-dependent relaxation or endothelial-independent relaxation response curve. Ang II (10⁻¹⁰~10⁻⁶ mol/l) was added cumulatively to conduct contraction curve. Further, AT₁R inhibitor losartan (10 μmol/l) was applied for 30 min before the Ang II-induced contraction was performed.

The extent of relaxation was expressed as

percentage of relaxation from a submaximal phenylephrine-induced constriction. Vasoconstriction was determined in relative values as the percentage of 60 mmol/l KCl contraction.

Measurement of plasma H₂S level

H₂S concentration in plasma was measured as previously described (Siegel 1965). Briefly, plasma (100 µl) was mixed with potassium phosphate buffer (pH 7.4, 2.5 ml) and zinc acetate (1 % w/v, 500 µl), then further incubated with *N,N*-dimethyl-*p*-phenylenediamine sulfate (20 mmol/l, 500 µl) in 7.2 mol/l HCl and FeCl₃ (30 mmol/l, 400 µl) in 1.2 mol/l HCl. 10 % trichloroacetic acid 1 ml was added to terminate the reaction after 20-min incubation. The absorbance of the mixture was measured at 665 nm. H₂S concentration was calculated against a calibration curve of NaHS (0.01 to 100 µmol/l). The results of H₂S concentration was expressed as µmol/l.

Measurement of plasma Ang II level

Ang II level in plasma was examined with an Ang II radioimmunoassay kit (Beijing Chemclin Biotech Co. Ltd, Beijing, China). Briefly, blood samples were collected in tubes containing 20 µl EDTA in ice. After centrifugation at 1000×g for 15 min at 4 °C, plasma (1 ml) was collected and mixed with enzyme inhibitor 1 (10 µl), enzyme inhibitor 2 (20 µl) following the instruction of the radioimmunoassay kit. Then aliquots (100 µl) were incubated with ¹²⁵I-AII (100 µl) and AII antibody (100 µl) for 15 h at 2-8 °C, followed by addition of separating reagent (including donkey anti-rabbit IgG serum and polyethylene glycol) 500 µl and incubated for 15 min at room temperature. The supernatant was discarded after centrifugation at 3,500 r/min for 20 min. The radioactive count was determined for 60 s. The results were expressed as pg/ml.

Western blot analysis

Thoracic aorta were homogenized with ice-cold RIPA lysing buffer and centrifuged at 20,000×g for 20 min. The supernatant was collected and analyzed for protein concentration using the bicinchoninic acid (BCA) method (Generay Biotechnology, Shanghai, China). Aliquots (100 µg protein) were suspended in six volumes of sample buffer with 5 % β-mercaptoethanol and denatured at 99 °C for 10 min. The protein samples were electrophoresed through 10 % SDS-polyacrylamide gel and then transferred onto an immobilon-P polyvinylidene

difluoride membrane using wet transfer at 100 V for 90 min at 4 °C. The membrane was blocked with 0.05 % Tween-20 Tris-buffered saline (TBST) containing 5 % non-fat milk for 60 min at room temperature. The blots were then incubated overnight at 4 °C with anti-AT₁R (1:800, Abcam, Hongkong), anti-Nox2 (1:1000, Abcam, Hongkong), anti-Nox4 (1:1000, Proteintech, Chicago, USA), anti-p67^{phox} (1:1000, EPITOMICS, California, USA) and anti-superoxide dismutase (SOD)-1 (1:1000, Proteintech, Chicago, USA) antibodies. All washes were performed in TBST. The membranes were then incubated with a horseradish peroxidase (HRP)-conjugated goat anti-rabbit antibody (1:2000, Proteintech, Chicago, USA) or HRP-conjugated rabbit anti-goat antibody (1:2000, Proteintech, Chicago, USA) for 1 h. Blots were developed with an enhanced chemiluminescence detection system (Sagecreation, Beijing, China). Densitometry was performed using lane-1 system (Sagecreation, Beijing, China).

Measurement of malondialdehyde (MDA) concentration

MDA concentration in aorta was measured by using thiobarbituric acid reactive substances (TBARS) assay kit (Nanjing Jiancheng Bioengineering Institute, Nanjing, Jiangsu, China) as previously described (Costa *et al.* 2009). Thoracic aorta was dissected out and snapped frozen at -80 °C, then homogenized on ice and centrifuged at 10,000×g for 5 min to collect the supernatant. The supernatant was mixed with 10 % trichloroacetic acid 1 ml and 0.67 % thiobarbituric acid 1 ml, followed by 30-min incubation in a boiling water bath. The absorbance of the mixture (532 nm) was measured. The results were expressed as nmol/mg protein.

Measurement of SOD activity

A SOD immunoassay kit was used to determine the total SOD activity in thoracic rings (Nanjing Jiancheng Bioengineering Institute, Nanjing, Jiangsu, China). Thoracic aorta was perfused with PBS to remove any red blood cells and homogenized on ice. The tissue lysate was centrifuged at 10,000×g for 5 min. The supernatant contains total SOD was then incubated with working solution for 20 min at 37 °C. Read the absorbance at 450 nm using a microplate reader. The results were expressed as U/mg protein.

Drugs

NaHS, Ang II, ACh, SNP were obtained from

Sigma (Vienna, Austria). Losartan was obtained from ALEXIS (Switzerland). Phe was obtained from TCI (Japan). NaHS was freshly prepared and used to generate H₂S in solution according to previous paper (Qu *et al.* 2006).

Statistical analysis

Data were expressed as the mean \pm SEM. The results were evaluated either by *t* test or 2-way ANOVA using Graphpad Prism. $p < 0.05$ was considered significant. The dose response curves were analyzed by 2-way ANOVA, followed by vertical contrast with Bonferroni adjustment.

Results

Effect of NaHS on plasma H₂S, Ang II level and hypertension in 2K1C rats

The plasma H₂S level in 2K1C rats (26.8 \pm 4.8 μ mol/l) was significantly lower than that in sham rats (50.6 \pm 6.9 μ mol/l, $p < 0.05$). 4 weeks of NaHS treatment elevated the plasma H₂S level to 56.4 \pm 9.2 μ mol/l ($p < 0.05$, Fig. 1A). However, Compared with sham rats, the plasma Ang II level was significantly higher in 2K1C rats (140.2 \pm 4.7 pg/ml vs. 101.1 \pm 1.9 pg/ml, $p < 0.01$). Treatment with NaHS for 4 weeks lowered the Ang II level to 111.5 \pm 3.4 pg/ml (Fig. 1B).




Fig. 1. Administration of NaHS for 4 weeks increased the plasma H₂S level (A), reduced the plasma Ang II level (B), and attenuated the development of hypertension (C) in 2K1C rats. Results are expressed as mean \pm SEM. $n = 6$, * $p < 0.05$, ** $p < 0.01$ vs. Sham; ## $p < 0.01$ vs. 2K1C

SBP was measured before and weekly after 2K1C surgery. No differences were observed in SBP among the groups before surgery. At the end of the first week, the SBP in 2K1C rats was increased from 105.0 \pm 1.2 mm Hg to 170.0 \pm 4.0 mm Hg, and remained elevated throughout the 4-week study period. Sham operation did not affect SBP. Intraperitoneally administrated with NaHS 56 μ mol/kg/day from the third day after surgery reduced the elevation of SBP (Fig. 1C) at the second week after surgery and remained lower than that of the non-treated 2K1C rats during the rest of the study period.

Effect of NaHS on Ang II-induced vasoconstriction

The Ang II-induced vasoconstriction was enhanced in the thoracic aorta of 2K1C rats. Such enhancement of vasoconstriction was suppressed by the 4-week treatment of NaHS (Fig. 2A,B). In all the three groups including sham and 2K1C without or with NaHS treatment, the vasoconstriction induced by Ang II was almost abolished by a 30-min preincubation of the aorta with the losartan (10 μ mol/l) (Fig. 2C).

Effect of NaHS on endothelium-dependent and-independent vasorelaxation

ACh-induced vasorelaxation was significantly

reduced in Phe-precontracted aorta in 2K1C rats (E_{max} 50.8±3.7 % vs. 100.2±0.3 % in sham rats, $p<0.05$). Treatment with NaHS for 4 weeks improved ACh-induced endothelium-dependent relaxation (64.8±9.4 %,

Fig. 3A,B). By contrast, the SNP-induced endothelium-independent relaxation showed no differences among sham, 2K1C, and NaHS-treated 2K1C rats. (Fig. 3C,D).




Fig. 2. The Ang II-induced contraction was attenuated by NaHS chronic treatment. NaHS chronic treatment ameliorated the contraction to Ang II (A, B). Pretreatment with losartan (10 μmol/l) for 30 min almost eliminated the Ang II-induced vasoconstriction in rat thoracic aorta (C). Results are expressed as mean ± SEM. n=6, * $p<0.05$, ** $p<0.01$ vs. Sham; ## $p<0.01$ vs. 2K1C




Fig. 3. Effect of NaHS chronic treatment on endothelium-dependent and endothelium-independent relaxation. NaHS 4-week administration enhanced ACh-induced vasorelaxation in 2K1C rats (A, B). No differences were detected in SNP-induced endothelium-independent relaxation (C, D). Results are expressed as mean ± SEM. n=6, ** $p<0.01$ vs. Sham; # $p<0.05$ vs. 2K1C

Effect of NaHS on oxidative stress-related proteins

Western blot analysis showed that protein level of AT₁R, NADPH oxidase isoform p67^{phox} and subunit Nox2, Nox4 was elevated in 2K1C rat thoracic aorta. The over-expression of AT₁R, p67^{phox}, Nox2 and Nox4 were reversed by chronic treatment with NaHS (Fig. 4).

Effect of NaHS on MDA level

The MDA level in the thoracic aorta of 2K1C rats was significantly elevated (671.6±94.6 vs. 339.9±77.6 nmol/gprot in sham rats, $p<0.05$). Treatment with NaHS for 4 weeks normalized the tissue level of MDA (356.9±42.9 nmol/gprot) (Fig. 5).




Fig. 4. NaHS chronic treatment reduces the exaggerated protein expression of AT₁R (A), p67^{phox} (B), Nox2 (C) and Nox4 (D) in thoracic aorta of 2K1C rats. Results are expressed as mean \pm SEM. n=4 or 5, * p<0.05 vs. Sham; # p<0.05 vs. 2K1C

Fig. 5. The MDA level in thoracic aorta was reduced by NaHS 4-week treatment. Results are expressed as mean \pm SEM. n=6, ** p<0.01 vs. Sham; ## p<0.01 vs. 2K1C

Effect of NaHS on SOD

The tissue SOD activity in rat thoracic aorta was markedly decreased, whereas NaHS chronic treatment reversed the decreased SOD activity in 2K1C rats (Fig. 6A). The downregulated protein expression of

SOD-1 was normalized by NaHS supplementation as well (Fig. 6B).

Discussion

The major findings of this study include: (1) NaHS reduces SBP in 2K1C hypertensive rats; (2) NaHS improves the endothelium-dependent relaxation and suppresses Ang II-induced contraction; (3) this study is the first time to show that H₂S downregulates AT₁R protein expression and oxidative stress in 2K1C hypertension. Taken together, the results suggest that H₂S protect endothelial function by decreasing oxidative stress in 2K1C rats. Figure 7 is a signal transduction diagram depicting the possible mechanisms and assumed pathways.

Previous studies have demonstrated that endothelial function is impaired in 2K1C hypertensive rats (Callera *et al.* 2000) and NaHS exhibits anti-hypertensive effect in this animal model (Lu *et al.* 2010), however, the effect of NaHS on endothelial function in renovascular hypertension remains poorly studied. In this study, we demonstrated that NaHS treatment lowered blood pressure in 2K1C hypertensive rats, which is

consistent to previous reports. Improvement of endothelium-dependent relaxation is a mechanism underlying the antihypertensive effect of NaHS. Suppression of vascular responsiveness to Ang II is also involved, evidenced by the attenuated contractile response of aorta to Ang II in rats treated with NaHS that is attributable to the lowered plasma level of Ang II and the downregulation of AT₁R in vascular cells.




Fig. 6. Treatment with NaHS for 4 weeks remarkably enhanced the tissue activity of SOD (A) and protein expression (B) of SOD-1. Results are expressed as mean ± SEM. n=4 or 5, * p<0.05, ** p<0.01 vs. Sham; # p<0.05, ## p<0.01 vs. 2K1C

The association between blood pressure elevation and vascular oxidative stress has been extensively reported in the development of hypertension, including essential, renovascular or malignant hypertension (Schulz *et al.* 2011). As a major source of ROS in vascular cells, NADPH oxidase can be activated by Ang II through acting on AT₁R (Drummond *et al.*

2011). Our results showed that NaHS treatment lowered plasma level of Ang II and downregulated aortic AT₁R protein expression in 2K1C hypertensive rats. As a consequence, the inhibition of Ang II-AT₁R action likely contributes to the inhibition of NADPH oxidases and the reduction of ROS production.




Fig. 7. Schematic diagram of mechanisms underlying the effect of NaHS against endothelial dysfunction in 2K1C rats. NaHS prevents endothelial dysfunction in thoracic aorta by inhibition of Ang II-AT₁R action and downregulation of NADPH oxidases, as well as upregulation of SOD.

Nox proteins represent the catalytic subunits of NADPH oxidase. In various animal models of hypertension including Ang II infusion, deoxycorticosterone acetate-salt, and renovascular hypertension as well as SHR, increased activity of Nox has been observed (Schulz *et al.* 2011). Ang II was reported to increase expression of several NADPH oxidase subunits, including Nox1 and Nox4 (Seshiah *et al.* 2002). In this study, we determined the expression of Nox2 and Nox4, the most important Nox isoforms in vascular cells (Brown and Griendling 2009), and found that in 2K1C rats, protein expressions of Nox2 and Nox4 were significantly increased in the aortic tissue. 4-week treatment with NaHS normalized the expression of Nox2 and Nox4. NaHS treatment also lowered the protein level of p67^{phox}, the regulatory cytosolic subunit of NADPH, in the aorta of 2K1C rats. These results suggested the direct

inhibitory effect of H₂S on NADPH oxidases. Furthermore, we showed that in NaHS-treated animals, the activity of the major antioxidant enzyme SOD is enhanced, accompanied by an upregulation of SOD-1 protein. Taken together, our results demonstrated the potent antioxidant effect of H₂S on vascular cells in renovascular hypertensive rats. The inhibition of vascular oxidative stress conferred by H₂S was further evidenced by the decreased MDA level in the aortic tissue.

NaHS is commonly used to deliver H₂S and increases H₂S concentrations rapidly (Li *et al.* 2008). NaHS incompletely dissociates into H₂S in Krebs solution, resulting in about 15-18.5 % being present as H₂S, with 81.5-85 % existing as hydrosulfide, HS⁻ and a trace of sulfide anion, S²⁻ (Dombkowski *et al.* 2004). Which form of H₂S (H₂S, HS⁻, or S²⁻) is physiologically active, or whether all three forms are active to varying extents are not known, however the active component is commonly termed as the chemical symbol "H₂S" (Al-Magableh *et al.* 2014). The physiological concentration of H₂S is difficult to measure and the measurement of H₂S not always gives the consistent values, varying from the previous micromolar range to the recent nanomolar range. Current literature reports several H₂S measurement methods, such as spectrophotometry, chromatography, ion-selective electrode and nanoparticles. The methylene blue method used in present study is the most common assay for detecting H₂S. This method revealed that human and rat serum contains 50-100 μM H₂S, which is in consistent with our results. However, this technique detects not only free H₂S but also other species such as HS⁻ and S²⁻, the true plasma concentration of H₂S is probably to be lower than tens of micromolar range. Although the methylene method has limitation, the present result is able to indicate that NaHS treatment increased H₂S concentration in 2K1C rats.

Previous studies showed that NaHS >50-100 μM is required to relax blood vessels in organ bath, 10-50 μmol/kg are needed to bring about significant falls in blood pressure. NaHS (56 μmol/kg/day, i.p.) treatment for 4 to 5 weeks decreased blood pressure and oxidative stress, inhibited atherogenesis and reduced atherosclerotic lesion size (Chen *et al.* 2011, Wang *et al.* 2009, Yan *et al.* 2004). Although lower doses of NaHS treatment (e.g. 10 or 30 μmol/kg/day, i.p.) were also effective in reducing blood pressure, a much longer treatment period (3 months) would be needed. Also, treatment with

different doses of NaHS (10, 30 and 90 μmol/kg/day) in SHR showed no obvious dose-related difference in blood pressure reduction (Shi *et al.* 2007). The present study showed the beneficial effect of NaHS in lowering blood pressure, improving endothelial function, reducing vascular superoxide production were all observed at a dose of 56 μmol/kg/day. However, the present study demonstrates limitation that we used a fixed dose of NaHS without performing a dose-dependent study, thus may miss the potentially toxic or beneficial effects of NaHS.

In summary, improvement of endothelial function is involved in the antihypertensive mechanism of H₂S. The protective effect of H₂S is attributable to, at least in part, suppression of vascular oxidative stress that involves inhibition of Ang II-AT₁R action, downregulation of NADPH oxidases, as well as upregulation of antioxidant enzyme SOD. H₂S may have therapeutic potential for renovascular hypertension.

Conflict of Interest

There is no conflict of interest.

Acknowledgements

This work was supported by National Natural Science Foundation of China (No.31171098), the Specialized Research Fund for the Doctoral Program of Higher Education of China (No.20121323110008) and the Natural Science Foundation of Hebei Province, China (No. C2013206080).

Abbreviations

ACh: acetylcholine; Ang II: angiotensin II; AT₁R: angiotensin II type1 receptor; BCA: bicinchoninic acid; CBS: cystathionine β-synthase; CSE: cystathionine γ-lyase; CAT: cysteine aminotransferase; EDHF: endothelial-dependent hyperpolarized factor; EDRF: endothelial-dependent relaxing factor; HRP: horseradish peroxidase; H₂S: hydrogen sulfide; MDA: malondialdehyde; 3-MST: 3-mercaptopyruvate sulfurtransferase; NADPH: nicotinamide adenine dinucleotide phosphate; Phe: phenylephrine; ROS: reactive oxygen species; NaHS: sodium hydrosulfide; SNP: sodium nitroprusside; SHR: spontaneously hypertensive rats; O₂⁻: superoxide anion; SOD: superoxide dismutase; SBP: systolic blood pressure; TBST: tris-buffered saline; 2K1C: two-kidney, one-clip

References

- AHMAD FU, SATTAR MA, RATHORE HA, TAN YC, AKHTAR S, JIN OH, PEI YP, ABDULLAH NA, JOHNS EJ: Hydrogen sulphide and tempol treatments improve the blood pressure and renal excretory responses in spontaneously hypertensive rats. *Ren Fail* **36**: 598-605, 2014.
- AL-MAGABLEH MR, KEMP-HARPER BK, NG HH, MILLER AA, HART JL: Hydrogen sulfide protects endothelial nitric oxide function under conditions of acute oxidative stress in vitro. *Naunyn Schmiedebergs Arch Pharmacol* **387**: 67-74, 2014.
- BAYRAKTUTAN U: Free radicals, diabetes and endothelial dysfunction. *Diabetes Obes Metab* **4**: 224-238, 2002.
- BROWN DI, GRIENGLING KK: Nox proteins in signal transduction. *Free Radic Biol Med* **47**: 1239-1253, 2009.
- CAI H, HARRISON DG: Endothelial dysfunction in cardiovascular diseases: the role of oxidant stress. *Circ Res* **87**: 840-844, 2000.
- CALLERA GE, VARANDA WA, BENDHACK LM: Impaired relaxation to acetylcholine in 2K-1C hypertensive rat aortas involves changes in membrane hyperpolarization instead of an abnormal contribution of endothelial factors. *Gen Pharmacol* **34**: 379-389, 2000.
- CHEN ZF, ZHAO B, TANG XY, LI W, ZHU LL, TANG CS, DU JB, JIN HF: Hydrogen sulfide regulates vascular endoplasmic reticulum stress in apolipoprotein E knockout mice. *Chin Med J (Engl)* **124**: 3460-3467, 2011.
- CHOI S, KIM HI, SANG HP, MI JL, JUN JY, KIM HL, CHUNG JH, YEUM CH: Endothelium-dependent vasodilation by ferulic acid in aorta from chronic renal hypertensive rats. *Kidney Res Clin Pract* **31**: 227-233, 2012.
- DIJKHORST-OEI LT, STROES ES, KOOMANS HA, RABELINK TJ: Acute simultaneous stimulation of nitric oxide and oxygen radicals by angiotensin II in humans in vivo. *J Cardiovasc Pharmacol* **33**: 420-424, 1999.
- DOMBKOWSKI RA, RUSSELL MJ, OLSON KR: Hydrogen sulfide as an endogenous regulator of vascular smooth muscle tone in trout. *Am J Physiol Regul Integr Comp Physiol* **286**: R678-R685, 2004.
- DRUMMOND GR, SELEMIDIS S, GRIENGLING KK, SOBEY CG: Combating oxidative stress in vascular disease: NADPH oxidases as therapeutic targets. *Nat Rev Drug Discov* **10**: 453-471, 2011.
- FORD A, AL-MAGABLEH M, GASPARI TA, HART JL: Chronic NaHS treatment is vasoprotective in High-Fat-Fed ApoE(-/-) mice. *Int J Vasc Med* **2013**: 915983, 2013.
- HIGASHI Y, SASAKI S, NAKAGAWA K, MATSUURA H, OSHIMA T, CHAYAMA K: Endothelial function and oxidative stress in renovascular hypertension. *N Engl J Med* **346**: 1954-1962, 2002.
- LI L, WHITEMAN M, GUAN YY, NEO KL, CHENG Y, LEE SW, ZHAO Y, BASKAR R, TAN CH, MOORE PK: Characterization of a novel, water-soluble hydrogen sulfide-releasing molecule (GYY4137): new insights into the biology of hydrogen sulfide. *Circulation* **117**: 2351-2360, 2008.
- LU M, LIU YH, GOH HS, WANG JJ, YONG QC, WANG R, BIAN JS: Hydrogen sulfide inhibits plasma renin activity. *J Am Soc Nephrol* **21**: 993-1002, 2010.
- MANCARDI D, PENNA C, MERLINO A, DEL SOLDATO P, WINK DA, PAGLIARO P: Physiological and pharmacological features of the novel gasotransmitter: hydrogen sulfide. *Biochim Biophys Acta* **1787**: 864-872, 2009.
- OLIVEIRA-SALES EB, NISHI EE, CARILLO BA, BOIM MA, DOLNIKOFF MS, BERGAMASCHI CT, CAMPOS RR: Oxidative stress in the sympathetic premotor neurons contributes to sympathetic activation in renovascular hypertension. *Am J Hypertens* **22**: 484-492, 2009.
- QU K, CHEN CP, HALLIWELL B, MOORE PK, WONG PT: Hydrogen sulfide is a mediator of cerebral ischemic damage. *Stroke* **37**: 889-893, 2006.
- SCHULZ E, GORI T, MUNZEL T: Oxidative stress and endothelial dysfunction in hypertension. *Hypertens Res* **34**: 665-673, 2011.
- SESHIAH PN, WEBER DS, ROCIC P, VALPPU L, TANIYAMA Y, GRIENGLING KK: Angiotensin II stimulation of NAD(P)H oxidase activity: upstream mediators. *Circ Res* **91**: 406-413, 2002.
- SHI YX, CHEN Y, ZHU YZ, HUANG GY, MOORE PK, HUANG SH, YAO T, ZHU YC: Chronic sodium hydrosulfide treatment decreases medial thickening of intramyocardial coronary arterioles, interstitial fibrosis, and ROS production in spontaneously hypertensive rats. *Am J Physiol Heart Circ Physiol* **293**: H2093-H2100, 2007.

-
- TADDEI S, VIRDIS A, MATTEI P, SALVETTI A: Vasodilation to acetylcholine in primary and secondary forms of human hypertension. *Hypertension* **21**: 929-933, 1993.
- TAKAHASHI N, SMITHIES O: Human genetics, animal models and computer simulations for studying hypertension. *Trends Genet* **20**: 136-145, 2004.
- TANIZAWA K: Production of H₂S by 3-mercaptopyruvate sulphurtransferase. *J Biochem* **149**: 357-359, 2011.
- VANHOUTTE PM, SHIMOKAWA H, TANG EH, FELETOU M: Endothelial dysfunction and vascular disease. *Acta Physiol (Oxf)* **196**: 193-222, 2009.
- WANG R: Two's company, three's a crowd: Can H₂S be the third endogenous gaseous transmitter? *FASEB J* **16**: 1792-1798, 2002.
- WANG R: Hydrogen sulfide: a new EDRF. *Kidney Int* **76**: 700-704, 2009.
- WANG Y, ZHAO X, JIN H, WEI H, LI W, BU D, TANG X, REN Y, TANG C, DU J: Role of hydrogen sulfide in the development of atherosclerotic lesions in apolipoprotein E knockout mice. *Arterioscler Thromb Vasc Biol* **29**: 173-179, 2009.
- YAN H, DU J, TANG C: The possible role of hydrogen sulfide on the pathogenesis of spontaneous hypertension in rats. *Biochem Biophys Res Commun* **313**: 22-27, 2004.
- YANG G, WU L, JIANG B, YANG W, QI J, CAO K, MENG Q, MUSTAFA AK, MU W, ZHANG S, SNYDER SH, WANG R: H₂S as a physiologic vasorelaxant: hypertension in mice with deletion of cystathionine gamma-lyase. *Science* **322**: 587-590, 2008.
- ZHAO W, ZHANG J, LU Y, WANG R: The vasorelaxant effect of H₂S as a novel endogenous gaseous K_{ATP} channel opener. *EMBO J* **20**: 6008-6016, 2001.
- ZHAO X, ZHANG LK, ZHANG CY, ZENG XJ, YAN H, JIN HF, TANG CS, DU JB: Regulatory effect of hydrogen sulfide on vascular collagen content in spontaneously hypertensive rats. *Hypertens Res* **31**: 1619-1630, 2008.
-