

Metabolic Status and Ghrelin Regulate Plasma Levels and Release of Ovarian Hormones in Layer Chicks

A. V. SIROTKIN^{1,2,3}, A. H. HARRATH³, R. GROSSMANN⁴

¹Department Zoology and Anthropology, Constantine the Philosopher University, Nitra, Slovakia,

²Department Genetics and Reproduction, Research Institute of Animal Production, Lužianky,

Slovakia, ³Zoology Department, College of Science, King Saud University, Riyadh, Saudi Arabia,

⁴Department Functional Genomics & Bioregulation, Friedrich Loeffler Institute, Mariensee, Neustadt, Germany

Received January 26, 2016

Accepted May 20, 2016

On-line October 26, 2016

Summary

The aim of the present study was to examine the role of nutritional status, the metabolic hormone ghrelin and their interrelationships in the control of chicken hormones involved in the regulation of reproduction. For this purpose, we identified the effect of food deprivation, administration of ghrelin 1-18 and their combination on plasma levels of testosterone (T), estradiol (E), arginine-vasotocin (AVT) and growth hormone (GH) as well as the release of these hormones by isolated and cultured ovarian fragments. It was observed that food deprivation reduces plasma T and E and increases plasma AVT and GH levels. Food restriction also reduced the amount of E produced by isolated ovaries, but it did not affect the ovarian secretion of T and AVT. No ovarian GH secretion was detected. Ghrelin administered to *ad libitum* fed chickens did not affect plasma T and E levels, but it did increase plasma GH and AVT concentrations. Moreover, it partially prevented the effect of food deprivation on plasma E and AVT levels, but not on T or GH levels. Ghrelin administration to control birds promoted ovarian T, but not E or AVT release and reduced T and no other hormonal outputs in birds subjected to food restriction. Our results (1) confirmed the ovarian origin of the main plasma T and E and the extra-ovarian origin of the main blood AVT and GH; (2) showed that food deprivation-induced suppression of reproduction may be caused by suppression of T and E and the promotion of AVT and GH release; (3) suggest the involvement of ghrelin in control chicken E, AVT and GH output; and (4) indicates that ghrelin can either mimic or modify the effect of the intake of low calories on chicken plasma and ovarian hormones, i.e. it can mediate the

effect of metabolic state on hormones involved in the control of reproduction.

Key words

Ghrelin • Ovary • Testosterone • Estradiol • GH • Arginine-vasotocin

Corresponding author

A. V. Sirotkin, Department Zoology and Anthropology, Constantine the Philosopher University, 949 74 Nitra, Slovakia. Fax: +421-37-6546361. E-mail: asirotkin@ukf.sk

Introduction

Reproduction is an energy-demanding process; therefore, it should be synchronized with optimal nutritional conditions. It has been postulated that the hypothetical mediator of the nutritional effect on reproduction (1) should be affected by nutrition and (2) controls reproduction, and (3) its changes should mimic and modify the effect of nutrition on reproduction. On the basis of these criteria, it is proposed that the nutritional status affects mammalian reproductive processes via the metabolic hormones, leptin, ghrelin and obestatin (Tena-Sempere 2008, Navarro and Kaiser 2013, Roa and Tena-Sempere 2014, Sirotkin 2014). In birds, malnutrition inhibits ovarian folliculogenesis, ovulation/egg laying and ovarian hormones release (Hocking 2004, Sirotkin and Grossmann 2015) most likely via the induction of

ovarian follicular cells apoptosis (Paczoska-Eliasiewicz *et al.* 2003). Malnutrition is associated with changes in plasma and brain ghrelin as well as its receptors (Kaiya *et al.* 2007, 2013, Sirotkin *et al.* 2013). Ghrelin i.m. administration in chicken is able to reduce plasma progesterone (P) levels (Sirotkin *et al.* 2015), which directly alters proliferation, apoptosis, steroidogenesis and protein kinases in cultured ovarian cells (Sirotkin *et al.* 2006, Sirotkin and Grossmann, 2007, 2008), and prevents the food restriction-induced decrease in ovarian testosterone (T), estradiol (E) and arginine-vasotocin (AVT) release (Sirotkin *et al.* 2015). Such data have demonstrated the importance of ghrelin in integrating nutrition and reproduction and its potential applicability for the improvement of farm avian reproduction. Nevertheless, the mediatory role of ghrelin in the metabolic control of ovarian functions has only been previously demonstrated in one study (Sirotkin *et al.* 2015). Moreover, in the described study, ghrelin and food restriction effects only on hormonal release by ovarian tissue *in vitro*, but not on plasma hormones under *in vivo* conditions, were examined.

The general aim of the present study was to examine the role of the nutritional status, metabolic hormone ghrelin and their interrelationships in the control of chicken steroid and peptide hormones involved in the regulation of reproduction. For this purpose, we used both *in-vivo* and *in-vitro* approaches to identify the effect of food deprivation, administration of ghrelin 1-18 and their combination on plasma level of hormones (T, E, AVT and growth hormone, GH) whose are known autocrine/paracrine and endocrine regulators of both mammalian and avian ovarian functions (Sirotkin 2005, 2014, Luna *et al.* 2014, Hrabia 2015) as well as the release of these hormones by isolated and cultured ovarian fragments.

Materials and Methods

Animal experiments, tissue collection and culture

Young (approximately 8 months of age) White Leghorn hens (LSL), weight 1.1-1.2 kg with an egg-laying rate of more than 95 %, were housed in individual cages under standard conditions at the Experimental Station of the Institute of Animal Science on a photoperiod 12L:12D (illumination 8.00-20.00). The conditions of their care, manipulations and use corresponded the instructions of the EC no. 178/2002 and related EC documents, and the protocols were

approved by the local ethics commission. After a two-month adaptation period to the conditions of the experimental farms, the hens were divided into four experimental groups: (1) the control group was fed *ad libitum*, no hormone treatment; (2) the group fed *ad libitum* and treated with i.m. injection of human recombinant research grade ghrelin 1-18 (Peptides International Inc., Louisville, Kentucky, USA) (this truncated ghrelin analogue ghrelin 1-18 mimicked the effect of full-length ghrelin 1-28 on chicken ovarian cells (Sirotkin and Grossmann 2008); (3) the group subjected to food deprivation, no hormone treatment; and (4) the group subjected to food deprivation and treated with i.m. injection of human recombinant ghrelin 1-18. The animals of the food-deprivation groups had no access to food during the entire experiment for 72 h, whereas all of the animals had permanent access to drinking water. Hormonal treatments combined with food deprivation began together with food restriction. Ghrelin was dissolved in sterile 0.7 % NaCl immediately prior to the start of the experiments and injected i.m. at doses 30 µg/animal in 1 ml of 0.7 % NaCl. This consecutive injection was done for 3 days, every 10-12 h, in the daytime (at 8.00 and 18.00). This dose, injection and sampling time (details are below) were comparable to the amount of hormones in the chicken organism and those treated with previously reported experiments (Kaiya *et al.* 2007, Sirotkin *et al.* 2013, Sirotkin and Grossmann 2015). Next, 1.5 h after the last injection (between 9.00 and 11.00 a.m.), the animals were killed by decapitation. Their blood was collected in heparinized tubes, and the plasma was separated by 10 min centrifugation at 500 g and frozen at -70°C until radioimmunoassay (RIA) or enzyme immunoassay (EIA). The largest (F1) follicles were isolated from the ovary. The stage of folliculogenesis was determined by recording the time of the last oviposition and by the size and position of the next ovarian follicle. Fragments of the follicular wall (5 mm in diameter, weight 24±8 mg) were isolated as previously described (Sirotkin and Grossmann 2003, 2006). After washing three times in sterile culture medium (DMEM/F-12 1:1 mixture supplemented with 10 % bovine fetal serum and 1 % antibiotic-antimycotic solution (all from Sigma, St. Louis, USA), these fragments were cultured without treatment for 2 days in 2 ml culture medium in Falcon 24-well plates (Becton Dickinson, Lincoln Park, USA) at 38.5 °C under 5 % CO₂ in humidified air. This

protocol yields the maximal accumulation of ovarian hormones in the culture medium, which is the most reliable characteristic of ovarian secretory activity (Sirotkin and Grossmann 2003).

Immunoassay

Concentrations of testosterone (T), estradiol (E), arginine-vasotocin (AVT) and growth hormone (GH) were determined in 25 µl aliquots of plasma or incubation medium by EIA and RIA, whose were previously validated for use in culture medium (Sirotkin *et al.* 2006, Sirotkin and Grossmann 2007). These hormones were considered as the indices of ovarian secretory activity, stress, response to hormonal stimuli and the key regulators of both mammalian (Sirotkin 2014) and chicken (Sirotkin and Grossmann 2006, 2007, 2008, Luna *et al.* 2014, Sirotkin 2014, Hrabia 2015) ovarian functions.

T was assayed according to Münster (1989) using antisera against steroids (produced in the Institute of Animal Science, Neustadt, Germany). The sensitivity of the assay was 10 pg/ml. The cross-reactivity of the T antiserum was ≤96 % to dihydrotestosterone, ≤3 % to androstenedione, ≤0.01 % to P₄ and E₂, ≤0.02 % to cortisol and ≤0.001 % to corticosterone. The inter- and intraassay coefficients of variation were 12.3 % and 6.8 %, respectively.

E concentrations were evaluated according to Münster (1989) using antisera against steroids (produced by the Institute of Animal Science, Neustadt, Germany) with an assay sensitivity of 5 pg/ml. The cross-reactivity of the E₂ antiserum was <2 % to estrone, ≤0.3 % to estriol, ≤0.004 % to T and ≤0.0001 % to P₄ and cortisol. The inter- and intraassay coefficients of variation did not exceed 16.6 % and 11.7 %, respectively.

AVT was determined using RIA according to Gray and Simon (1983). The anti-AVT antiserum was kindly provided by Dr. D.A. Gray (Max-Plank Institute for Physiological and Clinical Research, Bad Nauheim, Germany), which cross-reacted ≤1.0 % with mesotocin and angiotensin II. The sensitivity of the RIA was 0.3 pg/ml. The inter- and intraassay coefficients of variation did not exceed 8.8 % and 7.2 %, respectively.

GH was measured using the EIA based on the EIA used for porcine GH (Serpel *et al.* 1993) and adapted for determination of chicken GH (Zheng *et al.* 2007). Chicken GH for standards (AFP-9020C), iodination (AFP 7678B) and antiserum against chicken GH (AFP-551-11-1-86, dilution 1:720,000) were kindly

provided by Dr. A.P.F. Parlow (National Hormones and Pituitary Program, Bethesda, USA). This antiserum has 0.7 % cross-reactivity with chicken prolactin and <0.001 % cross-reactivity with P₄, T, E₂ and AVT. The sensitivity of the assay was 0.2 ng/ml. The inter- and intraassay coefficients of variation did not exceed 12.9 % or 10.8 %, respectively.

Statistics

The data shown are the mean of the values obtained in three separate experiments performed on separate days using independent animals (8 animals per group) and their ovaries. In each *in vitro* experiment, each experimental group consisted of six culture wells with ovarian fragments. Assays of the hormone levels in the incubation media were performed in duplicate. The values of the blank control were subtracted from the value determined using RIA/EIA in the cell-conditioned medium to exclude any non-specific background (less than 15 % of the total values). The rates of substance secretion were calculated per mg tissue/day. Significant differences between the experiments were evaluated using two-ways ANOVA. When effects of the treatments were revealed, data obtained from the experimental and control groups were compared using the Wilcoxon-Mann-Whitney multiple range test with Sigma Plot 11.0 statistical software (Systat Software, GmbH., Erkrath, Germany). Differences compared to control were considered significant if P<0.05.

Results

In chicken blood plasma T, E, AVT and GH were detected (Fig. 1). The culture medium conditioned by cultured ovarian fragments contained normal amounts of T, E and AVT, but not of measurable GH (Fig. 2). These parameters were affected by food restriction, administration of ghrelin and the combination of the above factors.

Food deprivation significantly reduced the concentrations of both T (Fig. 1A), E (Fig. 1B) and increased the level of both AVT (Fig. 1C) and GH (Fig. 1D) in plasma. Ghrelin administered to *ad libitum* fed chickens did not affect plasma T (Fig. 1A) or E (Fig. 1B) levels, but it did increase plasma AVT (Fig. 1C) and GH (Fig. 1D) concentrations. Moreover, ghrelin administration could partially prevent the effect of food restriction on plasma E (Fig. 1B) and AVT (Fig. 1C), but not on T (Fig. 1A) or GH (Fig. 1D) levels.

Analysis of hormones produced by ovarian tissue *in vitro* demonstrated that food deprivation reduced the amount of E produced by the ovary (Fig. 2B), but it did not affect the ovarian secretion of T (Fig. 1A) or AVT (Fig. 2C). Ghrelin administration to control birds promoted ovarian T (Fig. 2A), but not E (Fig. 2B) or AVT (Fig. 2C) release. Furthermore, ghrelin reduced T (Fig. 2A), but not E (Fig. 2B) or AVT (Fig. 2C) output by the ovaries of birds subjected to food deprivation.

Discussion

Our observations showed the availability of steroid hormones, AVT and GH in chicken blood plasma and the release of steroid hormones and AVT by cultured chicken ovarian tissue, but our assay failed to detect measurable GH production by chicken ovarian cells reported by other investigators (Luna *et al.* 2014, Hrabia 2015) which can be explained by low GH production by chicken ovarian cells in our experiments.




Fig. 1. Effect of food restriction and administration of ghrelin 1-18 (*in vivo*) on the levels of testosterone (A), estradiol (B), arginine-vasotocin (C) and growth hormone (D) in chicken blood plasma. Data are the mean \pm S.E.M. Differences between the groups at $P < 0.05$ were considered significant: a) effect of hormones administration (differences between control and hormone-treated chicken); b) effect of food restriction (differences between corresponding groups of chickens subjected and not subjected to food restriction).




Fig. 2. Effect of food restriction and administration of ghrelin *in vivo* on the release of testosterone (**A**), estradiol (**B**) and arginine-vasotocin (**C**) by isolated chicken ovarian fragments. Legends are similar to those presented in Fig. 1.

A comparison of the hormones that are available in general circulation and those that are produced by isolated ovarian tissues and its changes under the effect of extra-ovarian factors demonstrates that some hormones found in blood (E, T, AVT) are mainly ovarian, but other factors (GH) are mainly extra-ovarian in origin. Furthermore, differences in the mechanisms controlling the release of these hormones were observed. For example, food deprivation reduced E in both plasma and ovarian fragment-conditioned medium. This finding suggests that the negative metabolic state directly affects ovarian E output. In contrast, the effect of food restriction on plasma T and AVT, but not on their release by isolated ovarian tissue, suggest that the metabolic state controls

T and AVT release via upstream extra-ovarian regulatory mechanisms (probably via hypothalamo-hypophysial system involved in metabolic control of gonadal functions, Roa and Tena-Sempere 2014). One such extra-ovarian regulator could be ghrelin. Ghrelin could influence chicken ovarian steroid hormones in our previous (Sirotnik *et al.* 2006, 2015, Sirotnik and Grossmann 2007, 2008) and present studies. In some previous (Sirotnik *et al.* 2006, Sirotnik and Grossmann 2008), but not in our present experiments, ghrelin promoted chicken ovarian AVT output, which can be explained by variations in initial state of AVT producing cells between the experiments. Furthermore, our present observations of ghrelin-induced increase in plasma GH

confirmed previous findings (Baudet and Harvey 2003) that ghrelin is a physiological GH secretagogue not only in the mammalian, but also in the avian pituitary. The observed ghrelin-induced changes could be due to influence of ghrelin on hormonal regulators of reproduction at the level of the ovary (steroids), the upstream hypothalamo-hypophysial system (GH) or on the differentiation of CNS and ovarian tissue. The site and fine mechanisms of ghrelin influence on organisms require further studies, but the present observations suggest the action of ghrelin at both CNS and gonadal level. In our experiments, the pronounced effect of ghrelin on testosterone release by ovarian tissue (Fig. 2A) was not associated with the corresponding changes in plasma testosterone (Fig. 1A). It suggests that the direct action of ghrelin on the ovary could be masked by additional factors in CNS or general circulation affecting steroid transport, binging or degradation.

Our observations confirmed previous reports that food deprivation reduces ovarian steroid hormones (Paczoska-Eliasiewicz *et al.* 2003, Sirotkin and Grossmann 2015) and promotes GH (Buyse *et al.* 2000, 2002) levels in chicken plasma. The food deprivation-induced increase in blood AVT levels observed in our experiments and the fasting-induced reduction in ovarian AVT release observed in our previous (Sirotkin and Grossmann 2015) but not our present studies indicated that the metabolic state can affect blood and maybe ovarian AVT. Variations in initial state of ovarian AVT producing cells between the experiments could influence ovarian AVT response not only to ghrelin (details are above) but also to food restriction. The steroid hormones GH and AVT are known regulators of both mammalian and avian reproductive processes including ovarian steroidogenesis (Sirotkin 2005, 2014, Luna *et al.* 2014, Hrabia 2015). Thus, it is possible that food deprivation

can affect reproductive processes *via* changes in the release of these peptide and steroid hormones. In addition, the metabolic state can affect these hormonal regulators of reproduction via ghrelin. Food consumption affects the production of chicken ghrelin, ghrelin acylation and ghrelin receptor (Kaiya *et al.* 2007, 2013, Sirotkin *et al.* 2013, Sintubin *et al.* 2014). Furthermore, this is the first evidence that ghrelin can mimic the effect of food deprivation on plasma GH and AVT, and that ghrelin can modify the effect of food deprivation on chicken plasma hormones. Taken together, combined with the data on the importance of ghrelin in the control of basic ovarian functions (details are above), these evidence suggest that ghrelin can be the key hormone mediating the effect of the metabolic state on downstream hormonal regulators of avian ovarian functions.

Conflict of Interest

There is no conflict of interest.

Acknowledgements

The authors express their deep gratitude to G. Neuhaus for the skillful technical help in the arrangement of the experiments and AVT RIA, I. Stelter for EIA of the steroid hormones, K. Tothova and Z. Kuklova for assistance in the statistical processing of the data, as well as to Dr. D. A. Gray (Max-Plank Institute for Physiological and Clinical Research, Bad Nauheim, Germany) for the kind gift of antiserum against AVT. We thank Dr. A. P. F. Parlow (National Hormones and Pituitary Program, Bethesda, MD, USA) for kindly providing reagents for the chicken GH assay. This work was financially supported by the German and Slovak Ministries of Agriculture and by the Deanship of Scientific Research at King Saud University Research group No. RG#164.

References

- BAUDET ML, HARVEY S: Ghrelin-induced GH secretion in domestic fowl in vivo and in vitro. *J Endocrinol* **179**: 97-105, 2003.
- BUYSE J, DECUYPERE E, DARRAS VM, VLEURICK LM, KÜHN ER, VELDHUIS JD: Food deprivation and feeding of broiler chickens is associated with rapid and interdependent changes in the somatotrophic and thyrotrophic axes. *Br Poult Sci* **41**: 107-116, 2000.
- BUYSE J, JANSENS K, VAN DER GREYTEN S, VAN AS P, DUCUYPERE E, DARRAS VM: Pre-and postprandial changes in plasma hormone and metabolite levels and hepatic deiodinase activities in meal-fed broiler chickens. *Br J Nutr* **88**: 641-653, 2002.
- GRAY DA, SIMON E: Mammalian and avian antidiuretic hormone: studies related to possible species variation in osmoregulatory systems. *J Comp Physiol* **151**: 241-246, 1983.

- HOCKING PM: Roles of body weight and feed intake in ovarian follicular dynamics in broiler breeders at the onset of lay and after a forced molt. *Poult Sci* **83**: 2044-2050, 2004.
- HRABIA A: Growth hormone production and role in the reproductive system of female chicken. *Gen Comp Endocrinol* **220**: 112-118, 2015.
- KAIYA H, SAITO ES, TACHIBANA T, FURUSE M, KANGAWA K: Changes in ghrelin levels of plasma and proventriculus and ghrelin mRNA of proventriculus in fasted and refed layer chicks. *Domest Anim Endocrinol* **32**: 247-259, 2007.
- KAIYA H, KANGAWA K, MIYAZATO M: Update on ghrelin biology in birds. *Gen Comp Endocrinol* **190**: 170-175, 2013.
- LUNA M, MARTINEZ-MORENO CG, AHUMADA-SOLORZANO MS, HARVEY S, GARRANZA M, ARSMBURO C: Extrapituitary growth hormone in the chicken reproductive system. *Gen Comp Endocrinol* **203**: 60-68, 2014.
- MÜNSTER E: Entwicklung von enzymimmunologischen Messverfahren auf Mikrotitrationsplatten zur Bestimmung von Testosteron und Progesteron im Blutplasma. In: *Thesis. Institut for Animal Production and Breeding of the University of Hohenheim*. 1989, pp 154.
- PACZOSKA-ELIASIEWICZ HE, GERTLER A, PROSZKOWIEC M, PROUDMAN J, HRABIA A, SECHMAN A, MIKA M, JACEK T, CASSY S, RAVER N, RZASA J: Attenuation by leptin of the effects of fasting on ovarian function in hens (*Gallus domesticus*). *Reproduction* **126**: 739-751, 2003.
- PACZOSKA-ELIASIEWICZ HE, PROSZKOWIEC-WEGLARZ M, PROUDMN J, JACEK T, MIKA M, SECHMAN A, RZASA J, GERTLER A: Exogenous leptin advances puberty in domestic hen. *Domest Anim Endocrinol* **31**: 211-226, 2006.
- PRAKASH BS, MEYER HH, SCALLENBERER E, VAN DE WIEL DF: Development of a sensitive enzymeimmunoassay (EIA) for progesterone determination in unextracted bovine plasma using the second antibody technique. *J Steroid Biochem* **28**: 623-627, 1987.
- ROA J, TENA-SEMPERE M: Connecting metabolism and reproduction: Roles of central energy sensors and key molecular mediators. *Mol Cell Endocrinol* **397**: 4-14, 2014.
- SERPEK B, ELSAESER F, MEYER HH: Development of an enzyme immunoassay for the determination of porcine growth hormone in plasma. *Analytica Chimica Acta* **275**: 183-187, 1993.
- SINTUBIN P, GREENE E, COLLIN A, BORDAS A, ZERJAL T, TESSERAUD S, BUYSE J, DRIDI S: Expression profile of hypothalamic neuropeptides in chicken lines selected for high or low residual feed intake. *Neuropeptides* **48**: 213-220, 2014.
- SIROTKIN AV, GRISSMANN R: Role of tyrosine kinase- and MAP kinase-dependent intracellular mechanisms in control of ovarian functions in the Domestic fowl (*Gallus domesticus*) and in mediating effects of IGF-II. *J Reprod Dev* **49**: 99-106, 2003.
- SIROTKIN AV: Control of reproductive processes by growth hormone: extra- and intracellular mechanisms. *Vet J* **170**: 307-317, 2005.
- SIROTKIN AV, GROSSMANN R, MARIA-PEON MT, ROA J, TENA-SEMPERE M, KLEIN S: Novel expression and functional role of ghrelin in chicken ovary. *Mol Cell Endocrinol* **257-258**: 15-25, 2006.
- SIROTKIN AV, GROSSMANN R: The role of protein kinase A and cyclin-dependent (CDC2) kinase in the control of basal and IGF-II-induced proliferation and secretory activity of chicken ovarian cells. *Anim Reprod Sci* **92**: 169-181, 2006.
- SIROTKIN AV, GROSSMANN R: Leptin directly controls proliferation, apoptosis and secretory activity of cultured chicken ovarian cells. *Comp Biochem Physiol A Mol Integr Physiol* **148**: 422-429, 2007a.
- SIROTKIN AV, GROSSMANN R: The role of ghrelin and some intracellular mechanisms in controlling the secretory activity of chicken ovarian cells. *Comp Biochem Physiol A Mol Integr Physiol* **147**: 239-246, 2007b.
- SIROTKIN AV, GROSSMANN R: Effects of ghrelin and its analogues on chicken ovarian granulosa cells. *Domest Anim Endocrinol* **34**: 125-134, 2008.
- SIROTKIN AV, PAVLOVA S, TENA-SEMPERE M, GROSSMANN R, JIMENEZ MR, RODRIGUEZ JM, VALENZUELA F: Food restriction, ghrelin, its antagonist and obestatin control expression of ghrelin and its receptor in chicken hypothalamus and ovary. *Comp Biochem Physiol A Mol Integr Physiol* **164**: 141-153, 2013.

- SIROTKIN AV: *Regulators of Ovarian Functions*. Nova Science Publishers Inc., New York, 2014, pp 194.
- SIROTKIN AV, GROSSMANN R: Interrelationship between feeding level and the metabolic hormones leptin, ghrelin and obestatin in control of chicken egg laying and release of ovarian hormones. *Comp Biochem Physiol A Mol Integr Physiol* **184**: 1-5, 2015.
- ZHENG JX, LIU ZZ, YANG N: Deficiency of growth hormone receptor does not affect male reproduction in dwarf chickens. *Poultry Sci* **86**: 112-117, 2007.
-