

Vlasta T.: Jak může současná věda přispět k praktické druhové ochraně rostlin? Příběh kohátky kalíškaté (Živa 2021, 4: 162–166)

Literatura

- Bieker V. C. & Martin M. D. (2018) Implications and future prospects for evolutionary analyses of DNA in historical herbarium collections. – *Botany Letters* 165: 409–418.
- Eckert C. G., Samis K. E. & Loughheed S. C. (2008) Genetic variation across species' geographical ranges: The central-marginal hypothesis and beyond. – *Molecular Ecology* 17: 1170–1188.
- Frankham R., Ballou J. & Briscoe D. (2010) – *Introduction to Conservation Genetics* (2nd ed.). Cambridge University Press.
- Hamrick J.L. & Godt M. J. W. (1996) Effects of life history traits on genetic diversity in plant species. – *Philosophical Transactions: Biological Sciences* 351: 1291–1298.
- Honnay O. & Jacquemyn H. (2007) Susceptibility of Common and Rare Plant Species to the Genetic Consequences of Habitat Fragmentation. – *Conservation Biology* 21: 823–831.
- Kaplan Z., Danihelka J., Štěpánková J., Bureš P., Zázvorka J., Hroudová Z., Ducháček M., Grulich V., Řepka R., Dančák M., Prančl J., Šumberová K., Wild J., & Trávníček B. (2015) Distributions of vascular plants in the Czech Republic. Part 1. – *Preslia* 87: 417–500.
- Leimu R., Mutikainen P., Koricheva J. & Fischer M. (2006) How general are positive relationships between plant population size, fitness and genetic variation? – *Journal of Ecology* 94: 942–952.
- Münzbergová Z., Gabrielová J. & Černá L. Priority druhové ochrany cévnatých rostlin České republiky (2011) – *Příroda* 31: 5-9.
- Noel F., Machon N. & Porcher E. (2007) No genetic diversity at molecular markers and strong phenotypic plasticity in populations of *Ranunculus nodiflorus*, an endangered plant species in France. – *Annals of Botany* 99: 1203–1212.
- Petr L. & Novák J. (2014) High vegetation and environmental diversity during the Late Glacial and Early Holocene on the example of lowlands in the Czech Republic. – *Biologia* 69: 847–862.
- Plenk K., Bardy K., Höhn M. & Kropf M. (2019) Long-term survival and successful conservation? Low genetic diversity but no evidence for reduced reproductive success at the north-westernmost range edge of *Poa badensis* (*Poaceae*) in Central Europe. – *Biodiversity and Conservation* 28: 1245–1265.
- Pokorný P., Sádlo J. & Bernardová A. (2010) Holocene history of *Cladium mariscus* (L.) pohl in the Czech republic. Implications for species population dynamics and palaeoecology. – *Acta Palaeobotanica* 50: 65–76.
- Schönswetter P., Stehlík I., Holderegger R. & Tribsch A. (2005) Molecular evidence for glacial refugia of mountain plants in the European Alps. – *Molecular Ecology* 14: 3547–3555.
- Šmídová A., Münzbergová Z. & Plačková I. (2011) Genetic diversity of a relict plant species, *Ligularia sibirica* (L.) Cass. (*Asteraceae*). – *Flora* 20: 151–157.
- Štěpánková J. (ed). (2010) *Květena České republiky* 8. – Academia, Praha.
- Vlasta T., Jarošová A., Šurinová M. & Münzbergová Z. (2020) Development of microsatellite markers for the perennial plant *Tofieldia calyculata*. – *Applications in Plant Sciences* 8: 1–5.

Williams J. W. et al (2008) The Neotoma Paleocology Database: A multi-proxy international community-curated data resource – *Quaternary Research* 89, 156-177