



INSTITUTE OF MATHEMATICS

THE CZECH ACADEMY OF SCIENCES

**A Galvin-Hajnal theorem for generalized
cardinal characteristics**

Chris Lambie-Hanson

Preprint No. 21-2022

PRAHA 2022

A GALVIN-HAJNAL THEOREM FOR GENERALIZED CARDINAL CHARACTERISTICS

CHRIS LAMBIE-HANSON

ABSTRACT. We prove that a variety of generalized cardinal characteristics, including meeting numbers, the reaping number, and the dominating number, satisfy an analogue of the Galvin-Hajnal theorem, and hence also of Silver's theorem, at singular cardinals of uncountable cofinality.

1. INTRODUCTION

One of the seminal results in cardinal arithmetic, and one of the first indications that there are nontrivial ZFC constraints on the behavior of the continuum function at singular cardinals, is *Silver's theorem*.

Theorem 1.1 (Silver [18]). *Suppose that κ is a singular cardinal of uncountable cofinality, $\eta < \text{cf}(\kappa)$ is an ordinal, and the set of cardinals*

$$\{\mu < \kappa \mid 2^\mu \leq \mu^{+\eta}\}$$

is stationary in κ . Then $2^\kappa \leq \kappa^{+\eta}$.

Silver's original proof of this theorem involves a generic ultrapower argument; a purely combinatorial argument for the theorem was soon provided by Baumgartner and Prikry [1]. Around the same time, a generalization of Silver's theorem was proven by Galvin and Hajnal. The following statement of (a corollary of) their theorem involves the notion of the *Galvin-Hajnal rank* $\|\varphi\|_S$ of a function φ ; see Definition 2.1 below for its formal definition.

Theorem 1.2 (Galvin-Hajnal [4]). *Suppose that κ is a singular cardinal of uncountable cofinality, $\langle \kappa_i \mid i < \text{cf}(\kappa) \rangle$ is an increasing, continuous sequence of cardinals converging to κ , $S \subseteq \text{cf}(\kappa)$ is stationary, and $\varphi : S \rightarrow \text{On}$ is a function such that, for all $i \in S$, we have $2^{\kappa_i} \leq \kappa_i^{+\varphi(i)}$. Then $2^\kappa \leq \kappa^{+\|\varphi\|_S}$.*

This theorem does indeed generalize Silver's theorem, since, as we shall see, given any stationary subset S of a regular uncountable cardinal θ , and given any ordinal $\eta < \theta$, if φ is the constant function on S taking value η , then $\|\varphi\| = \eta$.

One of the central aspects of research into cardinal arithmetic is the study of certain methods of measuring the “size” of the power set of a cardinal κ that are in a sense *finer* than simply looking at the value of 2^κ . At singular cardinals, these methods come from two primary sources, with some overlap between the two:

- Shelah's PCF theory;

2010 *Mathematics Subject Classification*. 03E04, 03E05, 03E17.

Key words and phrases. singular cardinals, cardinal characteristics, cardinal arithmetic, Silver's theorem, Galvin-Hajnal theorem.

- generalizations of cardinal characteristics of the continuum to singular cardinals.

Certain of these methods are known to satisfy versions of Silver's theorem or the Galvin-Hajnal theorem. For example, in [17, §2, Claim 2.4], Shelah proves a variation of Theorem 1.2 involving PCF-theoretic pseudopowers $\text{pp}_J(\kappa)$ and $\text{pp}_J(\kappa_i)$ in place of the cardinals 2^κ and 2^{κ_i} , respectively. Also, Kojman, in [12], proves that certain *density* numbers satisfy an analogue of Silver's theorem (see Section 3 for details).

In this paper, we prove versions of the Galvin-Hajnal theorem for a variety of cardinal characteristics of the continuum generalized to singular cardinals of uncountable cofinality, focusing in particular on meeting numbers, the reaping number, and the dominating number. Before proceeding to a summary of our results, let us say a few words about our approach to cardinal characteristics at singular cardinals in general. There are often multiple natural ways to generalize familiar cardinal characteristics of the continuum to singular cardinals. For example, when defining the dominating number \mathfrak{d}_κ at a singular cardinal κ , any of the following possible definitions of \mathfrak{d}_κ seems potentially reasonable (see the end of this introduction for any undefined notation):

- $\text{cf}({}^\kappa\kappa, <_0)$, where, given $f, g \in {}^\kappa\kappa$, we let $f <_0 g$ if and only if $|\{i < \kappa \mid g(i) \leq f(i)\}| < \kappa$;
- $\text{cf}({}^\kappa\kappa, <_1)$, where, given $f, g \in {}^\kappa\kappa$, we let $f <_1 g$ if and only if $\{i < \kappa \mid g(i) \leq f(i)\}$ is bounded below κ ;
- $\text{cf}(\text{cf}({}^\kappa\kappa), <_2)$, where, given $f, g \in \text{cf}({}^\kappa\kappa)$, we let $f <_2 g$ if and only if $|\{i < \text{cf}(\kappa) \mid g(i) \leq f(i)\}| < \text{cf}(\kappa)$.

In all such choices that we face here, we opt for the definition that emphasizes the *cardinality* of κ over its *cofinality*, as, at least in this context, this seems to be what gives rise to the most genuinely new behavior at the singular cardinal κ . So, for instance, we will define \mathfrak{d}_κ to be what is called $\text{cf}({}^\kappa\kappa, <_0)$ above. (It is not difficult to show that what is called $\text{cf}(\text{cf}({}^\kappa\kappa), <_2)$ above is in fact nothing other than $\mathfrak{d}_{\text{cf}(\kappa)}$.)

We also note here that in this paper we are only considering cardinal characteristics at a singular cardinal κ that are provably strictly greater than κ . In particular, we are not considering the bounding number \mathfrak{b}_κ , the splitting number \mathfrak{s}_κ , or the almost disjointness number \mathfrak{a}_κ , since, at least when generalized in accordance with the principles laid out in the previous paragraph, these cardinal characteristics are provably at most $\mathfrak{b}_{\text{cf}(\kappa)}$, $\mathfrak{s}_{\text{cf}(\kappa)}$, and $\mathfrak{a}_{\text{cf}(\kappa)}$, respectively (though we will have more to say about the almost disjointness number in Section 6).

A slightly suboptimal but succinct summary of our main results can be stated as follows (we refer the reader to Section 2 for the definition of *canonical function* and to Section 5 for the precise definition of the cardinal characteristics under consideration):

Main Corollary. *Suppose that*

- κ is a singular cardinal with $\text{cf}(\kappa) = \theta > \omega$ such that $\mu^\theta \leq \kappa$ for all $\mu < \kappa$;
- $\langle \kappa_i \mid i < \theta \rangle$ is an increasing, continuous sequence of cardinals converging to κ ;
- β is an ordinal for which the canonical function on θ of rank β , φ_β^θ , exists;
- $S \subseteq \theta$ is stationary;

- \mathfrak{cc} is one of the cardinal characteristics $m(\theta, \kappa)$, $d(\theta, \kappa)$, \mathfrak{r}_κ , or \mathfrak{d}_κ , and, for all $i < \theta$, \mathfrak{cc}_i is the corresponding cardinal characteristic $m(\text{cf}(\kappa_i), \kappa_i)$, $d(\text{cf}(\kappa_i), \kappa_i)$, \mathfrak{r}_{κ_i} , or \mathfrak{d}_{κ_i} ;
- for all $i \in S$, we have $\mathfrak{cc}_i \leq \kappa_i^{+\varphi_\beta^\theta(i)}$.

Then $\mathfrak{cc} \leq \kappa^{+\beta}$.

The slight suboptimality in this statement comes from the assumption that $\mu^\theta \leq \kappa$ for all $\mu < \kappa$. As we will see, something weaker than this is needed for our results; the precise weakening depends on the specific cardinal characteristic under consideration and will require some further notation to state, so we leave the exact statements for later sections.

The structure of the remainder of the paper is as follows. In Section 2, we review the definitions and facts regarding canonical functions and the Galvin-Hajnal rank that we will need for our results. In Section 3, we recall certain notions of *density*. This is important for two reasons: first, because the analogue of Silver's theorem for density numbers proven in [12] was direct inspiration for this paper, and secondly and more immediately, these density numbers will appear in the precise formulations of our results. After this, we begin with the proof of our main theorem. The proofs of our various analogues of the Galvin-Hajnal theorem all have the same general shape, so in Section 4 we develop an abstract framework that will apply to all of our specific instances. In Section 5, we apply this abstract framework to our cardinal characteristics under consideration to obtain our Main Theorem, which is precisely stated at the end of the section. Finally, in Section 6, we record some questions that remain open.

1.1. Notation and conventions: Unless otherwise noted, we believe our notation and terminology to be standard. We refer the reader to [11] for any undefined notions or notations from set theory, and we refer the reader to [2] for an introduction to cardinal characteristics of the continuum, generalizations of which form the subject of this paper.

If X is a set and \rightsquigarrow is a binary relation on X , then $\text{cf}(X, \rightsquigarrow)$ denotes the minimal cardinality of a subset $Y \subseteq X$ such that, for all $x \in X$, there is $y \in Y$ for which $x \rightsquigarrow y$. If θ is a regular uncountable cardinal, then NS_θ denotes the nonstationary ideal on θ . If $S \subseteq \theta$ is a stationary set, then, formally, $\text{NS}_\theta \upharpoonright S$ is the ideal on θ generated by $\text{NS}_\theta \cup \{\theta \setminus S\}$; in practice, we will typically think of $\text{NS}_\theta \upharpoonright S$ as the ideal of nonstationary subsets of S , considered as an ideal on S . If X is a set and κ is a cardinal, then $[X]^\kappa := \{y \subseteq X \mid |y| = \kappa\}$.

2. CANONICAL FUNCTIONS AND THE GALVIN-HAJNAL RANK

Suppose that S is an infinite set and I is a proper ideal on S . Given two functions $\varphi, \psi \in {}^S\text{On}$, we write $\varphi <_I \psi$ to denote the assertion that the set $\{i \in S \mid \psi(i) \leq \varphi(i)\}$ is in I . Define $=_I$, \leq_i , etc. in the obvious way. We will be particularly interested in the case in which S is a stationary subset of a regular uncountable cardinal θ and $I = \text{NS}_\theta \upharpoonright S$, i.e., I is the collection of nonstationary subsets of S . In this context, given two functions $\varphi, \psi \in {}^S\text{On}$, we will write $\varphi <_S \psi$ instead of $\varphi <_{\text{NS}_\theta \upharpoonright S} \psi$ (and similarly with \leq_S , $=_S$, etc.). In particular, for functions $\varphi, \psi \in {}^\theta\text{On}$, $\varphi <_\theta \psi$ will denote $\varphi <_{\text{NS}_\theta} \psi$. Note that $\varphi <_S \psi$ if and only if there is a club $C \subseteq \theta$ such that, for all $i \in C \cap S$, we have $\varphi(i) < \psi(i)$.

Fix for the remainder of this section a regular uncountable cardinal θ . Given a stationary set $S \subseteq \theta$, the corresponding relation $<_S$ is well-founded and therefore has a rank function, which yields what is known as the *Galvin-Hajnal rank*.

Definition 2.1 ([4]). Suppose that θ is an uncountable regular cardinal and $S \subseteq \theta$ is stationary. The *Galvin-Hajnal rank* of a function $\varphi \in {}^S\text{On}$, denoted $\|\varphi\|_S$, is defined by recursion on $<_c$ by letting

$$\|\varphi\|_S = \sup\{\|\psi\|_S + 1 \mid \psi \in {}^S\text{On} \text{ and } \psi <_c \varphi\}$$

for all $\varphi \in {}^S\text{On}$.

It is readily verified by recursion on $\|\varphi\|_S$ that, for all stationary $T \subseteq S \subseteq \theta$ and all $\varphi \in {}^S\text{On}$, we have $\|\varphi\|_S \leq \|\varphi \upharpoonright T\|_T$. In general, it is quite possible to have strict inequality here. However, if φ is what is known as a *canonical function*, this inequality is in fact always an equality. With this in mind, let us now recall the definition of and some basic facts about canonical functions.

By recursion on ordinals α , attempt to define the *canonical function on θ of rank α* , $\varphi_\alpha^\theta \in {}^\theta\text{On}$, as follows. If β is an ordinal and φ_α^θ has been defined for all $\alpha < \beta$, then let φ_β^θ be the least upper bound for $\langle \varphi_\alpha^\theta \mid \alpha < \beta \rangle$ with respect to $<_\theta$, if such a least upper bound exists. In other words, $\varphi_\beta^\theta \in {}^\theta\text{On}$ is a function such that

- φ_β^θ is a $<_\theta$ -upper bound for $\langle \varphi_\alpha^\theta \mid \alpha < \beta \rangle$;
- if ψ is another $<_\theta$ -upper bound for $\langle \varphi_\alpha^\theta \mid \alpha < \beta \rangle$, then $\varphi_\beta^\theta \leq_\theta \psi$.

If such a least upper bound does not exist, then φ_β^θ is undefined (and therefore φ_γ^θ is undefined for all $\gamma > \beta$ as well).

Note that φ_β^θ is not uniquely determined, but is unique up to $=_\theta$ -equivalence. We will let Φ_β^θ denote the set of all canonical functions on θ of rank β . We will slightly abuse notation and use φ_β^θ to denote an arbitrary element of Φ_β^θ . We will always be working in contexts that are invariant under $=_\theta$ -equivalence, so this will not result in any loss of generality. The following well-known fact (see [14, §1] for an introduction to canonical functions of rank less than θ^+) shows that, for all $\beta < \theta^+$, there are canonical functions on θ of rank β .

Fact 2.2. *Let $\beta < \theta^+$, and let $e : \theta \rightarrow \beta$ be a surjection. Then the function $\varphi \in {}^\theta\theta$ defined by letting $f(i) = \text{otp}(e^{-1}i)$ for all $i < \theta$ is in Φ_β^θ .*

The following proposition follows almost immediately from the definition of *canonical function*.

Proposition 2.3. *Suppose that β is an ordinal for which φ_β^θ is defined, $\psi \in {}^\theta\text{On}$, and the set $S := \{i < \theta \mid \psi(i) < \varphi_\beta^\theta(i)\}$ is stationary in θ . Then there is a stationary $S' \subseteq S$ and an $\alpha < \beta$ such that $\psi(i) \leq \varphi_\alpha^\theta(i)$ for all $i \in S'$.*

Proof. Suppose not. Then, for all $\alpha < \beta$, there is a club $C_\alpha \subseteq \theta$ such that $\varphi_\alpha^\theta(i) < \psi(i)$ for all $i \in S \cap C_\alpha$. Define a function $\tau \in {}^\theta\text{On}$ by letting

$$\tau(i) = \begin{cases} \psi(i) & \text{if } i \in S \\ \varphi_\beta^\theta(i) & \text{if } i \in \theta \setminus S. \end{cases}$$

Then, by our assumptions, τ is a $<_\theta$ -upper bound for $\langle \varphi_\alpha^\theta \mid \alpha < \beta \rangle$, so, by the definition of *canonical function* we must have $\varphi_\beta^\theta \leq_\theta \tau$, contradicting the fact that $S \subseteq \theta$ is stationary and, for all $i \in S$, we have $\tau(i) = \psi(i) < \varphi_\beta^\theta(i)$. \square

The following basic facts will be relevant to our arguments.

Proposition 2.4. *Suppose that $\beta > 0$ is an ordinal such that φ_β^θ is defined.*

- (1) $\varphi_{\beta+1}^\theta$ is defined and $\varphi_{\beta+1}^\theta =_\theta \varphi_\beta^\theta + 1$.
- (2) If β is a limit ordinal, then there is a club $C \subseteq \theta$ such that $\varphi_\beta^\theta(i)$ is a limit ordinal for all $i \in C$.

Proof. (1) Clearly, $\varphi_\beta^\theta + 1$ is a $<_\theta$ -upper bound for $\langle \varphi_\alpha^\theta \mid \alpha \leq \beta \rangle$. Moreover, if ψ is any other $<_\theta$ -upper bound, then there must be a club $C \subseteq \theta$ such that $\varphi_\beta^\theta(i) + 1 \leq \psi(i)$ for all $i \in C$, and therefore $\varphi_\beta^\theta + 1 <_\theta \psi$. It follows that $\varphi_{\beta+1}^\theta =_\theta \varphi_\beta^\theta + 1$.

(2) Suppose for sake of contradiction that β is a limit ordinal and yet there is a stationary set $S \subseteq \theta$ such that $\varphi_\beta^\theta(i) = \gamma_i + 1$ is a successor ordinal for all $i \in S$. Define a function $\psi \in {}^\theta\text{On}$ by letting

$$\psi(i) = \begin{cases} \gamma_i & \text{if } i \in S \\ \varphi_\beta^\theta(i) & \text{otherwise} \end{cases}$$

for all $i < \theta$. By Proposition 2.3, we can find a stationary $S' \subseteq S$ and an ordinal $\alpha < \beta$ such that $\psi(i) \leq \varphi_\alpha^\theta(i)$ for all $i \in S'$, and hence, by removing a nonstationary subset from S' if necessary, we can assume that $\psi(i) + 1 \leq \varphi_{\alpha+1}^\theta(i)$ for all $i \in S'$. But, by our definition of ψ , we have $\psi(i) + 1 = \varphi_\beta^\theta(i)$ for all $i \in S'$, and hence $\varphi_{\alpha+1}^\theta \not<_\theta \varphi_\beta^\theta$, contradicting the fact that $\alpha + 1 < \beta$. \square

Proposition 2.5. *Suppose that θ is a regular uncountable cardinal, β is an ordinal such that φ_β^θ is defined, and $S \subseteq \theta$ is stationary. Then $\|\varphi_\beta^\theta \upharpoonright S\|_S = \beta$.*

Proof. The proof is by induction on β , so we assume that, for all $\alpha < \beta$ and all stationary $T \subseteq \theta$, we have $\|\varphi_\alpha^\theta \upharpoonright T\|_T = \alpha$. Since $\varphi_\alpha^\theta <_\theta \varphi_\beta^\theta$ for all $\alpha < \beta$, it follows that $\|\varphi_\beta^\theta \upharpoonright S\|_S \geq \beta$.

For the opposite inequality, fix a function $\psi \in {}^S\text{On}$ with $\psi <_S \varphi_\beta^\theta \upharpoonright S$; it suffices to show that $\|\psi\|_S < \beta$. An application of Proposition 2.3 yields a stationary $T \subseteq S$ and an $\alpha < \beta$ such that $\psi(i) \leq \varphi_\alpha^\theta(i)$ for all $i \in T$. By the induction hypothesis, we have $\|\varphi_\alpha^\theta \upharpoonright T\|_T = \alpha$, so it follows that $\|\psi \upharpoonright T\|_T \leq \alpha$. But then, since $T \subseteq S$, this implies that $\|\psi\|_S \leq \alpha < \beta$, as desired. \square

We now recall two results from [10], the first of which is already implicit in [17].

Theorem 2.6. [10, Corollary 2.3] *Suppose that A is an infinite set, I is an ideal on A , and $\langle \mu_a \mid a \in A \rangle$ is a sequence of regular cardinals such that $\mu_a > |A|^+$ for all $a \in A$. Then there exist a set $B \in I^+$, a regular cardinal $\lambda > |A|^+$, and a sequence $\vec{f} = \langle f_\alpha \mid \alpha < \lambda \rangle$ such that \vec{f} is $<_{I \upharpoonright B}$ -increasing and $<_{I \upharpoonright B}$ -cofinal in $\prod_{a \in B} \mu_a$.*

Our statement of the next theorem is less general than its statement in [10]; we focus on ideals of the form $\text{NS}_\theta \upharpoonright S$ rather than the arbitrary normal ideals of [10].

Theorem 2.7. [10, Main Theorem] *Suppose that*

- (1) κ is a singular cardinal and $\text{cf}(\kappa) = \theta > \omega$;
- (2) $\langle \kappa_i \mid i < \theta \rangle$ is an increasing, continuous sequence of cardinals converging to κ ;
- (3) $\langle \mu_i \mid i < \theta \rangle$ is an increasing sequence of regular cardinals such that, for some function $\varphi \in {}^\theta\text{On}$, we have $\mu_i = \kappa_i^{+\varphi(i)}$ for all $i < \theta$;
- (4) $S \subseteq \theta$ is stationary;

(5) λ is a regular cardinal and $\vec{f} = \langle f_\alpha \mid \alpha < \lambda \rangle$ is a $<_S$ -increasing and $<_S$ -cofinal sequence from $\prod_{i \in S} \mu_i$.

Then $\lambda \leq \kappa^{+\|\varphi \upharpoonright S\|_S}$.

Putting these two results together yields the following corollary.

Corollary 2.8. *Suppose that θ is a regular uncountable cardinal, $S \subseteq \theta$ is stationary, and β is an ordinal such that φ_β^θ is defined. Suppose also that κ is a singular cardinal, $\text{cf}(\kappa) = \theta$, and $\langle \kappa_i \mid i < \theta \rangle$ is an increasing, continuous sequence of cardinals converging to κ with $\kappa_0 > \theta$. Then there is a stationary $S' \subseteq S$ and a sequence $\vec{f} = \langle f_\alpha \mid \alpha < \lambda \rangle$ from $\prod_{i \in S'} \kappa_i^{+\varphi_\beta^\theta(i)+1}$ such that*

- (1) \vec{f} is $<_{S'}$ -increasing and $<_{S'}$ -cofinal in $\prod_{i \in S'} \kappa_i^{+\varphi_\beta^\theta(i)+1}$;
- (2) $\lambda \leq \kappa^{+\beta+1}$.

In particular, there exists a $<_{S'}$ -cofinal subset $\mathcal{F} \subseteq \prod_{i \in S'} \kappa_i^{+\varphi_\beta^\theta(i)+1}$ such that $|\mathcal{F}| \leq \kappa^{+\beta+1}$.

Proof. By Proposition 2.4, we have $\varphi_{\beta+1}^\theta = \varphi_\beta^\theta + 1$. For each $i \in S$, let $\mu_i := \kappa_i^{+\varphi_\beta^\theta(i)+1}$. Then, applying Theorem 2.6 to the set S , the ideal $\text{NS}_\theta \upharpoonright S$, and the sequence $\langle \mu_i \mid i \in S \rangle$ of regular cardinals, we obtain a stationary $S' \subseteq S$, a regular cardinal $\lambda > |A|^+$, and a sequence $\vec{f} = \langle f_\alpha \mid \alpha < \lambda \rangle$ such that \vec{f} is $<_{S'}$ -increasing and $<_{S'}$ -cofinal in $\prod_{i \in S'} \mu_i$. Then Theorem 2.7 implies that $\lambda \leq \kappa^{+\|\varphi_{\beta+1}^\theta \upharpoonright S'\|_{S'}}$, so, by Proposition 2.5, we have $\lambda \leq \kappa^{+\beta+1}$. \square

3. DENSITY

In this section, we recall some notions of *density* that will play a role throughout the paper. The first of these notions was the subject of Kojman's [12].

Definition 3.1. [12] Suppose that $\theta \leq \mu$ are infinite cardinals. The θ -density of μ , denoted $d(\theta, \mu)$, is the minimal cardinality of a set $\mathcal{Y} \subseteq [\mu]^\theta$ that is dense in $([\mu]^\theta, \subseteq)$.

If $\theta < \mu$, then $\underline{[\mu]}^\theta$ denotes the set of $x \in [\mu]^\theta$ such that $\sup(x) < \mu$, and the lower θ -density of μ , denoted $\underline{d}(\theta, \mu)$, is the minimal cardinality of a set $\mathcal{Y} \subseteq \underline{[\mu]}^\theta$ that is dense in $(\underline{[\mu]}^\theta, \subseteq)$.

Remark 3.2. In [12], the θ -density of μ is denoted by $\mathcal{D}(\mu, \theta)$. We have chosen the notation $d(\theta, \mu)$ to match the established notation $m(\theta, \mu)$ for *meeting numbers*, which are among the cardinal characteristics considered here. Our notation for both density numbers and meeting numbers follows [16].

Note that, if $\theta < \mu$, then $\underline{d}(\theta, \mu) = \mu \cdot \sum_{\nu < \mu} d(\theta, \nu)$. Therefore, if $\nu^\theta \leq \mu$ for all $\nu < \mu$, then $\underline{d}(\theta, \mu) = \mu$.

As remarked in [12], if $\text{cf}(\mu) \neq \text{cf}(\theta)$, then

$$d(\theta, \mu) = \underline{d}(\theta, \mu) = \mu \cdot \sum_{\nu < \mu} d(\theta, \nu).$$

In particular, if $\text{cf}(\mu) \neq \text{cf}(\theta)$ and $\nu^\theta \leq \mu$ for all $\nu < \mu$, then $d(\theta, \mu) = \mu$.

If $\text{cf}(\mu) = \text{cf}(\theta)$, then a routine diagonalization argument shows that $d(\theta, \mu) \geq \mu^+$.

The main result of [12] is a version of Silver’s theorem for the density number $d(\text{cf}(\kappa), \kappa)$; this result served as direct motivation for the initial work that led to the results of this paper. Our main result here, when applied to the density number, will generalize and slightly improve upon the results of [12].

If I is an ideal over a set X , then the *density* of I , which we will denote $d(I)$, is the minimal cardinality of a set $\mathcal{Y} \subseteq I^+$ such that, for all $S \in I^+$, there is $T \in \mathcal{Y}$ such that $T \setminus S \in I$. We will particularly be interested in densities of the form $d(\text{NS}_\theta \upharpoonright S)$, where S is a stationary set of a regular uncountable cardinal θ . Concretely, $d(\text{NS}_\theta \upharpoonright S)$ is the minimal cardinality of a collection \mathcal{T} of stationary subsets of S such that, for every stationary $S' \subseteq S$, there is $T \in \mathcal{T}$ such that $T \setminus S'$ is nonstationary in θ .

Finally, we introduce a notion of density that, in a sense, combines the two notions introduced in this section thus far.

Definition 3.3. Suppose that θ and κ are infinite cardinals, with θ regular, and suppose that $S \subseteq \theta$ is stationary. Then the *stationarity density of S_κ* , which we denote by $d_{\text{stat}}(S_\kappa)$, is the minimal cardinality of a family \mathcal{F} of functions such that

- (1) every $f \in \mathcal{F}$ is a function from a stationary subset of S to κ ;
- (2) for every function g from a stationary subset of S to κ , there is $f \in \mathcal{F}$ such that the set

$$\{i \in \text{dom}(f) \mid i \notin \text{dom}(g) \text{ or } f(i) \neq g(i)\}$$

is nonstationary. (Less precisely but more evocatively, f is contained in g modulo a nonstationary set.)

In analogy with lower density, we define the *lower stationary density of S_κ* , denoted $\underline{d}_{\text{stat}}(S_\kappa)$, in the same way as $d_{\text{stat}}(S_\kappa)$, except that, in item (2), we only consider functions whose ranges are bounded below κ (and hence we can require that all of our functions in \mathcal{F} also have ranges bounded below κ).

Remark 3.4. Note that $d(\text{NS}_\theta \upharpoonright S) \leq \underline{d}_{\text{stat}}(S_\kappa)$. Also, whenever $T \subseteq S$ are stationary subsets of θ , we have

- $d(\text{NS}_\theta \upharpoonright T) \leq d(\text{NS}_\theta \upharpoonright S)$;
- $\underline{d}_{\text{stat}}(T_\kappa) \leq \underline{d}_{\text{stat}}(S_\kappa)$;
- $\underline{d}_{\text{stat}}(T_\kappa) \leq \underline{d}_{\text{stat}}(S_\kappa)$.

4. THE GENERAL FRAMEWORK

The cardinal characteristics we consider in this paper all have the following form: a set X and a binary relation \rightsquigarrow on X are fixed, and the relevant cardinal characteristic is then $\text{cf}(\mathcal{X}, \rightsquigarrow)$, i.e., the minimal cardinality of a subset $\mathcal{Y} \subseteq \mathcal{X}$ such that, for all $x \in \mathcal{X}$, there is $y \in \mathcal{Y}$ such that $x \rightsquigarrow y$.

Because of the structural similarity of these cardinal characteristics, the inductive steps in the proofs of our main results end up being essentially the same, so in this section we prove a general lemma that we can directly apply to all of the specific situations under consideration here, and that we expect will find application beyond the scope of this paper, as well.

In order to state and prove our general lemma, let us fix some objects and notation for the remainder of this section:

- κ is a singular cardinal and $\text{cf}(\kappa) = \theta > \omega$;

- $\langle \kappa_i \mid i < \theta \rangle$ is an increasing, continuous sequence of cardinals converging to κ , with $\kappa_0 > \theta$;
- \mathcal{X} is a set and \rightsquigarrow is a binary relation on \mathcal{X} ;
- for each $i < \theta$, \mathcal{X}_i is a set and \rightsquigarrow_i is a binary relation on \mathcal{X}_i ;
- for each $i < \theta$, $\pi_i : \mathcal{X} \rightarrow \mathcal{X}_i$ is a function;
- $e : \text{NS}_\theta^+ \rightarrow \text{Card}$ is a function such that, for all stationary $T \subseteq S \subseteq \theta$, we have $e(T) \leq e(S)$.

Remark 4.1. To help orient the reader, let us preview here some typical interpretations of these objects. In a typical application, we might have $\mathcal{X} = \mathcal{P}(\kappa)$, $\mathcal{X}_i = \mathcal{P}(\kappa_i)$, and $\pi_i(x) = x \cap \kappa_i$, or $\mathcal{X} = {}^\kappa\kappa$, $\mathcal{X}_i = {}^{\kappa_i}\kappa_i$, and $\pi_i(x)$ is a modification of $x \upharpoonright \kappa_i$ to ensure that it takes values in κ_i . The function e will typically (though not always) output one of the density numbers introduced in Section 3.

In this context, if $S \subseteq \theta$ is stationary and β is an ordinal for which the canonical function φ_β^θ is defined, then let $\Psi(S, \beta)$ denote the following assertion:

If \mathcal{Z} and $\langle \mathcal{Y}_i \mid i \in S \rangle$ are such that

- (1) $\mathcal{Z} \subseteq \mathcal{X}$ and $\mathcal{Y}_i \subseteq \mathcal{X}_i$ for all $i \in S$;
- (2) $|\mathcal{Y}_i| \leq \kappa_i^{+\varphi_\beta^\theta(i)}$ for all $i \in S$;
- (3) for all $z \in \mathcal{Z}$, there is a club $C \subseteq \theta$ such that, for all $i \in C \cap S$, there is $y \in \mathcal{Y}_i$ for which $\pi_i(z) \rightsquigarrow_i y$;

then there is $\mathcal{Y} \subseteq \mathcal{X}$ such that $|\mathcal{Y}| \leq \kappa^{+\beta} + d(\text{NS}_\theta \upharpoonright S) + e(S)$ and, for all $z \in \mathcal{Z}$, there is $y \in \mathcal{Y}$ for which $z \rightsquigarrow y$.

Let $\Psi^*(S, \beta)$ be defined in the same way, except, in the conclusion, we only require $|\mathcal{Y}| \leq \kappa^{+\beta} + e(S)$.

Lemma 4.2. *Suppose that $S \subseteq \theta$ is stationary and $\Psi(T, 0)$ holds for all stationary $T \subseteq S$. Then, for all ordinals β for which φ_β^θ is defined, $\Psi(S, \beta)$ holds.*

Proof. The proof is by induction on β , simultaneously for all stationary $S \subseteq \theta$. Thus, fix an ordinal β for which φ_β^θ is defined and a stationary set $S \subseteq \theta$. By the hypothesis of the lemma, we can assume that $\beta > 0$, and by the inductive hypothesis, we can assume that $\Psi(T, \alpha)$ holds for all $\alpha < \beta$ and all stationary $T \subseteq S$. Fix \mathcal{Z} and $\langle \mathcal{Y}_i \mid i \in S \rangle$ as in the hypothesis of $\Psi(S, \beta)$; we will find $\mathcal{Y} \subseteq \mathcal{X}$ as in its conclusion. For each $i \in S$, enumerate \mathcal{Y}_i as $\langle y_{i,\xi} \mid \xi < \kappa_i^{+\varphi_\beta^\theta(i)} \rangle$ (with repetitions if $|\mathcal{Y}_i| < \kappa_i^{+\varphi_\beta^\theta(i)}$).

Suppose first that $\beta = \beta' + 1$ is a successor ordinal. By Proposition 2.4, we can assume that $\varphi_\beta^\theta = \varphi_{\beta'}^\theta + 1$. Then, by Corollary 2.8, we can fix a stationary $S' \subseteq S$ and a $<_{S'}$ -cofinal family $\mathcal{F} \subseteq \prod_{i \in S'} \kappa_i^{+\varphi_{\beta'}^\theta(i)}$ such that $|\mathcal{F}| \leq \kappa^{+\beta}$. For each $f \in \mathcal{F}$, let $\mathcal{Y}_{i,f} := \{y_{i,\xi} \mid \xi < f(i)\}$, and note that $|\mathcal{Y}_{i,f}| \leq \kappa_i^{+\varphi_{\beta'}^\theta(i)}$. Let \mathcal{Z}_f be the set of $z \in \mathcal{Z}$ for which there is a club $C \subseteq \theta$ such that, for all $i \in C \cap S'$, there is $y \in \mathcal{Y}_{i,f}$ such that $\pi_i(z) \rightsquigarrow_i y$. Recalling that $d(\text{NS}_\theta \upharpoonright S') \leq d(\text{NS}_\theta \upharpoonright S)$ and $e(S') \leq e(S)$, apply $\Psi(S', \beta')$ to \mathcal{Z}_f and $\langle \mathcal{Y}_{i,f} \mid i \in S' \rangle$ to find $\mathcal{Y}_f \subseteq \mathcal{X}$ such that $|\mathcal{Y}_f| \leq \kappa^{+\beta'} + d(\text{NS}_\theta \upharpoonright S) + e(S)$ and, for all $z \in \mathcal{Z}_f$, there is $y \in \mathcal{Y}_f$ such that $z \rightsquigarrow y$.

Let $\mathcal{Y} = \bigcup_{f \in \mathcal{F}} \mathcal{Y}_f$; we claim that \mathcal{Y} is as desired. It is evident that $\mathcal{Y} \subseteq \mathcal{X}$ and $|\mathcal{Y}| \leq \kappa^{+\beta} + d(\text{NS}_\theta \upharpoonright S) + e(S)$, so it remains to verify that, for all $z \in \mathcal{Z}$, there is $y \in \mathcal{Y}$ such that $z \rightsquigarrow y$. For this, it suffices to show that $\mathcal{Z} \subseteq \bigcup_{f \in \mathcal{F}} \mathcal{Z}_f$. To this

end, fix $z \in \mathcal{Z}$. By assumption, there is a club $C \subseteq \theta$ such that, for all $i \in C \cap S'$, there is $\xi_i < \kappa_i^{+\varphi_\beta^\theta(i)}$ for which $\pi_i(z) \rightsquigarrow_i y_{i,\xi_i}$. Define a function $g \in \prod_{i \in S'} \kappa_i^{+\varphi_\beta^\theta(i)}$ by letting

$$g(i) = \begin{cases} \xi_i & \text{if } i \in C \\ 0 & \text{otherwise} \end{cases}$$

for all $i \in S'$. Since \mathcal{F} is $<_c$ -cofinal in $\prod_{i \in S'} \kappa_i^{+\varphi_\beta^\theta(i)}$, we can find $f \in \mathcal{F}$ such that $g <_C f$, i.e., there is a club $D \subseteq \theta$ such that, for all $i \in D \cap S'$, we have $g(i) < f(i)$. But then, for all $i \in D \cap C \cap S'$, we have $\xi_i < f(i)$ and $\pi_i(z) \rightsquigarrow_i y_{i,\xi_i}$, so $D \cap C$ witnesses that z is in \mathcal{Z}_f , and we are done.

Finally, suppose that β is a limit ordinal. By Proposition 2.4, we can assume that $\varphi_\beta^\theta(i)$ is a limit ordinal for all $i \in S$. Let \mathcal{T} be a collection of stationary subsets of S such that $|\mathcal{T}| = d(\text{NS}_\theta \upharpoonright S)$ and, for every stationary $S' \subseteq S$, there is $T \in \mathcal{T}$ such that $T \setminus S' \in \text{NS}_\theta$. For each $\alpha < \beta$ and each $i \in S$, let $\mathcal{Y}_i^\alpha := \{y_{i,\xi} \mid \xi < \kappa_i^{+\varphi_\alpha^\theta(i)}\}$. For each $\alpha < \beta$ and each $T \in \mathcal{T}$, let $\mathcal{Z}_{T,\alpha}$ be the set of all $z \in \mathcal{Z}$ for which there is a club $C \subseteq \theta$ such that, for all $i \in C \cap T$, there is $y \in \mathcal{Y}_i^\alpha$ such that $\pi_i(z) \rightsquigarrow_i y$. Apply $\Psi(T, \alpha)$ to $\mathcal{Z}_{T,\alpha}$ and $\langle \mathcal{Y}_i^\alpha \mid i \in T \rangle$ to find $\mathcal{Y}_{T,\alpha} \subseteq \mathcal{X}$ such that $|\mathcal{Y}_{T,\alpha}| \leq \kappa^{+\alpha} + d(\text{NS}_\theta \upharpoonright S) + e(S)$ and, for all $z \in \mathcal{Z}_{T,\alpha}$, there is $y \in \mathcal{Y}_{T,\alpha}$ such that $z \rightsquigarrow y$.

Let $\mathcal{Y} = \bigcup \{\mathcal{Y}_{T,\alpha} \mid T \in \mathcal{T}, \alpha < \beta\}$; we claim that \mathcal{Y} is as desired. As in the successor case, it suffices to verify that $\mathcal{Z} \subseteq \bigcup \{\mathcal{Z}_{T,\alpha} \mid T \in \mathcal{T}, \alpha < \beta\}$. To this end, fix $z \in \mathcal{Z}$. By hypothesis, we can find a club $C \subseteq \theta$ such that, for all $i \in C \cap S$, there is $\xi_i < \kappa_i^{+\varphi_\beta^\theta(i)}$ for which $\pi_i(z) \rightsquigarrow_i y_{i,\xi_i}$. For each such i , use the fact that $\varphi_\beta^\theta(i)$ is a limit ordinal to find $\gamma_i < \varphi_\beta^\theta(i)$ such that $\xi_i < \kappa_i^{+\gamma_i}$. Define a function $\psi \in {}^\theta \text{On}$ by letting

$$\psi(i) = \begin{cases} \gamma_i & \text{if } i \in C \cap S \\ \varphi_\beta^\theta(i) & \text{otherwise} \end{cases}$$

for all $i < \theta$. By Proposition 2.3, we can find a stationary $S' \subseteq C \cap S$ and an $\alpha < \beta$ such that $\gamma_i = \psi(i) \leq \varphi_\alpha^\theta(i)$ for all $i \in S'$. We can subsequently find a $T \in \mathcal{T}$ and a club $D \subseteq \theta$ such that $D \cap T \subseteq S'$. Then, for all $i \in D \cap T$, we have $y_{i,\xi_i} \in \mathcal{Y}_i^\alpha$ and $\pi_i(z) \rightsquigarrow_i y_{i,\xi_i}$, so D witnesses that z is in $\mathcal{Z}_{T,\alpha}$, and we are done. \square

Notice that the only place in which the value of $d(\text{NS}_\theta \upharpoonright S)$ plays a role in the proof of Lemma 4.2 is in the case in which β is a limit ordinal (in the successor case it only makes an appearance via the inductive hypothesis). Therefore, if $\beta < \omega$ and $\Psi^*(T, 0)$ holds for all stationary $T \subseteq \theta$, then we can do away with $d(\text{NS}_\theta \upharpoonright S)$ in the conclusion of the lemma. More precisely, the proof of the successor case of Lemma 4.2 yields the following corollary.

Corollary 4.3. *Suppose that $S \subseteq \theta$ is stationary and $\Psi^*(T, 0)$ holds for all stationary $T \subseteq S$. Then $\Psi^*(S, n)$ holds for all $n < \omega$.*

The translation from Lemma 4.2 and Corollary 4.3 to our main results will happen via the following corollary.

Corollary 4.4. *Suppose that*

- (1) β is an ordinal for which φ_β^θ is defined;
- (2) $S := \{i < \theta \mid \text{cf}(X_i, \rightsquigarrow_i) \leq \kappa_i^{+\varphi_\beta^\theta(i)}\}$ is stationary in θ ;

(3) $\Psi(T, 0)$ holds for all stationary $T \subseteq S$.

Then $\text{cf}(X, \rightsquigarrow) \leq \kappa^{+\beta} + d(\text{NS}_\theta \upharpoonright S) + e(S)$. If, moreover, $\beta < \omega$ and $\Psi^*(T, 0)$ holds for every stationary $T \subseteq S$, then $\text{cf}(X, \rightsquigarrow) \leq \kappa^{+\beta} + e(S)$.

Proof. By Lemma 4.2, we know that $\Psi(S, \beta)$ holds. Let $\mathcal{Z} = \mathcal{X}$ and, for each $i \in S$, let $\mathcal{Y}_i \subseteq \mathcal{X}_i$ be such that $|\mathcal{Y}_i| \leq \kappa_i^{+\varphi_\beta^\theta(i)}$ and \mathcal{Y}_i is \rightsquigarrow_i -cofinal in \mathcal{X}_i . In particular, for all $z \in \mathcal{Z}$ and all $i \in S$, there is $y \in \mathcal{Y}_i$ such that $\pi_i(z) \rightsquigarrow_i y$. Therefore, applying $\Psi(S, \beta)$ to \mathcal{Z} and $\langle \mathcal{Y}_i \mid i \in S \rangle$ yields a set $\mathcal{Y} \subseteq \mathcal{X}$ such that $|\mathcal{Y}| \leq \kappa^{+\beta} + d(\text{NS}_\theta \upharpoonright S) + e(S)$ and, for all $z \in \mathcal{Z}$, there is $y \in \mathcal{Y}$ such that $z \rightsquigarrow y$, i.e., \mathcal{Y} is \rightsquigarrow -cofinal in \mathcal{X} .

For the “moreover” clause, if $\beta < \omega$ and $\Psi^*(T, 0)$ holds for every stationary $T \subseteq S$, then Corollary 4.3 implies that $\Psi^*(S, \beta)$ holds. Applying $\Psi^*(S, \beta)$ to the \mathcal{Z} and $\langle \mathcal{Y}_i \mid i \in S \rangle$ of the previous paragraph then yields $\mathcal{Y} \subseteq \mathcal{X}$ such that $|\mathcal{Y}| \leq \kappa^{+\beta} + e(S)$ and \mathcal{Y} is \rightsquigarrow -cofinal in \mathcal{X} . \square

5. SPECIFIC INSTANCES

We now turn to applications of the general framework introduced in the previous section to particular cardinal characteristics at singular cardinals. Let us fix for this entire section cardinals κ and θ such that $\omega < \theta = \text{cf}(\kappa) < \kappa$, as well as an increasing, continuous sequence of cardinals $\langle \kappa_i \mid i < \theta \rangle$ converging to κ , with $\kappa_0 > \theta$.

Each cardinal characteristic \mathfrak{cc}_κ we consider will entail a choice of a set \mathcal{X} and a binary relation \rightsquigarrow on \mathcal{X} such that $\mathfrak{cc}_\kappa = \text{cf}(\mathcal{X}, \rightsquigarrow)$. The sets $\langle \mathcal{X}_i \mid i < \theta \rangle$ and relations $\langle \rightsquigarrow_i \mid i < \theta \rangle$ will be defined analogously, so that $\mathfrak{cc}_{\kappa_i} = \text{cf}(\mathcal{X}_i, \rightsquigarrow_i)$ for each $i < \theta$. We will also have natural restriction operations $\pi_i : \mathcal{X} \rightarrow \mathcal{X}_i$ for each $i < \theta$. Together with an appropriate choice of function $e : \text{NS}_\theta^+ \rightarrow \text{Card}$, these assignments give rise to instances of the formulas $\Psi(S, \beta)$ and $\Psi^*(S, \beta)$ from the previous section. The primary work of this section will consist of proving that all of these instances of $\Psi^*(T, 0)$ hold; Corollary 4.4 will then directly yield our main results.

We will begin by introducing each of the cardinal characteristics we will be considering and specifying the appropriate assignments for \mathcal{X} , \rightsquigarrow , $\langle (\mathcal{X}_i, \rightsquigarrow_i, \pi_i) \mid i < \theta \rangle$, and e for each characteristic. We will then prove a lemma indicating that, for all of these assignments, the corresponding instance of $\Psi^*(T, 0)$ holds for every stationary $T \subseteq \theta$.

We note that for some of the cardinal characteristics, we will only define \mathcal{X}_i , \rightsquigarrow_i , and π_i for limit ordinals $i < \theta$. Since the clubs C in the statement of $\Psi(S, \beta)$ can always be assumed to consist entirely of limit ordinals, this is sufficient for our purposes.

5.1. The meeting number. The first cardinal characteristics we consider are the *meeting numbers*.

Definition 5.1 ([16]). Suppose that $\sigma \leq \lambda$ are infinite cardinals. Then the *meeting number* $m(\sigma, \lambda)$ is the minimal cardinality of a collection $\mathcal{Y} \subseteq [\lambda]^\sigma$ such that, for all $x \in [\lambda]^\sigma$, there is $y \in \mathcal{Y}$ such that $|x \cap y| = \sigma$.

The meeting number $m(\sigma, \lambda)$ is of special interest when $\text{cf}(\sigma) = \text{cf}(\lambda)$, in which case a routine diagonalization argument implies that $m(\sigma, \lambda) > \lambda$. A result of Matet

indicates that Shelah's Strong Hypothesis, a statement in PCF theory, is equivalent to the statement that all such meeting numbers take their minimal possible value:

Theorem 5.2 (Matet, [15, Theorem 1.1]). *The following are equivalent:*

- (1) *Shelah's Strong Hypothesis;*
- (2) *for every singular cardinal λ of countable cofinality, $m(\aleph_0, \lambda) = \lambda^+$;*
- (3) *for all infinite cardinals $\sigma < \lambda$, we have $m(\sigma, \lambda) = \lambda^+$ if $\text{cf}(\sigma) = \text{cf}(\lambda)$ and $m(\sigma, \lambda) = \lambda$ if $\text{cf}(\sigma) \neq \text{cf}(\lambda)$.*

We now specify assignments to define a version of the formula $\Psi^*(T, 0)$ appropriate for the meeting number. Let $\mathcal{X} = [\kappa]^\theta$ and, for each limit ordinal $i < \theta$, let $\mathcal{X}_i = [\kappa_i]^{\text{cf}(i)}$. Define relations \rightsquigarrow on \mathcal{X} and \rightsquigarrow_i on \mathcal{X}_i by letting $x \rightsquigarrow y$ iff $|x \cap y| = \theta$ and $x \rightsquigarrow_i y$ iff $|x_i \cap y_i| = \text{cf}(i)$ for all limit ordinals $i < \theta$. Then it is evident that $m(\theta, \kappa) = \text{cf}(\mathcal{X}, \rightsquigarrow)$ and $m(\text{cf}(i), \kappa_i) = m(\text{cf}(\kappa_i), \kappa_i) = \text{cf}(\mathcal{X}_i, \rightsquigarrow_i)$ for all limit $i < \theta$.

For each limit ordinal $i < \theta$, define a map $\pi_i : \mathcal{X} \rightarrow \mathcal{X}_i$ as follows. For any $i < \theta$ and $x \in \mathcal{X}$, if $\sup(x \cap \kappa_i) = \kappa_i$, then let $\pi_i(x)$ be an arbitrary unbounded subset of $x \cap \kappa_i$ of order type $\text{cf}(i)$. If $\sup(x \cap \kappa_i) < \kappa_i$, then simply let $\pi_i(x)$ be an arbitrary element of \mathcal{X}_i . Note that, for every $x \in \mathcal{X}$ that is unbounded in κ , the set $\{i < \theta \mid \sup(x \cap \kappa_i) = \kappa_i\}$ is a club in θ .

Let $e : \text{NS}_\theta^+ \rightarrow \text{Card}$ be the constant function taking value $\sum_{j < \theta} m(\theta, \kappa_j)$.

5.2. Density. As mentioned above, a version of Silver's theorem for density is proven in [12]. We include density here for completeness, since our results are slightly more general than those of [12].

We are interested in particular in the number $d(\theta, \kappa)$. The setup for density will be similar to that for the meeting number. Again let $\mathcal{X} = [\kappa]^\theta$ and, for each limit ordinal $i < \theta$, let $\mathcal{X}_i = [\kappa_i]^{\text{cf}(i)}$. Define relations \rightsquigarrow on \mathcal{X} and \rightsquigarrow_i on \mathcal{X}_i by letting $x \rightsquigarrow y$ or $x \rightsquigarrow_i y$ iff $x \supseteq y$. Then $d(\theta, \kappa) = \text{cf}(\mathcal{X}, \rightsquigarrow)$ and $d(\text{cf}(\kappa_i), \kappa_i) = \text{cf}(\mathcal{X}_i, \rightsquigarrow_i)$ for all limit $i < \theta$.

Define maps $\pi_i : \mathcal{X} \rightarrow \mathcal{X}_i$ for limit ordinals $i < \theta$ exactly as in the case of the meeting number in the previous subsection. Let $e : \text{NS}_\theta^+ \rightarrow \text{Card}$ be the constant function taking value $\underline{d(\theta, \kappa)} = \sum_{j < \theta} d(\theta, \kappa_j)$ (recall Definition 3.1).

5.3. The reaping number.

Definition 5.3. Let λ be an infinite cardinal.

- (1) If $x, y \in [\lambda]^\lambda$, then we say that x *splits* y if $|y \cap x| = |y \setminus x| = \lambda$.
- (2) A family $\mathcal{Y} \subseteq [\lambda]^\lambda$ is *unreaped* if there is no single $x \in [\lambda]^\lambda$ that splits every element of \mathcal{Y} .
- (3) The reaping number \mathfrak{r}_λ is the minimum cardinality of an unreaped family in $[\lambda]^\lambda$.

A standard diagonalization argument shows that $\mathfrak{r}_\lambda > \lambda$ for every infinite cardinal λ .

Let $\mathcal{X} = [\kappa]^\kappa$ and, for each $i < \theta$, let $\mathcal{X}_i = [\kappa_i]^{\kappa_i}$. Define a relation \rightsquigarrow on \mathcal{X} by letting $x \rightsquigarrow y$ iff x does not split y , i.e., either $|y \cap x| < \kappa$ or $|y \setminus x| < \kappa$. Similarly, for each $i < \theta$, define \rightsquigarrow_i on \mathcal{X}_i by letting $x \rightsquigarrow_i y$ iff x does not split y . Then it is evident that $\mathfrak{r}_\kappa = \text{cf}(\mathcal{X}, \rightsquigarrow)$ and $\mathfrak{r}_{\kappa_i} = \text{cf}(\mathcal{X}_i, \rightsquigarrow_i)$ for all $i < \theta$.

For each $i < \theta$, define a map $\pi_i : \mathcal{X} \rightarrow \mathcal{X}_i$ as follows. For all $x \in \mathcal{X}$, if $|x \cap \kappa_i| = \kappa_i$, then let $\pi_i(x) = x \cap \kappa_i$. Otherwise, let $\pi_i(x) = \kappa_i$. Note that, for all $x \in [\kappa]^\kappa$, the set of $i < \theta$ for which $|x \cap \kappa_i| = \kappa_i$, and hence for which $\pi_i(x) = x \cap \kappa_i$, is a club in

θ . Finally, as in the case of density, let $e : \text{NS}_\theta^+ \rightarrow \text{Card}$ be the constant function taking value $\underline{d}(\theta, \kappa)$.

5.4. The dominating number.

Definition 5.4. Suppose that λ is an infinite cardinal.

- (1) If $f, g \in {}^\lambda \text{On}$, then $f <^* g$ if and only if $|\{\eta < \lambda \mid g(\eta) \leq f(\eta)\}| < \lambda$.
- (2) The *dominating number* \mathfrak{d}_λ is the minimal cardinality of a family $\mathcal{F} \subseteq {}^\lambda \lambda$ such that for every $g \in {}^\lambda \lambda$, there is $f \in \mathcal{F}$ such that $g <^* f$.
- (3) More generally, for any limit ordinal σ , $\mathfrak{d}_{\lambda, \sigma}$ is the minimal cardinality of a family $\mathcal{F} \subseteq {}^\lambda \sigma$ such that, for every $g \in {}^\lambda \sigma$, there is $f \in \mathcal{F}$ such that $g <^* f$.

Proofs of the following basic facts can be found in [7].

Proposition 5.5 ([7, Claim 3.2 and Lemma 3.1]). *Suppose that λ is an infinite cardinal.*

- (1) $\mathfrak{d}_\lambda > \lambda$ and $\text{cf}(\mathfrak{d}_\lambda) > \lambda$.
- (2) $\mathfrak{d}_\lambda = \mathfrak{d}_{\lambda, \text{cf}(\lambda)}$.

The proofs of [7, Claim 3.2 and Lemma 3.1] can be routinely adapted to yield the following generalization of the preceding proposition.

Fact 5.6. *Suppose that λ is an infinite cardinal and σ is a limit ordinal.*

- (1) $\mathfrak{d}_{\lambda, \sigma} > \lambda$ and $\text{cf}(\mathfrak{d}_{\lambda, \sigma}) > \lambda$.
- (2) $\mathfrak{d}_{\lambda, \sigma} = \mathfrak{d}_{\lambda, \text{cf}(\sigma)}$.

Let $\mathcal{X} = {}^\kappa \theta$ and, for each limit ordinal $i < \theta$, let $\mathcal{X}_i = {}^{\kappa_i} i$. Let \rightsquigarrow and \rightsquigarrow_i be the relations $<^*$ on \mathcal{X} and \mathcal{X}_i , respectively. By Proposition 5.5 and Fact 5.6, we have $\mathfrak{d}_\kappa = \text{cf}(\mathcal{X}, \rightsquigarrow)$ and, for all limit $i < \theta$, we have $\mathfrak{d}_{\kappa_i} = \text{cf}(\mathcal{X}_i, \rightsquigarrow_i)$.

For each limit ordinal $i < \theta$, define a map $\pi_i : \mathcal{X} \rightarrow \mathcal{X}_i$ as follows. For all $x \in \mathcal{X}$ and all $\eta < \kappa_i$, let

$$\pi_i(x)(\eta) = \begin{cases} x(\eta) & \text{if } x(\eta) < i \\ 0 & \text{otherwise.} \end{cases}$$

Note that we do indeed have $\pi_i(x) \in {}^{\kappa_i} i = \mathcal{X}_i$, as desired. Finally, let $e : \text{NS}_\theta^+ \rightarrow \text{Card}$ be defined by letting $e(S) = \underline{d}_{\text{stat}}(S, \kappa)$ for all stationary $S \subseteq \theta$ (recall this notation from Definition 3.3).

5.5. The general lemma. We now show that, in all of the cases introduced in this section, the corresponding version of $\Psi^*(T, 0)$ holds for all stationary $T \subseteq \theta$. Note that we may assume that $\varphi_0^\theta(i) = 0$ for all $i < \theta$, so clause (2) in the definition of $\psi(T, 0)$ asserts that $|\mathcal{Y}_i| \leq \kappa_i$ for all $i \in T$.

Lemma 5.7. *For any cardinal characteristic $\mathfrak{cc} \in \{m(\theta, \kappa), \mathcal{D}(\kappa, \theta), \mathfrak{t}_\kappa, \mathfrak{d}_\kappa\}$, the corresponding formula $\Psi^*(T, 0)$ holds for every stationary $T \subseteq \theta$.*

Proof. We begin with some general preliminaries and then split into cases depending on the cardinal characteristic under consideration.

Fix a stationary set $T \subseteq \theta$; we may assume that every element of T is a limit ordinal. Fix assignments for \mathcal{X} , \rightsquigarrow , $(\langle \mathcal{X}_i, \rightsquigarrow_i, \pi_i \mid i < \theta \rangle)$, and e corresponding to one of the cardinal characteristics introduced in this section. To verify $\Psi^*(T, 0)$, fix a set $\mathcal{Z} \subseteq \mathcal{X}$ and, for each $i \in T$, a set $\mathcal{Y}_i \subseteq \mathcal{X}_i$ such that

- for all $i \in T$, $|\mathcal{Y}_i| \leq \kappa_i$;

- for all $z \in \mathcal{Z}$, there is a club $C \subseteq \theta$ such that, for all $i \in C \cap T$, there is $y \in \mathcal{Y}_i$ for which $\pi_i(z) \rightsquigarrow_i y$.

For each $i \in T$, enumerate \mathcal{Y}_i as $\langle y_{i,\xi} \mid \xi < \kappa_i \rangle$ (with repetitions if $|\mathcal{Y}_i| < \kappa_i$). We will find $\mathcal{Y} \subseteq \mathcal{X}$ such that $|\mathcal{Y}| \leq \kappa + e(T)$ and, for all $z \in \mathcal{Z}$, there is $y \in \mathcal{Y}$ for which $z \rightsquigarrow y$. Our method for doing this will depend on the precise cardinal characteristic.

Case 1: $m(\theta, \kappa)$. Recall that in this case $e(T) = \sum_{j < \theta} m(\theta, \kappa_j)$. Therefore, for each $j < \theta$, we can fix a family $\mathcal{W}_j \subseteq [T \times \kappa_j]^\theta$ such that

- $|\mathcal{W}_j| \leq e(T)$;
- for all $u \in [T \times \kappa_j]^\theta$, there is $w \in \mathcal{W}_j$ such that $|w \cap u| = \theta$.

Let $\mathcal{W} := \bigcup_{j < \theta} \mathcal{W}_j$. For each $w \in \mathcal{W}$, let $y_w^* := \bigcup \{y_{i,\xi} \mid (i, \xi) \in w\}$, and note that $y_w^* \in [\kappa]^{< \theta}$.

Also, for each $j < \theta$, let $\mathcal{Y}_j^* \subseteq [\kappa_j]^\theta$ be such that $|\mathcal{Y}_j^*| \leq e(T)$ and, for all $x \in [\kappa_j]^\theta$, there is $y \in \mathcal{Y}_j^*$ such that $|y \cap x| = \theta$. Finally, let

$$\mathcal{Y} := \bigcup_{j < \theta} \mathcal{Y}_j^* \cup (\{y_w^* \mid w \in \mathcal{W}_j\} \cap [\kappa]^\theta).$$

We claim that \mathcal{Y} is as desired. It is evident that $\mathcal{Y} \subseteq \mathcal{X}$ and $|\mathcal{Y}| \leq \kappa + e(T)$. It remains to show that, for all $z \in \mathcal{Z}$, there is $y \in \mathcal{Y}$ such that $z \rightsquigarrow y$, i.e., $|y \cap z| = \theta$. To this end, fix $z \in \mathcal{Z}$. If there is $j < \theta$ such that $z \subseteq \kappa_j$, then there is $y \in \mathcal{Y}_j^*$ such that $|y \cap z| = \theta$, and we are done. Therefore, we may assume that z is unbounded in κ . By assumption, there is a club $C \subseteq \theta$ such that, for all $i \in C \cap T$, we have

- $\sup(z \cap \kappa_i) = \kappa_i$, and hence $\pi_i(z)$ is an unbounded subset of $z \cap \kappa_i$ of order type $\text{cf}(i)$;
- there is $\xi_i < \kappa_i$ for which $\pi_i(z) \rightsquigarrow_i y_{i,\xi_i}$, i.e., $|y_{i,\xi_i} \cap \pi_i(z)| = \text{cf}(i)$.

Since each $i \in T$ is a limit ordinal, we can apply Fodor's lemma to find a fixed $j < \theta$ and a stationary $T' \subseteq C \cap T$ such that $\xi_i < \kappa_j$ for all $i \in T'$. Let $u = \{(i, \xi_i) \mid i \in T'\}$. Then $u \in [T \times \kappa_j]^\theta$, so, by our choice of \mathcal{W}_j , we can find $w \in \mathcal{W}_j$ such that $|w \cap u| = \theta$. It follows that the set $T'' := \{i \in T' \mid (i, \xi_i) \in w\}$ is unbounded in θ .

Consider the set $y_w^* = \bigcup \{y_{i,\xi} \mid (i, \xi) \in w\}$. Note that, for all $i \in T''$, we know that $\pi_i(z)$ is an unbounded subset of $z \cap \kappa_i$ of order type $\text{cf}(i)$, and we know that $|y_{i,\xi_i} \cap \pi_i(z)| = \text{cf}(i)$. It follows that $y_{i,\xi_i} \cap \pi_i(z)$ is an unbounded subset of $z \cap \kappa_i$, and therefore $y_w^* \cap z$ is an unbounded subset of z . In particular, $y_w^* \in \mathcal{Y}$ and $z \rightsquigarrow y_w^*$, as desired.

Case 2: $d(\theta, \kappa)$. The argument in this case is very similar to that in the previous case, so we suppress some details. Since $e(T) = \underline{d}(\theta, \kappa) = \sum_{j < \theta} d(\theta, \kappa_j)$, we can fix, for each $j < \theta$, a family $\mathcal{W}_j \subseteq [T \times \kappa_j]^\theta$ that is dense in $([T \times \kappa_j]^\theta, \subseteq)$ and has cardinality at most $e(T)$. Let $\mathcal{W} := \bigcup_{j < \theta} \mathcal{W}_j$ and, for each $w \in \mathcal{W}$, let $y_w^* := \bigcup \{y_{i,\xi} \mid (i, \xi) \in w\}$.

Also, for each $j < \theta$, fix $\mathcal{Y}_j^* \subseteq [\kappa_j]^\theta$ that is dense in $([\kappa_j]^\theta, \subseteq)$ and has cardinality at most $e(T)$. Let $\mathcal{Y} := \bigcup_{j < \theta} \mathcal{Y}_j^* \cup (\{y_w^* \mid w \in \mathcal{W}_j\} \cap [\kappa]^\theta)$.

It is evident that $\mathcal{Y} \subseteq \mathcal{X}$ and $|\mathcal{Y}| \leq \kappa + e(T)$. To verify that \mathcal{Y} is as desired, fix $z \in \mathcal{Z}$. We must find $y \in \mathcal{Y}$ such that $y \subseteq z$. If there is $j < \theta$ such that $z \subseteq \kappa_j$, then there is $y \in \mathcal{Y}_j^*$ such that $y \subseteq z$, and we are done. Thus, suppose that z is unbounded in κ . Then there is a club $C \subseteq \theta$ such that, for all $i \in C \cap T$, we have

- $\sup(z \cap \kappa_i) = \kappa_i$;

- there is $\xi_i < \kappa_i$ for which $y_{i,\xi_i} \subseteq \pi_i(z)$.

We can again find a stationary $T' \subseteq C \cap T$ and a fixed $j < \theta$ such that $\xi_i < \kappa_j$ for all $i \in T'$. Let $u = \{(i, \xi_i) \mid i \in T'\}$, and find $w \in \mathcal{W}_j$ such that $w \subseteq u$. As before, the set $T'' := \{i \in T' \mid (i, \xi_i) \in w\}$ is unbounded in θ ; moreover, w is *precisely* $\{(i, \xi_i) \mid i \in T''\}$. Therefore, $y_w^* = \bigcup \{y_{i,\xi_i} \mid i \in T''\} \subseteq z$. It also follows exactly as in the previous case that y_w^* is unbounded in κ and is therefore an element of \mathcal{Y} .

Case 3: \mathfrak{r}_κ . In this case, $e(T) = \underline{d}(\theta, \kappa)$. Therefore, for each $j < \theta$, we can fix as in the previous case a set $\mathcal{W}_j \subseteq [T \times \kappa_j]^\theta$ such that \mathcal{W}_j is dense in $([T \times \kappa_j]^\theta, \subseteq)$ and $|\mathcal{W}_j| \leq d(T)$. Let $\mathcal{W} = \bigcup_{j < \theta} \mathcal{D}_j$ and, for each $w \in \mathcal{D}$, let $y_w^* = \bigcup \{y_{i,\xi} \mid (i, \xi) \in w\}$. Finally, let $\mathcal{Y} = \{y_w^* \mid w \in \mathcal{D}\} \cap [\kappa]^\kappa$.

We claim that \mathcal{Y} is as desired. It is evident that $\mathcal{Y} \subseteq [\kappa]^\kappa$ and $|\mathcal{Y}| \leq e(T)$. It remains to verify that no element of \mathcal{Z} splits every element of \mathcal{Y} . To this end, fix $z \in \mathcal{Z}$. By assumption, we can find a club $C \subseteq \theta$ such that, for all $i \in C \cap T$, we can find $\xi_i < \kappa_i$ such that either $|y_{i,\xi_i} \cap \pi_i(z)| < \kappa_i$ or $|y_{i,\xi_i} \setminus \pi_i(z)| < \kappa_i$. We can also assume that, for all $i \in C$, we have $|z \cap \kappa_i| = \kappa_i$, and hence $\pi_i(z) = z \cap \kappa_i$.

Find a stationary $S_0 \subseteq C \cap T$ such that either

- (1) for all $i \in S_0$, $|y_{i,\xi_i} \cap \pi_i(z)| < \kappa_i$; *or*
- (2) for all $i \in S_0$, $|y_{i,\xi_i} \setminus \pi_i(z)| < \kappa_i$.

Without loss of generality, assume that (1) holds (the proof is symmetric if (2) holds). For each limit ordinal $i \in S_0$, we can find $j(i) < i$ such that $\max\{\xi_i, |y_{i,\xi_i} \cap \pi_i(z)|\} < \kappa_{j(i)}$. Since S_0 is a stationary subset of θ , we can therefore find a stationary $S_1 \subseteq S_0$ and a fixed $j < \theta$ such that $j(i) = j$ for all $i \in S_1$.

Let $u = \{(i, \xi_i) \mid i \in S_1\}$. Then $u \in [\theta \times \kappa_j]^\theta$, so we can find $w \in \mathcal{W}_j \subseteq \mathcal{W}$ such that $w \subseteq u$. Note that $\{i \in S_1 \mid (i, \xi_i) \in w\}$ must be unbounded in θ , so we have $|y_w^*| = \kappa$, and hence $y_w^* \in \mathcal{Y}$. Moreover,

$$y_w^* \cap z = \bigcup_{(i,\xi) \in w} (y_{i,\xi} \cap z) = \bigcup_{(i,\xi) \in w} (y_{i,\xi} \cap \pi_i(z)) \subseteq \bigcup_{i \in S_1} (y_{i,\xi_i} \cap \pi_i(z)).$$

Since $|y_{i,\xi_i} \cap \pi_i(z)| < \kappa_j$ for all $i \in S_1$ and $\kappa_j > \theta = |S_1|$, it follows that $|y_w^* \cap z| < \kappa_j < \kappa$, so z does not split y_w^* , i.e., we have $z \rightsquigarrow y_w^*$, as desired.

Case 4: \mathfrak{d}_κ . In this case, $e(T) = \underline{d}_{\text{stat}}(T, \kappa)$. Therefore, we can fix a family \mathcal{H} such that

- $|\mathcal{H}| = e(T)$;
- every element of \mathcal{H} is a function from a stationary subset of T to κ whose range is bounded below κ ;
- for every function g from a stationary subset of T to κ such that the range of g is bounded below κ , there is a function $h \in \mathcal{H}$ such that $\{i \in \text{dom}(h) \mid i \notin \text{dom}(g) \text{ or } h(i) \neq g(i)\}$ is nonstationary in θ .

Recall also that, for each $i \in T$ and each $\xi < \kappa_i$, we have $y_{i,\xi} \in {}^{\kappa_i}i$.

For each $h \in \mathcal{H}$, define a function $y_h^* \in {}^\kappa\theta$ as follows. Let $T' := \text{dom}(h)$. Since the range of h is bounded below κ , we know that, for all sufficiently large $i \in T'$, we have $h(i) < \kappa_i$, and hence $y_{i,h(i)}$ is defined. Therefore, for all sufficiently large $i \in T'$ and all $\eta < \kappa_i$, we have $y_{i,h(i)}(\eta) < i$. Therefore, by Fodor's Lemma, for each $\eta < \kappa$, we can find a $j < \theta$ such that

$$\{i \in T' \mid \eta < \kappa_i, h(i) < \kappa_i \text{ and } y_{i,h(i)}(\eta) = j\}$$

is stationary in θ . Let $y_h^*(\eta)$ be the least such j .

Let $\mathcal{Y} = \{y_h^* \mid h \in \mathcal{H}\}$. We claim that \mathcal{Y} is as desired. It is evident that $\mathcal{Y} \subseteq {}^\kappa\theta$ and $|\mathcal{Y}| \leq e(T)$. It remains to verify that, for every $z \in \mathcal{Z}$, there is $y \in \mathcal{Y}$ such that $z <^* y$. To this end, fix $z \in \mathcal{Z}$. By assumption, we can find a club $C \subseteq \theta$ such that, for all $i \in C \cap T$, there is $\xi_i < \kappa_i$ for which $\pi_i(z) <^* y_{i,\xi_i}$. For each $i \in C \cap T$, let

$$E_i := \{\eta < \kappa_i \mid y_{i,\xi_i}(\eta) \leq \pi_i(z)(\eta)\},$$

and note that $|E_i| < \kappa_i$. By two applications of Fodor's lemma, we can find a $j < \theta$ and a stationary $T' \subseteq C \cap T$ such that $\max\{\xi_i, |E_i|\} < \kappa_j$ for all $i \in T'$. Then the map from T' to κ defined by sending each $i \in T'$ to ξ_i has range bounded below κ , so we can find $h \in \mathcal{H}$ and a club $D \subseteq \theta$ such that, letting $T'' := \text{dom}(h)$, the following statement holds: for all $i \in D \cap T''$, we have $i \in T'$ and $h(i) = \xi_i$.

We claim that $z <^* y_h^*$. To see this, first let $E := \bigcup_{i \in D \cap T''} E_i$, and note that $|E| \leq \theta \cdot \kappa_j < \kappa$. It therefore suffices to show that $z(\eta) < y_h^*(\eta)$ for all $\eta \in \kappa \setminus E$. To this end, fix $\eta \in \kappa \setminus E$. Fix $\ell < \theta$ such that $\eta < \kappa_\ell$ and $z(\eta) < \ell$. Then, for all $i \in D \cap T'' \setminus \ell$, we have $\pi_i(z)(\eta) = z(\eta)$. Moreover, for all such i , we have $\eta \notin E_i$, and hence $z(\eta) < y_{i,\xi_i}(\eta) = y_{i,h(i)}(\eta)$. Recall that $y_h^*(\eta)$ was defined in such a way that there is a stationary set $T^* \subseteq T''$ such that $y_h^*(\eta) = y_{i,h(i)}(\eta)$ for all $i \in T^*$. Since D is a club in θ , we can fix $i^* \in (D \cap T^*) \setminus \ell$. But then we have $z(\eta) < y_{i^*,h(i^*)}(\eta) = y_h^*(\eta)$, as desired. \square

Combining the results of this and the previous section, we obtain the precise statement of our main result.

Main Theorem. *Suppose that*

- κ is a singular cardinal and $\text{cf}(\kappa) = \theta > \omega$;
- $\langle \kappa_i \mid i < \theta \rangle$ is an increasing, continuous sequence of cardinals converging to κ ;
- β is an ordinal for which φ_β^θ exists;
- $S \subseteq \theta$ is stationary;
- \mathbf{cc} is one of the cardinal characteristics $m(\theta, \kappa)$, $d(\theta, \kappa)$, \mathfrak{r}_κ , or \mathfrak{d}_κ , and, for each $i < \theta$, \mathbf{cc}_i is the corresponding cardinal characteristic $m(\text{cf}(\kappa_i), \kappa_i)$, $d(\text{cf}(\kappa_i), \kappa_i)$, \mathfrak{r}_{κ_i} , or \mathfrak{d}_{κ_i} ;
- for all $i \in S$, we have $\mathbf{cc}_i \leq \kappa_i^{+\varphi_\beta^\theta(i)}$.

Then:

- (1) If $\mathbf{cc} = m(\theta, \kappa)$, then $\mathbf{cc} \leq \kappa^{+\beta} + \sum_{j < \theta} m(\theta, \kappa_j) + d(\text{NS}_\theta \upharpoonright S)$. Moreover, if $\beta < \omega$, then $\mathbf{cc} \leq \kappa^{+\beta} + \sum_{j < \theta} m(\theta, \kappa_j)$.
- (2) If $\mathbf{cc} = d(\theta, \kappa)$ or $\mathbf{cc} = \mathfrak{r}_\kappa$, then $\mathbf{cc} \leq \kappa^{+\beta} + \underline{d(\kappa, \theta)} + d(\text{NS}_\theta \upharpoonright S)$. Moreover, if $\beta < \omega$, then $\mathbf{cc} \leq \kappa^{+\beta} + \underline{d(\theta, \kappa)}$.
- (3) If $\mathbf{cc} = \mathfrak{d}_\kappa$, then $\mathbf{cc} \leq \kappa^{+\beta} + \underline{d_{\text{stat}}(S, \kappa)}$.

Proof. This follows directly from Lemma 5.7 and Corollary 4.4. \square

6. OPEN QUESTIONS

Throughout this section, κ will denote an arbitrary infinite cardinal. We feel that the most prominent cardinal characteristic at singular cardinals that is not covered by our results here is the *ultrafilter number*, a close relative of the reaping number.

Definition 6.1. The ultrafilter number \mathfrak{u}_κ is the minimal size of a base for a uniform ultrafilter over κ . In other words, it is the minimal cardinal λ for which there exists a uniform ultrafilter U over κ and a family $\mathcal{X} \subseteq U$ such that $|\mathcal{X}| = \lambda$ and, for all $Y \in U$, there is $X \in \mathcal{X}$ such that $|X \setminus Y| < \kappa$.

It is provable that $\mathfrak{u}_\kappa > \kappa$, and the ultrafilter number at singular cardinals has been extensively studied (cf. [5], [7], [9], among others).

Question 6.2. *Does a version of our Main Theorem hold for the ultrafilter number?*

We briefly mentioned the almost disjointness number at the beginning of Section 5; we feel that some interesting questions can be formulated around it. We first recall the relevant definitions.

Definition 6.3. An *almost disjoint family* over κ is a family $\mathcal{A} \subseteq [\kappa]^\kappa$ such that, for all distinct $A, B \in \mathcal{A}$, we have $|A \cap B| < \kappa$. Such a family is a *maximal almost disjoint family* (MAD family) over κ if, moreover, there is no almost disjoint family \mathcal{B} over κ with $\mathcal{A} \subsetneq \mathcal{B}$.

There are trivial ways to form MAD families over κ (as an extreme case, $\{\kappa\}$ is a MAD family over κ). The *almost disjointness number* \mathfrak{a}_κ is defined to be the minimal cardinality of a *nontrivial* MAD family over κ . It remains to specify what *nontriviality* means. The most natural solution seems to be to say that a MAD family is nontrivial if and only if its cardinality is greater than $\text{cf}(\kappa)$ (this is the approach taken, for instance, in [13]). Under this definition, it is not difficult to prove that $\mathfrak{a}_\kappa \leq \mathfrak{a}_{\text{cf}(\kappa)}$. However, there always exist MAD families over κ of cardinality strictly greater than κ , so one could also declare that a MAD family over κ is nontrivial if and only if its cardinality is greater than κ . Let us denote the version of the almost disjointness number arising from this more stringent definition of nontriviality by \mathfrak{a}_κ^* .

Question 6.4. *Does a version of our Main Theorem hold for \mathfrak{a}_κ^* ?*

The most immediate specific incarnation of this question would be the following:

Question 6.5. *Suppose that κ is a singular strong limit cardinal of uncountable cofinality and there is a stationary subset $S \subseteq \kappa$ consisting of singular cardinals such that, for all $\mu \in S$, there is a MAD family over μ of cardinality μ^+ . Must there be a MAD family over κ of cardinality κ^+ .*

Definition 6.6. A graph G is *universal for graphs of size κ* if, for every graph H with at most κ -many vertices, there is an induced subgraph of G that is isomorphic to H . Let \mathfrak{ug}_κ denote the minimal number of vertices in a graph G that is universal for graphs of size κ .

Question 6.7. *Does a version of our Main Theorem hold for \mathfrak{ug}_κ ?*

We are also interested in whether analogues of Silver's theorem hold for statements that are not naturally formulated as statements about cardinal characteristics but which are consequences of $2^\kappa = \kappa^+$. We record here some particularly prominent examples.

Definition 6.8. The polarized partition relation $\binom{\kappa^+}{\kappa} \rightarrow \binom{\kappa^+}{\kappa}_2^{1,1}$ is the assertion that, for every function $c : \kappa^+ \times \kappa \rightarrow 2$, there are sets $A \in [\kappa^+]^{\kappa^+}$ and $B \in [\kappa]^\kappa$

such that $c \upharpoonright A \times B$ is constant. The negation of this relation is denoted by $\left(\begin{smallmatrix} \kappa^+ \\ \kappa \end{smallmatrix}\right) \not\rightarrow \left(\begin{smallmatrix} \kappa^+ \\ \kappa \end{smallmatrix}\right)_2^{1,1}$.

Erdős, Hajnal, and Rado prove in [3] that, if $2^\kappa = \kappa^+$, then $\left(\begin{smallmatrix} \kappa^+ \\ \kappa \end{smallmatrix}\right) \not\rightarrow \left(\begin{smallmatrix} \kappa^+ \\ \kappa \end{smallmatrix}\right)_2^{1,1}$. On the other hand, Garti and Shelah prove in [6] that, assuming the consistency of a supercompact cardinal, the positive relation $\left(\begin{smallmatrix} \kappa^+ \\ \kappa \end{smallmatrix}\right) \rightarrow \left(\begin{smallmatrix} \kappa^+ \\ \kappa \end{smallmatrix}\right)_2^{1,1}$ consistently holds for a singular strong limit cardinal κ (in their result, κ can have either countable or uncountable cofinality).

Question 6.9. *Suppose that κ is a singular cardinal of uncountable cofinality and there is a stationary set $S \subseteq \kappa$ consisting of singular cardinals such that, for all $\mu \in S$, we have $\left(\begin{smallmatrix} \mu^+ \\ \mu \end{smallmatrix}\right) \not\rightarrow \left(\begin{smallmatrix} \mu^+ \\ \mu \end{smallmatrix}\right)_2^{1,1}$. Must it be the case that $\left(\begin{smallmatrix} \kappa^+ \\ \kappa \end{smallmatrix}\right) \not\rightarrow \left(\begin{smallmatrix} \kappa^+ \\ \kappa \end{smallmatrix}\right)_2^{1,1}$?*

Recall that, for a regular uncountable cardinal λ , a λ -Aronszajn tree is a tree of height λ with no levels or branches of cardinality λ . By a result of Specker [19], if μ is regular and $\mu^{<\mu} = \mu$, then there is a μ^+ -Aronszajn tree. In particular, if $2^\kappa = \kappa^+$, then there is a κ^{++} -Aronszajn tree. Therefore, the nonexistence of Aronszajn trees at the double successor of a singular strong limit cardinal requires a failure of the Singular Cardinals Hypothesis. For a construction of a model in which there is a singular strong limit cardinal κ of uncountable cofinality such that there are no κ^{++} -Aronszajn trees, see [8]. For our final question, we ask whether the existence of κ^{++} -Aronszajn trees satisfies a version of Silver’s theorem.

Question 6.10. *Suppose that κ is a singular cardinal of uncountable cofinality and $\{\mu < \kappa \mid \text{there exists a } \mu^{++}\text{-Aronszajn tree}\}$ is stationary in κ . Must there exist a κ^{++} -Aronszajn tree?*

REFERENCES

- [1] J. E. Baumgartner and K. Prikry. On a theorem of Silver. *Discrete Math.*, 14(1):17–21, 1976.
- [2] Andreas Blass. Combinatorial cardinal characteristics of the continuum. In *Handbook of set theory. Vols. 1, 2, 3*, pages 395–489. Springer, Dordrecht, 2010.
- [3] P. Erdős, A. Hajnal, and R. Rado. Partition relations for cardinal numbers. *Acta Math. Acad. Sci. Hungar.*, 16:93–196, 1965.
- [4] Fred Galvin and András Hajnal. Inequalities for cardinal powers. *Ann. of Math. (2)*, 101:491–498, 1975.
- [5] S. Garti and S. Shelah. The ultrafilter number for singular cardinals. *Acta Math. Hungar.*, 137(4):296–301, 2012.
- [6] Shimon Garti and Saharon Shelah. A strong polarized relation. *J. Symbolic Logic*, 77(3):766–776, 2012.
- [7] Shimon Garti and Saharon Shelah. Remarks on generalized ultrafilter, dominating and reaping numbers. *Fund. Math.*, 250(2):101–115, 2020.
- [8] Mohammad Golshani and Rahman Mohammadpour. The tree property at double successors of singular cardinals of uncountable cofinality. *Ann. Pure Appl. Logic*, 169(2):164–175, 2018.
- [9] Radek Honzik and Šárka Stejskalová. Small $u(\kappa)$ at singular κ with compactness at κ^{++} . *Arch. Math. Logic*, pages 1–22, 2021.
- [10] Thomas Jech. A variation on a theorem of Galvin and Hajnal. *Bull. London Math. Soc.*, 25(2):97–103, 1993.
- [11] Thomas Jech. *Set theory*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003. The third millennium edition, revised and expanded.

- [12] Menachem Kojman. On the arithmetic of density. *Topology Appl.*, 213:145–153, 2016.
- [13] Menachem Kojman, Wiesław Kubiś, and Saharon Shelah. On two problems of Erdős and Hechler: new methods in singular madness. *Proc. Amer. Math. Soc.*, 132(11):3357–3365, 2004.
- [14] John Krueger and Ernest Schimmerling. An equiconsistency result on partial squares. *J. Math. Log.*, 11(1):29–59, 2011.
- [15] Pierre Matet. Meeting numbers and pseudopowers. *MLQ Math. Log. Q.*, 67(1):59–76, 2021.
- [16] Pierre Matet. Towers and clubs. *Arch. Math. Logic*, 60(6):683–719, 2021.
- [17] Saharon Shelah. *Cardinal arithmetic*, volume 29 of *Oxford Logic Guides*. The Clarendon Press, Oxford University Press, New York, 1994. Oxford Science Publications.
- [18] Jack Silver. On the singular cardinals problem. In *Proceedings of the International Congress of Mathematicians (Vancouver, B. C., 1974)*, Vol. 1, pages 265–268, 1975.
- [19] E. Specker. Sur un problème de Sikorski. *Colloq. Math.*, 2:9–12, 1949.

INSTITUTE OF MATHEMATICS OF THE CZECH ACADEMY OF SCIENCES, ŽITNÁ 25, PRAHA 1, CZECHIA

Email address: `lambiehanson@math.cas.cz`

URL: `http://math.cas.cz/lambiehanson`