

Effects of Voluntary Running Exercise on Skeletal Muscle Properties in Nonobese Rats with Type 2 Diabetes

Isuzu NAKAMOTO¹, Akihiko ISHIHARA¹

¹Laboratory of Cell Biology and Life Science, Graduate School of Human and Environmental Studies, Kyoto University, Kyoto, Japan

Received April 4, 2019

Accepted September 25, 2019

Epub Ahead of Print December 19, 2019

Summary

The skeletal muscles of animals and humans with type 2 diabetes have decreased oxidative capacity. Aerobic exercise can improve muscle oxidative capacity, but no data are available on the amount of exercise required. We investigated the effects of voluntary running exercise and running distance on the skeletal muscle properties of nonobese rats with type 2 diabetes. Six-week-old male diabetic Goto-Kakizaki rats were divided into nonexercised (GK) and exercised (GK-Ex) groups. The rats in the GK-Ex group were permitted voluntary running exercise on wheels for 6 weeks. Age-matched male Wistar rats (WR) were used as nondiabetic controls. Fasting blood glucose and HbA1c levels were higher in the GK and GK-Ex groups than in the WR group and lower in the GK-Ex group than in the GK group. Succinate dehydrogenase (SDH) activity and peroxisome proliferator-activated receptor γ coactivator-1α (*Pgc-1α*) mRNA levels in the soleus and plantaris muscles were higher in the WR and GK-Ex groups than in the GK group. HbA1c and total cholesterol levels were negatively correlated with running distance and SDH activity and *Pgc-1α* mRNA levels in the soleus muscle were positively correlated with running distance. The onset and progression of diabetes in nonobese diabetic rats were effectively inhibited by running longer distances.

Key words

Goto-Kakizaki rat • Oxidative capacity • *Pgc-1α* mRNA • Skeletal muscle • Type 2 diabetes • Voluntary running exercise

Corresponding author

A. Ishihara, Laboratory of Cell Biology and Life Science, Graduate School of Human and Environmental Studies, Kyoto University, Kyoto 606-8501, Japan. Fax: +81-75-753-6771. E-mail: ishihara.akihiko.8s@kyoto-u.ac.jp

Introduction

The metabolic properties of skeletal muscle, an organ responsible for a major portion of glucose disposal, play an important role in the onset and progression of type 2 diabetes (Phielix and Mensink 2008). The skeletal muscles of experimental animals with type 2 diabetes have decreased oxidative capacity; that is, reduced succinate dehydrogenase (SDH) activity and peroxisome proliferator-activated receptor γ coactivator-1α (*Pgc-1α*) mRNA levels (Nagatomo *et al.* 2009, Nagatomo *et al.* 2011a).

Aerobic exercise has been shown to effectively prevent and/or improve insulin resistance and impaired glucose metabolism (Ishihara *et al.* 2012, Jenkins and Jenks 2017). Aerobic exercise restored the decreased oxidative capacity of skeletal muscles and improved hyperglycemia in obese rats with type 2 diabetes (Yasuda *et al.* 2006).

Pgc-1α is a major regulator of oxidative metabolism in skeletal muscles (Puigserver 2005, Wende *et al.* 2005). High *Pgc-1α* mRNA levels induced increases in mitochondrial content and the percentage of high-oxidative fibers in skeletal muscles (Lin *et al.* 2002, Schuler *et al.* 2006, Wu *et al.* 2002). Diabetic rats showed growth-related decreases in the percentage of high-oxidative fibers and SDH activity in skeletal muscles (Yasuda *et al.* 2007). Therefore, an exercise-induced increase in *Pgc-1α* mRNA levels in skeletal muscles is expected to lead to a fiber type shift and improve muscle SDH activity. Previous studies (Fu *et al.* 2019, Kim *et al.* 2011, Macia *et al.* 2018, Morifuji *et al.* 2012, Qi *et al.* 2011) examined the effects of running exercise on

diabetes; however, these studies used forced running training on a treadmill, and the amount of exercise was constant for all exercised rats. Furthermore, treadmill exercise is stressful because the rats are forcibly run at a constant speed. In this study, we investigated the effects of voluntary running exercise and running distance on the properties of skeletal muscle in diabetic rats, because it is uncertain whether voluntary running distance is related to the improvement of hyperglycemia and decreased muscle oxidative capacity in diabetic rats. We investigated the effects of voluntary running exercise and running distance on the properties of the slow soleus and fast plantaris muscles in diabetic Goto-Kakizaki (GK) rats.

Methods

Ethical approval

All experimental and animal care procedures were conducted in accordance with the Guide for the Care and Use of Laboratory Animals issued by the Institutional Animal Experiment Committee of Kyoto University (Kyoto, Japan) as well as national guidelines and relevant national laws on the protection of animals.

Experimental animals

The nonobese experimental animals with type 2 diabetes used in this study were GK rats, which have low oxidative capacity in skeletal muscles compared to nondiabetic rats (Nagatomo *et al.* 2009, Nagatomo *et al.* 2011a). These rats were established by repeated inbreeding of normal Wistar rats with glucose tolerance at the upper limit of the normal range (Goto *et al.* 1975, 1976). Six-week-old male GK rats were assigned to either the non-exercised (GK, n=5) or exercised (GK-Ex, n=10) group based on their body weights and fasting blood glucose levels, such that the mean values of these parameters in the two groups were similar. Age-matched male Wistar rats were used as nondiabetic controls (WR, n=5). All rats had free access to a standard diet (MF; Oriental East, Tokyo, Japan) and water. The rats were housed under controlled conditions at 22±2 °C with a 12-h light/dark cycle (light from 08:00 to 20:00) and 45–55 % relative humidity. The body weight and food intake of each rat were measured once every 2 weeks.

Voluntary running exercise

Each rat in the GK-Ex group was allowed to run voluntarily on a running wheel (diameter, 31.8 cm; width, 10 cm) that was attached to a breeding cage for small

animals at its own pace 24 h a day for 6 weeks (Ishihara *et al.* 1998). The number of revolutions of each running wheel was recorded daily.

Fasting blood glucose and HbA1c analyses

Fasting blood glucose was measured once every 2 weeks after fasting for 15 h using blood samples collected from the tail vein of conscious rats with a blood glucose meter (GT-1650; Arkray, Kyoto, Japan). At 12 weeks of age, HbA1c levels were analyzed using the same blood glucose meter and a DCA Vantage analyzer (Siemens Healthcare Diagnostics, Erlangen, Germany).

Serum biochemical analyses

After collecting blood samples from the tail vein at 12 weeks of age, the rats were deeply anesthetized, and blood samples were obtained from the abdominal aorta. Total cholesterol, high density lipoprotein (HDL) and low density lipoprotein (LDL) cholesterol, and triglyceride levels were measured using routine laboratory methods. Insulin, high molecular weight adiponectin, and leptin levels were measured using an enzyme-linked immunosorbent assay kit (Shibayagi Corporation, Gunma, Japan).

Muscle histochemical analyses

After collecting blood samples, the soleus and plantaris muscles were bilaterally dissected, cleaned of excess fat and connective tissue, wet-weighed, and subjected to histochemical, biochemical, and mRNA analyses. The right soleus and plantaris muscles were pinned on a corkboard near their approximate *in vivo* length and rapidly frozen in isopentane, which was previously cooled using a mixture of dry ice and acetone, and stored at -80 °C until analysis. The muscle was mounted onto a specimen chuck with Tissue-Tek OCT compound (Sakura Finetek Japan, Tokyo, Japan). Serial transverse sections (16 µm thick) were cut with a cryostat at -25 °C, brought to room temperature, air dried for 30 min, and preincubated in acidic (pH 4.5) and/or alkaline (pH 10.4) conditions for ATPase staining. Soleus muscle fibers were classified as type I (positive response at pH 4.5 and negative response at pH 10.4), type IIA (negative response at pH 4.5 and positive response at pH 10.4), or type IIC (positive response at pH 4.5 and 10.4) (Nagatomo *et al.* 2011b, Nagatomo *et al.* 2012a, Nagatomo *et al.* 2012b). Plantaris muscle fibers were classified as type I (positive response at pH 4.5), type IIA (negative response at pH 4.5), or type IIB (intermediate

response at pH 4.5) (Nagatomo *et al.* 2011a, Nagatomo *et al.* 2011b).

The fiber type percentage and cross-sectional area (CSA) were determined for approximately 100 adjacent fibers in the central region of the muscle sections. Fiber CSA was measured by tracing the outline of each fiber in the muscle section using a computer-assisted image processing system (Neuroimaging System, Kyoto, Japan) (Nagatomo *et al.* 2011b).

Muscle biochemical analyses

The left soleus and plantaris muscles were divided into the proximal and distal regions for biochemical and histochemical analyses, respectively. The proximal regions of the left soleus and plantaris muscles were frozen rapidly in liquid nitrogen for the evaluation of SDH activity (Nagatomo *et al.* 2011b). The muscle was homogenized in five volumes of ice-cold 0.3 M phosphate buffer (pH 7.4) using a glass tissue homogenizer. The final concentrations of the reaction mixture components were as follows: sodium succinate, 17 mM; sodium cyanide, 1 mM; aluminum chloride, 0.4 mM; and calcium chloride, 0.4 mM. Reduction of cytochrome *c* in this reaction mixture was analyzed by measuring the increase in extinction at 550 nm with a spectrophotometer. SDH activity was calculated from the ferricytochrome *c* concentration and protein content.

Muscle mRNA analysis

The distal regions of the left soleus and plantaris muscles were rapidly frozen in liquid nitrogen, and total RNA was extracted from the tissues using the QuickGene RNA Tissue Kit SII (Fujifilm, Tokyo, Japan). Reverse transcription was performed using the High-Capacity cDNA Archive Kit (Applied Biosystems, CA, USA), and the resultant cDNA samples were stored. *Pgc-1α* expression levels were quantified by TaqMan Gene Expression Assays (Applied Biosystems, CA, USA) as described previously (Nagatomo *et al.* 2011b). Each TaqMan probe and primer set was validated by quantitative real-time polymerase chain reaction with a series of cDNA template dilutions to obtain standard curves of the threshold cycle against the relative concentration of *S18* (housekeeping gene) as an internal standard. All sample and non-template control reactions were performed in a 7500 Fast Sequence Detection System (Applied Biosystems, CA, USA). Target mRNA levels were normalized to the levels in the WR group.

Statistical analyses

Values are expressed as mean \pm standard deviation (SD). Analysis of variance followed by Scheffé's *post hoc* test was used to evaluate differences among the WR, GK, and GK-Ex groups. Correlations were assessed using Pearson's correlation coefficient. Statistical significance was set at $p < 0.05$.

Results

Voluntary running distance

The voluntary running distances in the GK-Ex group increased every week (Fig. 1). The mean running distance during the 6 weeks of voluntary running exercise was 4.7 ± 1.2 km/day.




Fig. 1. Voluntary running distance in exercised Goto-Kakizaki rats. Each rat in the GK-Ex group was allowed to voluntarily run in a running wheel that was attached to a breeding cage for small animals at its own pace for 24 h a day during the 6 weeks of voluntary running exercise. Values are expressed as mean \pm SD ($n=10$).

Body weight

The body weights of the WR, GK, and GK-Ex groups increased every week (Fig. 2A). There were no differences in body weight among the age-matched three groups, irrespective of age.

Food intake

At 6 weeks of age, food intake was lower in the GK-Ex group than in the WR and GK groups ($p < 0.05$, Fig. 2B). At 8 weeks of age, food intake was higher in the GK-Ex group than in the WR group ($p < 0.05$). At 10 and 12 weeks of age, food intake was higher in the GK-Ex group than in the WR and GK groups ($p < 0.05$).

Fasting blood glucose levels

At 8, 10, and 12 weeks of age, fasting blood

glucose levels were higher in the GK group than in the WR group ($p<0.05$) and lower in the GK-Ex group than in the GK group ($p<0.05$, Fig. 2C).




Fig. 2. Body weight (A), food intake (B), and fasting blood glucose (C) in the Wistar (WR), Goto-Kakizaki (GK), and exercised Goto-Kakizaki (GK-Ex) groups. Blood glucose levels were measured after fasting for 15 h using blood samples collected from the tail vein of conscious rats. Values are expressed as mean \pm SD (n=5 for WR and GK; n=10 for GK-Ex). ^a $p<0.05$ compared with WR; ^b $p<0.05$ compared with GK.

Serum biochemical parameters

HbA1c (Fig. 3A) and total cholesterol (Fig. 3B) levels were higher in the GK group than in the WR and GK-Ex groups ($p<0.05$) and higher in the GK-Ex group than in the WR group ($p<0.05$). HDL (Fig. 3C) and LDL (Fig. 3D) cholesterol levels were higher in the GK group than in the WR group ($p<0.05$) and higher in the GK group than in the GK-Ex group ($p<0.05$).

Triglyceride (Fig. 3E) and insulin (Fig. 3F) levels were lower in the GK and GK-Ex groups than in the WR group ($p<0.05$). There were no differences in adiponectin (Fig. 3G) and leptin (Fig. 3H) levels among the three groups.

Muscle weight

The soleus muscle weight was higher in the GK-Ex group than in the WR and GK groups (Fig. 4A). The soleus muscle weight per body weight was higher in the GK-Ex group than in the GK group (Fig. 4B). The plantaris muscle weight (Fig. 4C) and muscle weight per body weight (Fig. 4D) were higher in the GK-Ex group than in the WR and GK groups.

Muscle histochemical properties

The soleus muscles in the WR group contained three types of fibers: I, IIA, and IIC (Fig. 5A, B), whereas these muscles in the GK group contained only type I fibers (Fig. 5C, D). In the GK-Ex group, seven rats had only type I fibers (Fig. 5E, F), whereas the other three had fiber types: I, IIA, and IIC (Fig. 5G, H). The percentage of type I fibers was higher in the GK and GK-Ex groups than in the WR group ($p<0.05$, Fig. 6A), whereas the percentage of type IIA fibers was lower in the GK-Ex group than in the WR group ($p<0.05$). The CSA of type I fibers was larger in the GK-Ex group than in the WR and GK groups ($p<0.05$, Fig. 6B), whereas the CSA of type IIA fibers was larger in the GK-Ex group than in the WR group ($p<0.05$).

The plantaris muscles in all three groups contained three types of fibers: I, IIA, and IIB (Fig. 7). The percentages of type IIA and type IIB fibers were lower in the GK group than in the WR group ($p<0.05$, Fig. 6C), whereas these percentages were higher in the GK-Ex group than in the GK group ($p<0.05$). There were no differences in the CSAs among the three groups, irrespective of fiber type (Fig. 6D).

Muscle SDH activity and *Pgc-1α* mRNA levels

SDH activity in the soleus (Fig. 8A) and plantaris (Fig. 8B) muscles was lower in the GK group than in the WR group ($p<0.05$) and higher in the GK-Ex group than in the GK group ($p<0.05$).

Pgc-1α mRNA levels in the soleus (Fig. 8C) and plantaris (Fig. 8D) muscles were lower in the GK group than in the WR group ($p<0.05$) and higher in the GK-Ex group than in the GK group ($p<0.05$).




Fig. 3. HbA1c (**A**), total cholesterol (**B**), high density lipoprotein (HDL) cholesterol (**C**), low density lipoprotein (LDL) cholesterol (**D**), triglyceride (**E**), insulin (**F**), adiponectin (**G**), and leptin (**H**) levels in the Wistar (WR), Goto-Kakizaki (GK), and exercised Goto-Kakizaki (GK-Ex) groups. Values are expressed as mean \pm SD (n=5 for WR and GK; n=10 for GK-Ex). ^ap<0.05 compared with WR; ^bp<0.05 compared with GK.




Fig. 4. Muscle weight (**A, B**) and relative weight per body weight (**C, D**) of the soleus and plantaris muscles in the Wistar (WR), Goto-Kakizaki (GK), and exercised Goto-Kakizaki (GK-Ex) groups. Values are expressed as mean \pm SD (n=5 for WR and GK; n=10 for GK-Ex). ^ap<0.05 compared with WR; ^bp<0.05 compared with GK.




Fig. 5. Serial transverse sections of the soleus muscles of rats in the control (WR; **A-B**), Goto-Kakizaki (GK; **C-D**), and exercised Goto-Kakizaki (GK-Ex; **E-H**) groups stained for ATPase activity after preincubation at pH 10.4 (**A, C, E**, and **G**) and pH 4.5 (**B, D, F**, and **H**). In the GK-Ex group, the soleus muscles in seven rats contained only type I fibers (**E, F**), whereas these muscles in three rats contained three fiber types (**G, H**). 1, type I; 2, type IIA; 3, type IIC. Scale bar=200 μ m.




Fig. 6. Fiber type percentages (**A, C**) and cross-sectional areas (CSAs; **B, D**) in the soleus and plantaris muscles of the Wistar (WR), Goto-Kakizaki (GK), and exercised Goto-Kakizaki (GK-Ex) groups. Values are expressed as mean \pm SD (n=5 for WR and GK; n=10 for GK-Ex). ^a p<0.05 compared with WR; ^b p<0.05 compared with GK.

Relationship between voluntary running distance and the abovementioned parameters

In the GK-Ex group, there were negative relationships between voluntary running distance and HbA1c ($r=-0.81, p<0.05$, Fig. 9A) and total cholesterol ($r=-0.83, p<0.05$, Fig. 9B) levels. There were positive relationships between voluntary running distance and both SDH activity ($r=0.98, p<0.05$, Fig. 9C) and *Pgc-1α* mRNA levels ($r=0.70, p<0.05$, Fig. 9D) in the soleus muscle.




Fig. 7. Serial transverse sections of the plantaris muscles in rats of the Wistar (WR; **A**), Goto-Kakizaki (GK; **B**), and exercised Goto-Kakizaki (GK-Ex; **C**) groups stained for ATPase activity after preincubation at pH 4.5. 1, type I; 2, type IIA; 3, type IIB. Scale bar=200 μ m.

Discussion

Voluntary running exercise and diabetes

Some previous studies (Macia *et al.* 2018, Morifuji *et al.* 2012, Qi *et al.* 2011) observed that running training did not reduce the increased blood glucose levels in diabetic GK rats. In contrast, other studies (Fu *et al.* 2019, Kim *et al.* 2011) showed that the high blood glucose levels in diabetic GK rats were decreased by running training. However, all these aforementioned studies (Fu *et al.* 2019, Kim *et al.* 2011, Macia *et al.* 2018, Morifuji *et al.* 2012, Qi *et al.* 2011) used forced running training on a treadmill at a constant speed and time, and therefore, did not consider the effects of different running speeds and times. In this study, the growth-related increase in the fasting blood glucose levels of diabetic GK rats was inhibited by voluntary running exercise (Fig. 2C). Furthermore, a negative relationship between the amount of voluntary running distance and HbA1c was observed in diabetic GK rats (Fig. 9A), indicating that the onset and progression of type 2 diabetes were effectively inhibited when diabetic GK rats ran longer distances.

This study showed that nonobese diabetic GK rats have higher total and HDL cholesterol levels (Fig. 3B, C) and lower triglyceride levels than nondiabetic WR rats (Fig. 3E). Previous studies showed that metabolic syndrome (Takemura *et al.* 2017) and diabetic Otsuka Long-Evans Tokushima Fatty (OLETF) (Nagatomo *et al.* 2018) rats had higher total cholesterol, HDL cholesterol, and triglyceride levels. OLETF rats, which are obese and have hypertension, hyperlipidemia, hyperglycemia, hyperinsulinemia, and insulin resistance, were developed as obese diabetic models (Kawano *et al.* 1992). The increased levels of HDL cholesterol in metabolic syndrome and diabetic rats may be due to increased total cholesterol levels. The differences in triglyceride levels between diabetic OLETF and GK rats may be due to the respective presence and absence of obesity in the rats.

Voluntary running exercise and muscle fiber type

In diabetic GK rats, the slow soleus muscles contained only low-oxidative type I fibers, whereas in nondiabetic WR rats, these muscles contained three fiber types: low-oxidative type I and high-oxidative types IIA and IIC (Fig. 6A). Furthermore, the soleus muscles in diabetic GK rats had a higher percentage of low-oxidative type I fibers than those in nondiabetic Wistar rats. The

fast plantaris muscles in diabetic GK rats had an increased percentage of low-oxidative type IIB fibers and a decreased percentage of high-oxidative type IIA fibers when compared to the corresponding percentages in WR and GK-Ex rats (Fig. 6C). These results correspond to those obtained in previous studies using nonobese GK (Gu *et al.* 2010, Nagatomo *et al.* 2011a, Yasuda *et al.* 2002, Yasuda *et al.* 2007) and obese OLETF (Nagatomo *et al.* 2018, Yasuda *et al.* 2001) rats.

Animals and humans with type 2 diabetes show a low percentage of high-oxidative fibers and a high percentage of low-oxidative fibers in skeletal muscles; a high percentage of low-oxidative fibers, especially type IIB fibers, was observed in the vastus lateralis muscles of patients with type 2 diabetes (Gaster *et al.* 2001, Märin *et al.* 1994, Nyholm *et al.* 1997, Oberbach *et al.* 2006). Obese Zucker rats with type 2 diabetes had a lower percentage of high-oxidative fibers in the soleus muscle than nonobese Zucker rats without type 2 diabetes, whereas obese Zucker rats without type 2 diabetes had

a normal fiber type distribution (Adachi *et al.* 2007). The altered percentages in skeletal muscles are directly linked to blood glucose levels, but not to insulin levels or body weight, because the growth-related increase in blood glucose levels in rats with type 2 diabetes paralleled the increased percentage of low-oxidative type I fibers in the soleus muscle (Yasuda *et al.* 2002). Thus, the altered patterns of fiber types in the skeletal muscles of rats with type 2 diabetes are tightly linked to impaired glucose metabolism.

Voluntary running exercise inhibited the type shift from high-oxidative fibers to low-oxidative fibers in the plantaris muscle of diabetic GK rats (Fig. 6C). Inhibition of the fiber type shift induced by voluntary running exercise is thought to be due to enhanced oxidative metabolism in cells and tissues. However, we did not elucidate the reason why the diabetes-induced alteration in fiber type distribution was shifted to normal in the fast plantaris muscle, but not in the slow soleus muscle, by voluntary running exercise.




Fig. 8. Succinate dehydrogenase (SDH) activity (**A, B**) and peroxisome proliferator-activated receptor γ co-activator-1 α (*Pgc-1 α*) mRNA levels (**C, D**) in the soleus and plantaris muscles of the Wistar (WR), Goto-Kakizaki (GK), and exercised Goto-Kakizaki (GK-Ex) groups. Values are expressed as mean \pm SD (n=5 for WR and GK; n=10 for GK-Ex). ^ap<0.05 compared with WR; ^bp<0.05 compared with GK.

Voluntary running, muscle SDH activity, and *Pgc-1 α* mRNA levels

Diabetic GK rats showed decreased muscle SDH activity (Fig. 8A, B) and *Pgc-1 α* mRNA levels (Fig. 8C, D) when compared with nondiabetic WR rats. PGC-1 α is a member of a family of transcription coactivators that plays central roles in the regulation of oxidative metabolism in cells and tissues (Liang and Ward 2006, Southgate *et al.* 2007). In skeletal muscles, PGC-1 α accelerates mitochondrial synthesis and promotes muscle fiber remodeling (Mortensen *et al.* 2006, Wu *et al.* 1999).

Transgenic mice with increased PGC-1 α expression in skeletal muscles showed an increased number of mitochondria and a fiber type shift, i.e. more oxidative and less glycolytic fibers (Miura *et al.* 2003, Miura *et al.* 2006). The decrease in oxidative activity is due to low *Pgc-1 α* mRNA levels, which may be induced by hyperglycemia. Diabetic GK rats exhibited hyperglycemia (Figs 2 and 3A), but not hypertriglyceridemia (Fig. 3E) or hyperinsulinemia (Fig. 3F). Obese diabetic OLETF rats also showed lower muscle oxidative capacity than nondiabetic Long-Evans Tokushima Otsuka (LETO)

rats (Yasuda *et al.* 2001), but OLETF rats also exhibited hyperglycemia, hypertriglyceridemia, and hyperinsulinemia. Both diabetic GK and OLETF rats exhibit hyperglycemia. Therefore, the decreased oxidative capacity in the skeletal muscles of diabetic rats is thought to be due to hyperglycemia, but not hypertriglyceridemia or hyperinsulinemia. We concluded that the diabetes-induced decrease in SDH activity was inhibited by voluntary running exercise, which effectively maintained and improved oxidative metabolism in skeletal muscles.

Relationships between voluntary running distance and the abovementioned parameters

We observed that the running distances of individual rats were negatively related to HbA1c (Fig. 9A) and total cholesterol (Fig. 9B) levels and positively related to SDH activity (Fig. 9C) and *Pgc-1α* mRNA levels (Fig. 9D) in the soleus muscle, indicating that type 2 diabetes is effectively inhibited when diabetic GK rats ran longer distances. Voluntary running exercise, which is different from forced exercise, such as treadmill

running, is effective as an exercise program because of the low stress.

A previous study (Yasuda *et al.* 2006) observed a negative relationship between voluntary running distance and the percentage of low-oxidative type I fibers in the soleus muscle of 10 OLETF rats. Furthermore, five OLETF rats with voluntary running distances <7000 m/day had no high-oxidative type IIA fibers in the soleus muscle, whereas five OLETF rats with running distances >7000 m/day showed a positive relationship between voluntary running distance and the percentage of type IIA fibers in the soleus muscle. In this study, seven diabetic GK rats had no high-oxidative type II fibers (Fig. 5E, F) after voluntary running exercise, whereas three had high-oxidative type IIA and type IIC fibers as well as low-oxidative type I fibers in the soleus muscle (Fig. 5G, H) after voluntary running exercise. However, there was no relationship between voluntary running distance and the fiber type percentages in the soleus and plantaris muscles in diabetic GK rats.




Fig. 9. Relationships between voluntary running distance and the HbA1c (**A**) and total cholesterol (**B**) levels, succinate dehydrogenase (SDH) activity (**C**), and peroxisome proliferator-activated receptor γ coactivator-1α (*Pgc-1α*) mRNA levels (**D**) in the soleus muscles of exercised Goto-Kakizaki rats ($n=10$). Running distance was negatively related to both HbA1c ($r=-0.81$, $p<0.05$) and total cholesterol ($r=-0.83$, $p<0.05$) levels and positively related to both SDH activity ($r=0.98$, $p<0.05$) and *Pgc-1α* mRNA levels ($r=0.70$, $p<0.05$).

Conclusions

Growth-related increases in fasting blood glucose and HbA1c levels and decreases in the percentage of low-oxidative fibers and SDH activity in

the skeletal muscles of diabetic rats were inhibited by voluntary running exercise. The observed inhibition of the decreased muscle oxidative capacity induced by voluntary running exercise was thought to be due to increased *Pgc-1α* mRNA levels, which are influenced by

blood glucose levels.

HbA1c levels were negatively related to running distance, and SDH activity and *Pgc-1α* mRNA levels in the soleus muscle were positively related to running distance in the diabetic rats, indicating that the onset and progression of type 2 diabetes are effectively inhibited in nonobese diabetic rats that ran longer distances.

Conflict of Interest

There is no conflict of interest.

References

- ADACHI T, KIKUCHI N, YASUHARA K, ANAHARA R, GU N, MATSUNAGA T, YAMAMURA T, MORI C, TSUJIMOTO G, TSUDA K, ISHIHARA A: Fibre type distribution and gene expression levels of both succinate dehydrogenase and peroxisome proliferator-activated receptor- γ coactivator-1 α of fibres in the soleus muscle of Zucker diabetic fatty rats. *Exp Physiol* 92: 449-455, 2007. <https://doi.org/10.1113/expphysiol.2006.035451>
- FU S, MENG Y, ZHANG W, WANG J, HE Y, HUANG L, CHEN H, KUANG J, DU H: Transcriptomic responses of skeletal muscle to acute exercise in diabetic Goto-Kakizaki rats. *Front Physiol* 10: 872, 2019. <https://doi.org/10.3389/fphys.2019.00872>
- GASTER M, STAEBER P, BECK-NIELSEN H, SCHRØDER HD, HANDBERG A: GLUT4 is reduced in slow muscle fibers of type 2 diabetic patients: is insulin resistance in type 2 diabetes a slow, type 1 fiber disease? *Diabetes* 50: 1324-1329, 2001. <https://doi.org/10.2337/diabetes.50.6.1324>
- GOTO Y, KAKIZAKI M, MASAKI N: Spontaneous diabetes produced by selective breeding of normal Wistar rats. *Proc Jpn Acad* 51: 80-85, 1975. <https://doi.org/10.2183/pjab1945.51.80>
- GOTO Y, KAKIZAKI M, MASAKI N: Production of spontaneous diabetic rats by repetition of selective breeding. *Tohoku J Exp* 119: 85-90, 1976. <https://doi.org/10.1620/tjem.119.85>
- GU N, NAGATOMO F, FUJINO H, TAKEDA I, TSUDA K, ISHIHARA A: Hyperbaric oxygen exposure improves blood glucose level and muscle oxidative capacity in rats with type 2 diabetes. *Diabetes Technol Ther* 12: 125-133, 2010. <https://doi.org/10.1089/dia.2009.0104>
- ISHIHARA A, NAGATOMO F, FUJINO H, KONDO H, TSUDA K: Lifestyle-related disease and skeletal muscle: a review. *J Phys Fitness Sports Med* 1: 17-27, 2012. <https://doi.org/10.7600/jpfsm.1.17>
- ISHIHARA A, ROY RR, OHIRA Y, IBATA Y, EDGERTON VR: Hypertrophy of rat plantaris muscle fibers after voluntary running with increasing loads. *J Appl Physiol* 84: 2183-2189, 1998. <https://doi.org/10.1152/jappl.1998.84.6.2183>
- JENKINS DW, JENKS A: Exercise and diabetes: a narrative review. *J Foot Ankle Surg* 56: 968-974, 2017. <https://doi.org/10.1053/j.jfas.2017.06.019>
- KAWANO K, HIRASHIMA T, MORI S, SAITO Y, KUROSUMI M, NATORI T: Spontaneous long-term hyperglycemic rat with diabetic complications. Otsuka Long-Evans Tokushima Fatty (OLETF) strain. *Diabetes* 41: 1422-1428, 1992. <https://doi.org/10.2337/diabetes.41.11.1422>
- KIM SS, KOO JH, KWON IS, OH YS, LEE SJ, KIM EJ, KIM WK, LEE J, CHO JY: Exercise training and selenium or a combined treatment ameliorates aberrant expression of glucose and lactate metabolic proteins in skeletal muscle in a rodent model of diabetes. *Nutr Res Pract* 5: 205-213, 2011. <https://doi.org/10.4162/nrp.2011.5.3.205>
- LIANG H, WARD WF: PGC-1 α : a key regulator of energy metabolism. *Adv Physiol Educ* 30: 145-151, 2006. <https://doi.org/10.1152/advan.00052.2006>

Acknowledgements

This study was supported by a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science.

Authors Contributions

Isuzu Nakamoto and Akihiko Ishihara conceived, designed, and performed the experiments, analyzed the data, and wrote the paper.

- LIN J, WU H, TARR PT, ZHANG CY, WU Z, BOSS O, MICHAEL LF, PUIGSERVER P, ISOTANI E, OLSON EN, LOWELL BB, BASSEL-DUBY R, SPIEGELMAN BM: Transcriptional co-activator PGC-1 α drives the formation of slow-twitch muscle fibres. *Nature* 418: 797-801, 2002. <https://doi.org/10.1038/nature00904>
- MACIA M, PECCHI E, DESROIS M, LAN C, VILMEN C, PORTHA B, BERNARD M, BENDAHAN D, GIANNESINI B: Exercise training impacts exercise tolerance and bioenergetics in gastrocnemius muscle of non-obese type-2 diabetic Goto-Kakizaki rat in vivo. *Biochimie* 148: 36-45, 2018. <https://doi.org/10.1016/j.biochi.2018.02.014>
- MÅRIN P, ANDERSSON B, KROTKIEWSKI M, BJÖRNTORP P: Muscle fiber composition and capillary density in women and men with NIDDM. *Diabetes Care* 17: 382-386, 1994. <https://doi.org/10.2337/diacare.17.5.382>
- MIURA S, KAI Y, ONO M, EZAKI O: Overexpression of peroxisome proliferator-activated receptor γ coactivator-1 α (PGC-1 α) down-regulates GLUT4 mRNA in skeletal muscles. *J Biol Chem* 278: 31385-31390, 2003. <https://doi.org/10.1074/jbc.M304312200>
- MIURA S, TOMITSUKA E, KAMEI Y, YAMAZAKI T, KAI Y, TAMURA M, KITA K, NISHINO I, EZAKI O: Overexpression of peroxisome proliferators-activated receptor γ co-activator-1 α leads to muscle atrophy with depletion of ATP. *Am J Pathol* 169: 1129-1139, 2006. <https://doi.org/10.2353/ajpath.2006.060034>
- MORIFUJI T, MURAKAMI S, FUJITA N, KONDO H, FUJINO H: Exercise training prevents decrease in luminal capillary diameter of skeletal muscles in rats with type 2 diabetes. *Sci World J* 2012: 645891, 2012. <https://doi.org/10.1100/2012/645891>
- MORTENSEN OH, FRANDSEN L, SCHJERLING P, NISHIMURA E, GRUNNET N: PGC-1 α and PGC-1 β have both similar and distinct effects upon myofiber switching toward an oxidative phenotype. *Am J Physiol Endocrinol Metab* 291: E807-E816, 2006. <https://doi.org/10.1152/ajpendo.00591.2005>
- NAGATOMO F, FUJINO H, KONDO H, GU N, TAKEDA I, ISHIOKA N, TSUDA K, ISHIHARA A: PGC-1 α mRNA level and oxidative capacity of the plantaris muscle in rats with metabolic syndrome, hypertension, and type 2 diabetes. *Acta Histochem Cytochem* 44: 73-80, 2011a. <https://doi.org/10.1267/ahc.10041>
- NAGATOMO F, FUJINO H, KONDO H, KOUZAKI M, GU N, TAKEDA I, TSUDA K, ISHIHARA A: The effects of running exercise on oxidative capacity and PGC-1 α mRNA levels in the soleus muscle of rats with metabolic syndrome. *J Physiol Sci* 62: 105-114, 2012a. <https://doi.org/10.1007/s12576-011-0188-1>
- NAGATOMO F, FUJINO H, KONDO H, SUZUKI H, KOUZAKI M, TAKEDA I, ISHIHARA A: PGC-1 α and FOXO1 mRNA levels and fiber characteristics of the soleus and plantaris muscles in rats after hindlimb unloading. *Histol Histopathol* 26: 1545-1553, 2011b. <https://doi.org/10.14670/HH-26.1545>
- NAGATOMO F, FUJINO H, KONDO H, TAKEDA I, TSUDA K, ISHIHARA A: High-fat diet-induced reduction of peroxisome proliferator-activated receptor- γ coactivator-1 α mRNA levels and oxidative capacity in the soleus muscle of rats with metabolic syndrome. *Nutr Res* 32: 144-151, 2012b. <https://doi.org/10.1016/j.nutres.2011.12.015>
- NAGATOMO F, GU N, FUJINO H, TAKEDA I, TSUDA K, ISHIHARA A: Skeletal muscle characteristics of rats with obesity, diabetes, hypertension, and hyperlipidemia. *J Atheroscler Thromb* 16: 576-585, 2009. <https://doi.org/10.5551/jat.1065>
- NAGATOMO F, TAKEMURA A, ROY RR, FUJINO H, KONDO H, ISHIHARA A: Mild hyperbaric oxygen inhibits the growth-related decline in skeletal muscle oxidative capacity and prevents hyperglycemia in rats with type 2 diabetes mellitus. *J Diabetes* 10: 753-763, 2018. <https://doi.org/10.1111/1753-0407.12666>
- NYHOLM B, QU Z, KAAL A, PEDERSEN SB, GRAVHOLT CH, ANDERSEN JL, SALTIN B, SCHMITZ O: Evidence of an increased number of type IIb muscle fibers in insulin-resistant first-degree relatives of patients with NIDDM. *Diabetes* 46: 1822-1828, 1997. <https://doi.org/10.2337/diabetes.46.11.1822>
- OBERBACH A, BOSSENZ Y, LEHMANN S, NIEBAUER J, ADAMS V, PASCHKE R, SCHÖN MR, BLÜHER M, PUNKT K: Altered fiber distribution and fiber-specific glycolytic and oxidative enzyme activity in skeletal muscle of patients with type 2 diabetes. *Diabetes Care* 29: 895-900, 2006. <https://doi.org/10.2337/diacare.29.04.06 dc05-1854>
- PHIELIX E, MENSIN M: Type 2 diabetes mellitus and skeletal muscle metabolic function. *Physiol Behav* 94: 252-258, 2008. <https://doi.org/10.1016/j.physbeh.2008.01.020>

- PUIGSERVER P: Tissue-specific regulation of metabolic pathways through the transcriptional coactivator PGC-1 α . *Int J Obes* 29 (Suppl 1): S5-S9, 2005. <https://doi.org/10.1038/sj.ijo.0802905>
- QI Z, HE J, ZHANG Y, SHAO Y, DING S: Exercise training attenuates oxidative stress and decreases p53 protein content in skeletal muscle of type 2 diabetic Goto-Kakizaki rats. *Free Rad Biol Med* 50: 794-800, 2011. <https://doi.org/10.1016/j.freeradbiomed.2010.12.022>
- SCHULER M, ALI F, CHAMBON C, DUTEIL D, BORNERT JM, TARDIVEL A, DESVERGNE B, WAHLI W, CHAMBON P, METZGER D: PGC1 α expression is controlled in skeletal muscles by PPAR β , whose ablation results in fiber-type switching, obesity, and type 2 diabetes. *Cell Metab* 4: 407-414, 2006. <https://doi.org/10.1016/j.cmet.2006.10.003>
- SOUTHGATE RJ, NEILL B, PRELOVSEK O, EL-OSTA A, KAMEI Y, MIURA S, EZAKI O, McLOUGHLIN TJ, ZHANG W, UNTERMAN TG, FEBBRAIO MA: FOXO1 regulates the expression of 4E-BP1 and inhibits mTOR signaling in mammalian skeletal muscle. *J Biol Chem* 282: 21176-21186, 2007. <https://doi.org/10.1074/jbc.M702039200>
- TAKEMURA A, ISHIHARA A: Mild hyperbaric oxygen inhibits growth-related decrease in muscle oxidative capacity of rats with metabolic syndrome. *J Atheroscler Thromb* 24: 26-38, 2017. <https://doi.org/10.5551/jat.34686>
- WENDE AR, HUSS JM, SCHAEFFER PJ, GIGUÈRE V, KELLY DP: PGC-1 α coactivates PDK4 gene expression via the orphan nuclear receptor ERR α : a mechanism for transcriptional control of muscle glucose metabolism. *Mol Cell Biol* 25: 10684-10694, 2005. <https://doi.org/10.1128/MCB.25.24.10684-10694.2005>
- WU H, KANATOUS SB, THURMOND FA, GALLARDO T, ISOTANI E, BASSEL-DUBY R, WILLIAMS RS: Regulation of mitochondrial biogenesis in skeletal muscle by CaMK. *Science* 296: 349-352, 2002. <https://doi.org/10.1126/science.1071163>
- WU Z, PUIGSERFVER P, ANDERSSON U, ZHANG C, ADELMANT G, MOOTHA V, TROY A, CINTI S, LOWELL B, SCARPULLA RC, SPIEGELMAN BM: Mechanisms controlling mitochondrial biogenesis and respiration through the thermogenic coactivator PGC-1. *Cell* 98: 115-124, 1999. [https://doi.org/10.1016/S0092-8674\(00\)80611-X](https://doi.org/10.1016/S0092-8674(00)80611-X)
- YASUDA K, ADACHI T, GU N, MATSUMOTO A, MATSUNAGA T, TSUJIMOTO G, TSUDA K, ISHIHARA A: Effects of hyperbaric exposure with high oxygen concentration on glucose and insulin levels and skeletal muscle-fiber properties in diabetic rats. *Muscle Nerve* 35: 337-343, 2007. <https://doi.org/10.1002/mus.20692>
- YASUDA K, ADACHI T, KIKUCHI N, TSUJIMOTO G, AOKI N, TSUDA K, ISHIHARA A: Effects of running exercise on fibre-type distribution of soleus and plantaris muscles in diabetic Otsuka Long-Evans Tokushima fatty rats. *Diabetes Obes Metab* 8: 311-321, 2006. <https://doi.org/10.1111/j.1463-1326.2005.00507.x>
- YASUDA K, ISHIHARA A, ADACHI T, SHIHARA N, SEINO Y, TSUDA K: Growth-related changes in skeletal muscle fiber type and insulin resistance in diabetic Otsuka Long-Evans Tokushima Fatty rats. *Acta Histochem Cytochem* 34: 371-382, 2001. <https://doi.org/10.1267/ahc.34.371>
- YASUDA K, NISHIKAWA W, IWANAKA N, NAKAMURA E, SEINO Y, TSUDA K, ISHIHARA A: Abnormality in fibre type distribution of soleus and plantaris muscles in non-obese diabetic Goto-Kakizaki rats. *Clin Exp Pharmacol Physiol* 29: 1001-1008, 2002. <https://doi.org/10.1046/j.1440-1681.2002.03757.x>