

## REVIEW

---

# Jan Evangelista Purkyně and the Cerebellum Then and Now

F. VOŽEH<sup>1,2</sup>

<sup>1</sup>Department of Pathophysiology, Faculty of Medicine in Pilsen, Charles University, Pilsen, Czech Republic, <sup>2</sup>Laboratory of Neurodegenerative Disorders, Biomedical Centre, Faculty of Medicine in Pilsen, Charles University, Pilsen, Czech Republic

Received March 31, 2015

Accepted June 25, 2015

On-line December 15, 2015

---

### Summary

The name of Jan Evangelista Purkyně and the cerebellum belong inseparably together. He was the first who saw and described the largest nerve cells in the brain, *de facto* in the cerebellum. The most distinguished researchers of the nervous system then showed him the highest recognition by naming these neurons as Purkinje cells. Through experiments by J. E. Purkyně and his followers properly functionally was attributed to the cerebellum share in precision of motor skills. Despite ongoing and fruitful research, after a relatively long time, especially in the last two decades, scientists had to constantly replenish and re-evaluate the traditional conception of the cerebellum and formulate a new one. It started in the early 1990s, when it was found that cerebellar cortex contains more neurons than the cerebral cortex. Shortly thereafter it was gradually revealed that such enormous numbers of neural cells are not without an impact on brain functions and that the cerebellum, except its traditional role in the motor skills, also participates in higher nervous activity. These new findings were obtained thanks to the introduction of modern methods of examination into the clinical praxis, and experimental procedures using animal models of cerebellar disorders described below.

### Key words

Animal models • Ataxia • Cerebellum • Neurodegenerations • Purkinje cells

### Corresponding author

F. Vožeh, Department of Pathophysiology, Faculty of Medicine in Pilsen, Charles University, Alej Svobody 1655/76, 323 00 Plzeň, Czech Republic. E-mail: frantisek.vozeh@lfp.cuni.cz

### Introduction

Jan Evangelista Purkyně was the most formidable figure in Czech physiology on the world who brought many scientific discoveries into the biological, physiological and biomedical sciences. Among the systems to which he paid attention it was mainly the nervous system and especially the cerebellum. J. E. Purkyně was the first scientist to see and describe the largest nerve cells in the brain, *de facto* in the cerebellum. He hand-illustrated them at his memorable lecture to the professional public in Prague in 1837. Moreover, Purkyně depicted and precisely described the individual three layers of the cerebellar cortex when the largest neurons were named ganglial bodies (Druga 1986). Later, S. R. y Cajal (1911), one of the most distinguished researchers of the nervous system, recommended in his monumental monography to designate these large neurons according an author who described them for the first time. Since that time also other researchers used this eponyme and the Purkinje cells (in English-language texts) represent one of the most cited scientific terms in the neuroscientific literature and the name Purkyně is inseparably connected with the cerebellum. On the basis of experiments with animals of many species (sheep, rabbits, chickens, pigeons, ortolans, fish) and also by examination of human brains J. E. Purkyně (Valentin 1836, Druga 1986, Rokyta 2011) and later his followers (Friede 1963, Eccles *et al.* 1967, Szentagothai 1968, Ito 1970, 1972, 1984, Palay and Chan-Palay 1974) attributed to the function of the cerebellum a share in the control and precision of motor skills. Despite their enormous

scientific discoveries the cerebellum resisted for some time further knowledge even though it prepared for another surprise. The first revelation came in the early 1990s, when it was found that cerebellar cortex contains more neurons than the cerebral cortex (Glickstein 1992) and that the cerebellum has bidirectional connections with almost all significant structures of the brain (Schmahmann 1996, Schmahmann and Pandya 1997, Middleton and Strick 2001, Ito 2006). Shortly thereafter it was revealed that such an enormous number of nerve cells as well as functionally highly potential cerebellar connections exert some impact on brain functions. Shortly thereafter it was confirmed that the cerebellum, except its traditional role in the precision of motor skills, also participates in the higher nervous functions (Manto and Lorivel 2011, Buckner 2013).

### Functional and morphological characteristics of the cerebellum

The cerebellum is a brain structure well known even to people not educated in biology or biomedical science. This is first of all thanks to its beautiful texture which resembles the Biblical tree of life or *Arbor vitae* (in Latin). And, of course, Purkinje cells with their shrubby dendrites belong to the most beautiful creations of living nature of all.

As a part of the hindbrain (rhombencephalon) the cerebellum is localized in the posterior part of the cranial cavity, behind the pons and medulla oblongata. The surface of the cerebellum is formed by vermis, paravermal zones and hemispheres. From a functionally-developmental point of view, the cerebellum consists of three lobes. Flocculonodular lobe (archicerebellum, vestibulocerebellum), smallest and phylogenetically oldest component of the cerebellum, situated at its bottom part. It receives afferentation from the vestibular nerve and nuclei, and is responsible for balance and posture.

The anterior lobe (paleocerebellum, spinocerebellum) is a phylogenetically younger part of the cerebellum and represents 1/3 of its hemispheres. Afferentation comes mainly from the spinal cord and enables the spinocerebellum to be responsible for muscular tonus.

The posterior lobe (neocerebellum, pontocerebellum), the largest and developmentally youngest part of the cerebellum with afferentation from the neocortex, is responsible for the coordination of movements, motor stereotypes and also participates in

higher nervous activity. The bidirectional communication of the cerebellum with the brainstem and other parts of the CNS provide the three pairs of the cerebellar peduncle: the inferior, which contains both afferent and efferent fibers; the middle, with afferent fibers coming from the nuclei pontis, and the superior which consists mainly from efferent fibers from the deep cerebellar nuclei (Kandel *et al.* 2000).

The cerebellar cortex is composed of gray matter, which is largely folding so that the cerebellar surface is formed by transverse folia (which in sagittal sections resemble above mentioned *Arbor vitae*). This is the reason that 85 % of the surface is hidden, despite the size of this area being fully comparable to the cortical area of the big brain. The cerebellar cortex consists of three layers viewed from the surface: the molecular layer, the Purkinje cell layer and the granule cell layer (deepest) (Kandel *et al.* 2000). The input to the cerebellum comes *via* the mossy fibers and *via* the climbing fibers. Both of these are glutamatergic (excitatory). Nevertheless, synapses between cerebellar cortical interneurons and Purkinje and granule cells are GABAergic (inhibitory). Also the output from Purkinje cells to the deep cerebellar nuclei is GABAergic. Therefore, the functional effect of the cerebellum on the other part of the CNS is inhibitory (Danbolt *et al.* 2004).

Under the gray matter the central part of the cerebellum is composed of white matter – medulla, which consists of nervous fibers. Inside of the cerebellar hemispheres there are four pairs of cerebellar nuclei embedded in the medullary center. From the midline to the lateral side there are: the fastigial nucleus, the globose nucleus, emboliform nucleus and the largest of them, the dentate nucleus. The input to the cerebellar nuclei comes from the Purkinje cells (they represent the only output from the cerebellar cortex) and from sources outside the cerebellum (pontocerebellar, spinocerebellar, and olivocerebellar fibers, together with fibers from reticular nuclei). Most of these afferents are collaterals of fibers proceeding to the cerebellar cortex. A few rubrocerebellar fibers end in the globose and emboliform nucleus, and the fastigial nucleus receives afferents from the vestibular nerve and nuclei. The outputs from the cerebellar nuclei go to the brain stem (from fastigial nucleus through the inferior cerebellar peduncle descending through the reticular formation and the lateral vestibular nuclei to the spinal cord and ascending through ventrolateral nucleus of the thalamus to the primary motor cortex). Efferents from the other nuclei leave the cerebellum through the

superior peduncle and terminate in the brain stem and neurons of the ventral lateral nucleus of the thalamus projecting to the limb areas of motor cortex (Barr and Kiernan 1988, Kandel *et al.* 2000).

Although the human cerebellum represents only 1/10 of the whole brain volume, recent research has shown, that number of neurons in this structure represents 80 % of all neurons in the brain. Except humans, the same situation is valid for agouti and galago, whereas e.g. in the mouse the cerebellum holds 60 %, in rat, guinea pig and macaque it represents 70 % of all brain neurons (Herculano-Houzel 2010). Because such huge functional potential was not surprising, that except its traditional role in maintaining balance, posture control and motor function, the cerebellum participates in cognition, language, learning, memory and emotions which is a core of the new cerebellar conception (Buckner 2013, Schutter 2013). Within the framework of this conception, it is possible to sum up that the cerebellum receives sensory and predictive stimuli, which include signals from receptors of muscles, tendons and joints (proprioceptors). These then integrate with motor commands so that it leads to precise control of movements and proper modulation of cognitive-emotional activities (Schutter 2013).

### **The cerebellar disorders and methods of examination and research of the cerebellum**

The cerebellar disorders include mainly motor deficits but also mental and behavioral abnormalities.

*Inhibitory cerebellar syndrome* as a consequence of the destruction of this structure (due to injuries, ischemia, hemorrhage, tumors, inflammation, intoxication and inherited neurodegenerative conditions) is characterized by:

- increased muscle passivity and weakness, intention tremor and ataxia
- asynergy and hypermetry of movements (lack of movements coordination and overreaching the intended object or goal)
- adiadochokinesia (inability to perform rapid alternating movements)
- speech disorders (slow, incoherent, slurred) (Kandel *et al.* 2000).

Mental and behavioral abnormalities in the cerebellar disorders are known in humans as *cognitive-affective syndrome*, which is typical with:

- deficit in the formation of words,

- diminished ability of planning and abstraction,
- disorder of working memory,
- altered personality with blunted emotions and inappropriate behavior (Schmahmann and Sherman 1997).

There is also evidence for a probable pathogenetic connection with schizophrenia (findings of small vermis and asymmetric hemispheres in these patients) and with autism (Andreasen and Pierson 2008, Fatemi *et al.* 2012).

Less frequent are the cerebellar disorders characterized by the so-called *irritative cerebellar syndrome*, which represents abnormalities in the sense of hyperfunction due to pathologic irritation. It is characterized by:

- muscle hypertonia
- resting tremor
- flexion posture
- hypokinesia.

These symptoms resemble the hypertonic-hypokinetic syndrome in parkinsonism (Záhlava 1994, Kishore *et al.* 2014). It is useful to stress, that unilateral cerebellar lesions cause functional disorders on the ipsilateral side in contrast to cerebral defects when motor abnormalities are contralateral. The reason is crossing of both the cerebellar (decussation of the superior cerebellar peduncle) and cerebral (cross in the rubrospinal and corticospinal systems) pathways (Kandel *et al.* 2000).

It is obvious that the knowledge for formulation of the new conception of the cerebellum would not be obtained without incredible effort of clinical and experimental researchers, and of course, thanks to modern devices as well as the introduction of new special clinical and laboratory methods put into praxis.

#### *Clinical and neuro-imaging procedures*

Clinical medical examination of the cerebellar function is made mostly in persons suspected of alcohol intake as a complement to the blood alcohol determination or/and as a preliminary neurological examination before further diagnostic procedures. It involves the assessment of posture and gait with eyes closed, coordination and accuracy of movements, quality of speech and handwriting. Further, common studies with psychologists on cerebellum-injured patients have reported e.g. deficits in generating words, performing tasks requiring anticipatory planning and the occurrence of dysgraphia and agrammatism (Fiez *et al.* 1992, Silveri *et al.* 1994, Schmahmann 1996, Leggio *et al.* 2008). Very

significant progress both in diagnostics and research was then done thanks to the introduction of modern imaging methods e.g. PET and MRI (positron emission tomography and magnetic resonance imaging). Using these methods, for example, the participation of the cerebellar dentate nucleus in tasks related to planning was proved in healthy subjects tested in procedural learning abilities (Kim *et al.* 1994, Van Mier *et al.* 1998, 2002, Sharma *et al.* 2015).

## The experimental research on animals

Experiments on animals play irreplaceable role in neuroscience research. In particular the use of animal models of cerebellar disorders has brought valuable insights into the physiology and pathophysiology of the cerebellum (Akita and Arai 2009, Manto and Marmolino 2009, Ebner and Chen 2013). The first was the confirmed participation of the cerebellum in different types of learning and memory (Bickford 1993, Lalonde *et al.* 1993a,b, 1996, Cendelín *et al.* 2008, Hilber and Caston 2001, Křížková and Vožeh 2004, Thullier *et al.* 1997, Vožeh *et al.* 1997, 1999, 2001). Also some pathogenetic mechanisms of cerebellar disorders were clarified including the possibilities of influencing them (e.g. by means of non-invasive procedures, forced motor activity, enriched environment, pharmacologically or by transplantation of embryonic cerebellar tissue as well as stem cells). It was possible thanks to a battery of sophisticated behavioral tests, electrophysiological investigation together with use not only classic neuro-histological procedures but also modern immunofluorescence methods (Cendelín and Vožeh 2013, Purkartová and Vožeh 2013).

There are two groups of these models of various cerebellar dysfunctions, known also in human beings, used in this research:

1. mouse models derived from naturally arising mutations in genome of individuals that were then further bred and prepared for experimental research,
2. mouse models that were developed by means of genetical engineering procedures as transgenic animals which have to be treated according to rules for the manipulation with genetically modified organisms (GMO).

## Naturally arising animal models

### *Lurcher mutant mice*

Lurcher mice represent the first described

mutant animal of the 1st group (Phillips 1960), gradually derived from four strains (DBA, C57B6, C3H, CBA). The mice suffer from olivocerebellar degeneration due to semi-dominant spontaneous mutation *Grid2<sup>Lc</sup>* in the gene for delta 2 glutamate receptor localized on chromosome 6 (Zuo *et al.* 1997, De Jager *et al.* 1997). Heterozygous animals (+/Lc) exhibit almost complete postnatal loss of Purkinje cells and significantly decreased the number of granule cells and inferior olive neurons (ION) during 3 postnatal months (PM). The death of Purkinje cells is a type of excitotoxic apoptosis (Norman *et al.* 1995, Zuo *et al.* 1997, Selimi *et al.* 2000, Purkartová and Vožeh 2013) due to an inborn mutation. Nevertheless, some features are present which indicate necrotic cell death (Dumesnil-Bousez and Sotelo 1992, Dusart *et al.* 2006) and autophagy (Wang *et al.* 2006). It means that multiple Purkinje cell death pathways play role in Lurchers (Nishiyama and Yuzaki 2010, Zanjani *et al.* 2013). The degeneration of granule cells and ION is a consequence of the lost target of their axons (Purkinje cells) similarly, as described below, decrease of cerebellar interneurons (Golgi, stellate and basket cells) (Zanjani *et al.* 2006). Homozygous Lurcher mice die shortly after birth due to massive loss of brain-stem neurons during late embryogenesis (Resibois *et al.* 1997). Homozygous wild type littermates of Lurcher mutants (+/+) represent about one half of newborns in the litter. They are completely healthy and serve as ideal controls. Degeneration of the deep cerebellar nuclei is relatively mild in Lurchers (Heckroth 1994, Sultan *et al.* 2002). Functionally, the Lurchers suffer from multiple neural problems. Except ataxia, which is more or less a common symptom of all cerebellar ataxic mice, they exhibit higher excitability, impairment of learning and memory, higher stress response and worse maternal behavior (Lalonde *et al.* 1988, Lalonde and Thifault 1994, Frederic *et al.* 1997, Tuma *et al.* 2013, Caston *et al.* 1998, Hilber *et al.* 2004). Further, the Lurcher mice exhibit a lower resistance of ION against neurotoxic 3-acetylpyridine already before the Purkinje cell degeneration starts (Caddy and Vožeh 1997). There were also described changes in classical eyelid conditioning in Lurchers (Porrás-García *et al.* 2005, 2010). Nevertheless, they are capable of learning to a certain degree both cognitive and motor tasks. This was proven in spatial memory using the Morris water maze but also with some motor tests whereby significant improvement was observed in trained animals (Lalonde *et al.* 1993a,b, 1996, Hilber and Caston 2001, Křížková and Vožeh 2004). Also the positive effect of forced motor activity was evident in motor and cognitive functions in

general, and especially in the impact on age dependent decline of these abilities (Cendelín *et al.* 2008, Cendelín and Vožeh 2013).

#### *Purkinje cell degeneration mice*

Purkinje cell degeneration (*pcd*) mutant mice are similar to Lurchers, characterized by autosomal recessive *Agtbp1pcd/J* mutation in the gene encoding cytosolic ATP/GTP binding protein 1 (cytosolic carboxypeptidase-like protein, CCP1) on chromosome 13 (resulting in lack of this protein) (Mullen *et al.* 1976, Fernandez-Gonzalez *et al.* 2002). Therefore the *pcd* mice are homozygous. They exhibit almost a complete loss of Purkinje cells at the end of the first postnatal month (PM) (Mullen *et al.* 1976) and their death is of apoptotic origin together with autophagy (Kyuhou *et al.* 2006, Chakrabarti *et al.* 2009, Berezniuk and Fricker 2010). The animals suffer also from progressive extinction of 90 % of the granule cells (between PM 3-20 %) because of the loss of their target – Purkinje cells (Ghetti *et al.* 1987, Triarhou 2010). Probably, due to the missing input from Purkinje cells there is also about a 20 % reduction of the deep cerebellar nuclei (in PM 10) (Triarhou *et al.* 1987). At PM 10 there is also a significant decrease (almost 50 %) of ION as a consequence of the lost Purkinje cells, their target. Except for these defects, these mice are affected by degeneration of retinal photoreceptors (Blanks *et al.* 1982, Blanks and Spee 1992, LaVail *et al.* 1982), bulb mitral olfactory neurons and selected thalamic neurons (O’Gorman and Sidman 1985). In addition, there is also defective spermatogenesis, which results in the *pcd* males being sterile (Kim *et al.* 2011). Except the cerebellar ataxia, *pcd* mice exhibit subtle habitus and poor health. Functionally they show worse performance in the rotarod, coat-hanger tests but also in stationary beam (Le Marec and Lalonde 1997). In the Morris water maze they are unable to successfully navigate to the hidden platform compared to reaching a visible goal where they are better (Goodlett *et al.* 1992). The *pcd* mice are impaired in the delay eye-blink classical conditioning when compared with wild type controls (Chen *et al.* 1996), while in trace eye-blink conditioning there were no differences between them (Brown *et al.* 2009). The results indicate that the essential neural circuitry for trace conditioning bypasses the cerebellar cortex and differs from the one for classical eye blink conditioning.

#### *Staggerer mice*

The Staggerers, similarly like the *pcd* mice

suffer from an autosomal recessive mutation (*Rorasg*) in the gene encoding the retinoid-related orphan receptor alpha on chromosome 9 (Hamilton *et al.* 1996). They are characterized by a staggering gait, hypotony tremor, smaller body size and except for Purkinje cells also granule cells and ION degeneration is present. At the end of PM 1, 60-90 % of Purkinje cells are missing (in dependence on region), likewise more than 90 % of granule cells and about 60 % of ION (Landis and Sidman 1978, Herrup and Mullen 1979a,b, Blatt and Eisenman 1985a,b). Despite a 30 % reduction in the volume of the deep cerebellar nuclei, the number of neurons remains unchanged (Roffler-Tarlov and Sidman 1978, Roffler-Tarlov and Herrup 1981). The development of Purkinje cells in Staggerer mice is characterized by a delayed growth of their spines and surviving Purkinje cells exhibit smaller somata and dendritic trees (Landis and Sidman 1978, Herrup and Mullen 1979a,b).

#### *Hot-foot mutant mice*

Hot foot mice are characterized by autosomally recessive mutation in the gene for delta 2 glutamate receptor (*Grid 2ho*) localized on chromosome 6 which is identical with the one in Lurcher mutants (Lalouette *et al.* 1998). They display defective presynaptic innervation of Purkinje cells due to ectopic spine and partial necrotic loss of granule cells. They suffer from ataxia with typically irregular movements of hind limbs (as walking on a hot plate) (Guastavino *et al.* 1990). Hot-foot mutants further exhibit poor performance in the rotating grid, wooden beam, coat-hanger, rotarod tests and also in Morris water maze (Lalonde *et al.* 1996). Nevertheless, they are capable of partial learning in the rotating grid and coat hanger tests (Lalonde *et al.* 1995, 1996, 2003).

#### *Nervous mice*

Nervous mice represent autosomal recessive mutants who suffer from severe degeneration of Purkinje cells and ION. The nervous mutation (*nr*) is located on chromosome 8 (Campbell and Hess 1996, De Jager *et al.* 1998). Purkinje cells exhibit abnormally rounded mitochondria (Landis 1973) and 90 % of them undergo to necrotic degeneration. About 10 % of Purkinje cells survive. Nevertheless 1/3 of ION die by retrograde degeneration because of the loss of their target. Nervous mutants are typical with ataxia and hyperactivity (Lalonde and Strazielle 2003) and they display bad performance in the beam, coat-hanger and rotarod tests. In the Morris water maze, they fail in experiments with

the hidden platform but not in those with a visible goal (Lalonde and Strazielle 2003). Nervous mice also suffer from retinal regeneration which is in early postnatal period prompt but then slows down with an almost complete loss of photoreceptors in PM 17 (Mullen and LaVail 1975, LaVail *et al.* 1993).

#### *Weaver mice*

Weaver mice are affected by autosomal semidominant Grikv mutation in the gene (encoding a G-protein) located on chromosome 2 (Patil *et al.* 1995). The consequences of the mutation result in an impairment of the cerebellar and some extracerebellar structures. In contrast to other mutants described, there mainly cerebellar granule cells suffer from apoptotic degeneration, while Purkinje cells are relatively unimpaired (Migheli *et al.* 1995, Wullner *et al.* 1995). Weaver homozygous have 72 % and heterozygous up to 86 % of the normal number of Purkinje cells when ION in both of them are actually intact (Blatt and Eisenman 1985a,b). On the other hand, homozygous Weaver mutants exhibit 23-25 % decrease in the number of deep cerebellar nuclei neurons (Maricich *et al.* 1997). Except the cerebellum there were still further described: the degeneration of dopaminergic neurons in the substantia nigra together with a decreased concentration of dopamine in the nucleus caudatus and in the striatum (Roffler-Tarlov and Graybiel 1984). Morphological changes (thickening of the pyramidal cell layer) were observed in the hippocampus of Weaver mice. From the functional point of view the Weaver failed in the spatial learning in the Morris water maze, and they also had lower exploratory activity as well as worse performance on the wooden beam and grid tests. Just like some other cerebellar mutant mice (e.g. pcd mice) the Weaver homozygous males are sterile due to the death of germ cells in their testes.

#### *Reeler mice*

Reeler mice are typical for autosomal recessive mutation in the gene encoding extracellular matrix protein reelin, which is important for neural cell migration (D'Arcangelo *et al.* 1995). This mutation located on chromosome 5 leads to abnormal migration of neurons during brain development (Beckers *et al.* 1994). There are therefore ectopic cell localizations not only in the cerebellum (Hamburgh 1963, Terashima *et al.* 1983), but also in some other brain structures (hippocampus, neocortex, inferior olive, olfactory bulb, superior

colliculus, substantia nigra) (Mikoshiba *et al.* 1980, Wyss *et al.* 1980, Kang *et al.* 2010). In homozygous Reeler mice, the cerebellum is reduced in size (Mariani *et al.* 1977) with Purkinje cells reduced to less than half the normal amount, but only about 5 % of them are localized normally. About 10 % of Purkinje cells are placed within the granular layer while, some granule cells are located in the molecular layer (Terashima *et al.* 1985, Heckroth *et al.* 1989). Also the inferior olivary complex is reduced in size by 22.6 %. Despite all these abnormalities, the specificity of the cerebellar connections is mostly preserved (Mariani *et al.* 1977). Functionally the Reeler mice exhibit poor performance in the active avoidance task, water maze, wooden beam, coat-hanger and rotarod tests (Lalonde *et al.* 2004). The lower neurogenesis rates together with a higher inclination to ischemic brain injury and epilepsy were described in Reeler mice (Patrylo *et al.* 2006, Won *et al.* 2006). Because of special behavioral abnormalities in heterozygous Reeler mice, some authors suggest considering them as a model of schizophrenia (Costa *et al.* 2002, Schmitt *et al.* 2013).

#### *Scrambler mice*

Scrambler mice are characterized by the spontaneous autosomal recessive scrambler mutation (Dab1scm-3J) in the gene on chromosome 4 (Sweet *et al.* 1996). The cerebellum is reduced in size already in 1M old homozygous mice. The number of Purkinje cells is decreased and only 5 % of them are normally located. The number of granule cells is reduced by 80 %. Despite these abnormalities, the cerebellar afferent systems are preserved (Goldowitz *et al.* 1997). Homozygous Scrambler mice display ataxia and whole body tremor as early as the 2nd postnatal week. From the functional point of view, they exhibit poor performance in the rotarod, wooden beam and coat-hanger tests. In the Morris water maze test of spatial navigation, they failed in both variants with the submerged or visible platform (Jacquelin *et al.* 2012, 2013).

#### *Toppler mutant mice*

Toppler mice suffer from autosomal recessive mutation in a gene on chromosome 8. The dramatic loss of Purkinje cells with typical abnormalities of their dendritic trees is evident already from 2nd postnatal week. Besides Purkinje cells there is present also degeneration of Bergmann glia and these both processes are evidently in a functional relationship. The Toppler mutants exhibit serious ataxia, which worsens with age

and leads to a shorter lifespan (8-12 months). They represent a useful model for investigation of the developmental interaction of Purkinje cells and Bergmann glia (Duchala *et al.* 2004).

#### *Pogo mutant mice*

Pogo mice are naturally occurring mutants derived from an inbred strain of the Korean wild mouse. The autosomal recessive mutation affects a gene, which is located on chromosome 8.

The mice show ataxia and difficulty in maintaining their posture. It is further characterized by a wobbly gait and a tendency to fall over, which appear at about 2 postnatal weeks and continue throughout life without progression. Morphologically in Pogo mice is the extent loss of Purkinje cells with ectopic spines, emanating from their primary dendrites (Lee and Jeong 2009). In addition, parallel fiber varicosities were larger than in the control mice and a single fiber often established synaptic contacts with up to four dendritic spines of a Purkinje cell. These multiple synapses were observed in both the cerebellar vermis and the hemispheres. Moreover, condensed and vacuolated granule cells nuclei were observed in the granular layer of the ataxic Pogo but not in the control cerebella. These morphological defects suggest that there could be massive synaptic abnormalities between Purkinje cells and the cerebellar afferent pathways in the Pogo cerebellum (Lee *et al.* 2011).

### **Transgenic animal models**

Animals of this group represent mainly transgenic mice arising from the same type of hereditary determined neurodegeneration in humans resulting from repeating CAG triplets in the genetic information. Besides Huntington disease also dentatorubral-pallidoluysian atrophy, spinal and bulbar muscular atrophy, Niemann-Pick disease, Friedreich ataxia and also 7 types of spinocerebellar ataxias (SCA 1, 2, 3, 6, 7, 17, 23) from more than many scores of those in humans belong to this group (Yamada *et al.* 2008, Cendelin 2014). The aforementioned triplet codes for polyglutamine (polyQ) in each respective gene, resulting in mutants to the expansion of these sections mutated proteins. The severity of disability depends on the number of CAG triplet repeats (Koeppen 2005, Yamada *et al.* 2008, Fan *et al.* 2014). In any case, these transgenic animals represent more precise models of the above-

mentioned diseases.

#### *Mouse models of human hereditary cerebellar ataxias*

##### *SCA1*

Spinocerebellar ataxia type 1 (SCA1) represents one of the autosomal dominant hereditary ataxias. SCA1 causes about 3 to 16 % of autosomal dominant cerebellar ataxias. In addition to ataxia, SCA1 is associated with the difficulty of speaking, swallowing, impaired cognition and increased reflexes are also common. SCA1 usually starts in the mid-30s and progresses more rapidly than other SCA subtypes (Manto 2005). It is caused by an enlarged region of CAG repeat in the gene for ataxin-1 which results in the expansion of the polyQ tract in the ataxin-1 protein. Normally the length is between 6-35 repeats, while in affected people it is 39-83 (Taroni *et al.* 2013). To understand the pathophysiology of this disease, the transgenic mice expressing the human SCA1 gene were generated with either a normal or an expanded CAG tract. Both transgenes were stable in parents to offspring transmission and therefore six transgenic lines of mice for both normal or an expanded CAG tract were obtained. While all six transgenic lines expressing the unexpanded human allele had normal Purkinje cells, in transgenic animals from five of six lines with the expanded SCA1 allele (carrying 82 CAG repeats), ataxia and Purkinje cell degeneration developed (Burrigh *et al.* 1995). There is a marked loss of Purkinje cells, and those surviving have abnormal dendritic trees. Bergmann glia proliferation, a shrunk molecular layer with gliosis and ectopic Purkinje cells in it and also in the granular layer are further typical findings in the SCA1 mice cerebellar cortex. Ataxia with shorter strides start at PM 3, with worse performance on the rotarod test observed during PM 2 and serious gait abnormalities subsequently at PM 12 (Clark *et al.* 1997). The data obtained contributed to the clarification of some important pathogenetic mechanisms of SCA1 as a role of nuclear localization of the pathological ataxin-1 (Klement *et al.* 1998), its proteasomal degradation (Cummings 1999) as well as the effect of expanded ataxin-1 in the downregulation of genes acting in signal transduction and calcium homeostasis prior to manifestation of the pathology (Lin *et al.* 2000).

##### *SCA2*

SCA2 type of the autosomal dominant spinocerebellar ataxia causes about 6 to 18 % of all types of this disease. Besides progressive ataxia SCA2 is

characterized by progressive ataxia, dysarthria, posture tremor, slow saccades, hyporeflexia of the upper limbs, autonomic dysfunction, sleep disturbances, ophthalmoparesis, dementia, and parkinsonism (Huynh *et al.* 2000). The onset age varies from 2 to 65 years, and onset before the age of 20 years correlates with a more aggressive disease course. SCA2 is caused by another CAG trinucleotide repeat, this time encoding a protein called ataxin 2. Normal ataxin-2 usually contains 22 or 23 glutamines, while SCA2 patients have 32-77 repeats. SCA2 transgenic mice generated in several lines differ in lengths of the CAG repeat segment and vary in severity of symptoms of this disease. Those with 58 CAG repeats are characterized by about 50 % reduction of Purkinje cell numbers in PM 6. In animals from all three generated lines, SCA2 mice exhibited a reduced stride length to the end of PM 4. Some interesting findings in animals of two lines were observed in the rotarod test when 6 weeks old transgenic animals did not differ in their performance from wild type controls, while SCA2 homozygotes and heterozygotes failed by PM 4 or PM 6 respectively (Huynh *et al.* 2000). From the above mentioned phenomena it is evident that a more severe manifestation is in mouse-lines with prolonged polyQ tracts. It is in agreement with the facts obtained in humans when prolonged CAG repeats characterized an earlier onset of symptoms and vice versa. Research performed on transgenic SCA2 animals has brought important knowledge about the pathogenetic mechanisms of this disease. It was shown that nuclear localization and inclusion body formation of ataxin-2 was not necessary for disease development. (Huynh *et al.* 2000). Further glutamate induced death of Purkinje cells and its partial prevention by dantrolene was proved. This a calcium-ion stabilizer used in long-term treatment of SCA2 mice also alleviated motor deficits and Purkinje cell loss. These observations confirmed calcium signaling role in the pathogenesis of SCA2 (Liu *et al.* 2009).

### SCA3

SCA3 type of spinocerebellar ataxia is better known as Machado-Joseph disease. SCA3 is characterized by progressive cerebellar ataxia, areflexia, peripheral amyotrophy, muscle atrophy, parkinsonian features, dystonia, and spasticity (Bettencourt *et al.* 2008). However, some minor presentations, such as external progressive ophthalmoplegia are also present. Its onset varies from 5 to 75 years of age and it represents around 1 % from all SCA cases. In normal alleles, CAG

trinucleotides repeat length varies between 12 and 44, in SCA3 patients, the length is 52-86 (Taroni *et al.* 2013). SCA3 transgenic mice were generated with polyQ tracts of 64, 67, 72, 76 and 84 repeats (Cemal *et al.* 2002). The severity of symptoms depends on the number of CAG repeats. Therefore, in the mice expressing ataxin-3 with an expanded polyglutamine tract with 79 repeats suffer except from a wide gait, tremor and lower activity also from progressive ataxia starting at PM 5-6 despite the fact that the loss of cerebellar neurons is mild (Chou *et al.* 2008, 2011). These findings suggest that the decline in cerebellar function is more dependent on down-regulation of cerebellar expressions of proteins and neuronal dysfunction than on Purkinje cell loss (Chou *et al.* 2011). Research results obtained from experiments on SCA3 mice contributed to the understanding of more pathogenetic mechanisms of this disease mainly by pointing out the role of altered voltage-activated potassium channels (Shakkottai *et al.* 2011), calcium dependent calpain-type proteases (Hubener *et al.* 2013) and the disruption of dendritic development as well as metabotropic glutamate receptor signaling in Purkinje cells by mutant ataxin-3 protein (Konno *et al.* 2014).

### SCA6

The spinocerebellar ataxia type 6 (SCA6) is characterized by CAG polyQ repeat expansion in the CACNA1A gene encoding the alpha 1A-voltage-dependent calcium channel (CaV2.1) (Zhuchenko *et al.* 1997) and it represents the most frequent type of all SCAs (from 5 to 18 %). SCA6 is characterized by a slowly progressive ataxia, dysarthria, intention tremor, gaze-evoked and/or downbeat nystagmus, dysphagia, positional vertigo, and sensory, pyramidal, and extrapyramidal motor deficits (Stevanin *et al.* 1997). Normal alleles have 4-18 repeats, while alleles in diseased people contain 20-33 polyQ repeats. SCA6 generated transgenic mice which have the 84 polyQ repeats tract are characterized by progressive motor abnormalities and the aggregation of the pathological protein CACNA1A. Homozygous animals have shown hypoactivity and worse performance on the accelerating rotarod (Watase *et al.* 2008). Heterozygotes did not differ from wild type mice according to visual assessment up to PM 20 despite performing worse than controls on the rotarod test at PM 19. Regardless of motor disturbances no neuronal loss and morphological changes on Purkinje cells were observed in these mice at PM 20. From the point of view of pathophysiological mechanisms, the



observations in SCA6 mice with 28 or 84 polyQ repeats suggest that alteration of CaV2.1 channel properties does not play any role in the SCA6 pathology which may be linked to the accumulation of mutant channels (Saegusa *et al.* 2007, Watase *et al.* 2008).

#### SCA7

Spinocerebellar ataxia type 7 (SCA7) is a neurodegenerative disease caused by the expansion of a CAG repeat within the gene encoding ataxin-7. The normal range is 7-19 repeats. Pathological alleles contain from 37 to more than 400 CAG triplets. SCA7, with the onset in childhood, is characterized by progressive ataxia, macular or retinal degeneration with visual loss, slow saccades, ophthalmoplegia, dysphagia, somatosensory and neuropsychiatric impairment (David *et al.* 1998). Among SCAs, it is the only one characterized by severe ataxia, and at the same time by blindness. (Taroni *et al.* 2013). Generated transgenic mice with 90 polyQ repeats have nuclear inclusions of mutant protein ataxin-7 in Purkinje cells and photoreceptors, and exhibit both abnormal motor coordination together with vision impairment. Nevertheless, there are big differences in the spectrum of findings in dependence on the polyQ tract length. For instance the mice with 266 CAG repeats suffer from an infantile form of SCA7 characterized by ataxia, visual impairment, abnormal short-term synaptic potentiation and untimely death (Yoo *et al.* 2003). On the other hand, the mice with 92 polyQ repeats and ataxin-7 expression in all CNS neurons, except Purkinje cells, display severe degeneration just of these cells, ataxic gait disorder and typical ataxin-7 nuclear aggregates which correlate with visible symptoms including an earlier death (Garden *et al.* 2002). Conversely, transgenic mice with 52 CAG repeats suffer from ataxia, without a significant loss of Purkinje cells. All these findings suggest that the Purkinje cell degeneration in SCA7 is not cell-autonomous and that in the pathogenesis of this disease yet other factors play a role, for instance changes in gene expression with an impact on glutamatergic transmission, signal transduction, myelin formation, axonal transport neuronal differentiation and glial function (Chou *et al.* 2010). This hypothesis was confirmed by observation when the suppression of the mutant gene (to 50 %), one month after the ataxia started, was effective in delaying motor disorders, reducing ataxin-7 pathological protein aggregation in Purkinje cells and preventing the synaptic loss between Purkinje cells and the climbing fibers (Furrer *et al.* 2013).

#### SCA17

SCA17 type of spinocerebellar ataxia late-onset degenerative disorder is caused by an expanded polyQ repeat in the TATA-box-binding protein (TBP). The disease is characterized by progressive gait and limb ataxia, seizure, cognitive dysfunction, neuropsychiatric impairment, and pyramidal and extrapyramidal features such as spasticity, dystonia, chorea, and Parkinson's disease. While healthy people have 23-43 CAG polyQ repeats, patients suffering from this pathology have 45-63 repeats polyQ tract. The disease is very rare, i.e. below 1 % of all SCAs (Lasek *et al.* 2006). Despite that, two mouse models of this disease were generated (Friedman *et al.* 2007, Chang *et al.* 2011), it is only type of SCA when a rat TBPQ64 model SCA17 has shown to be more advantageous. Similarly as in mouse models for SCA17, TBPQ64 rats show a severe neurological symptomatology including ataxia, impairment of postural reflexes and hyperactivity in the early stages followed by a reduced activity, loss of body weight, and early death. Neuropathological findings showed neuronal loss, particularly in the cerebellum, degeneration of Purkinje, basket, and stellate cells as well as, changes in the morphology of the dendrites. In addition, nuclear TBP-positive immunoreactivity and axonal torpedoes were found by light and electron microscopy. It was also shown that some crucial characteristics of SCA17 are better mirrored in TBP rats than in existing mouse models. The use of this model for the first time by means of PET and diffusion tensor imaging (DTI) were also replicated recent PET studies in human SCA17 patients. These results further confirmed that DTI are potentially useful correlates of neuropathological changes in TBPQ64 rats and raise the hope that DTI imaging could provide a biomarker for SCA17 patients (Kelp *et al.* 2013).

#### SCA23

Spinocerebellar ataxia type 23 (SCA23) is an adult-onset extremely rare neurodegenerative disorder characterized by slow progressive gait and limb ataxia, with variable additional features, including peripheral neuropathy and dysarthria (Bakalkin *et al.* 2010). In the literature a Dutch family was reported with autosomal dominant late-onset spinocerebellar ataxia affecting at least 13 members spanning 3 generations. Only 1 patient, with a disease duration of 23 years, was wheelchair-bound. The MRI of that patient showed severe cerebellar atrophy with memory deficits beginning around age 50.

*Postmortem* examination of one patient displayed frontotemporal atrophy, atrophy of the cerebellar vermis, pons, and spinal cord. There was also neuronal loss in the cerebellar vermis, dentate nuclei, and inferior olives, but not in the pons. Moreover, there was thinning of the cerebellopontine tracts and demyelination of the posterior and lateral columns of the spinal cord. SCA23 maps to chromosome region 20p12.3-p13 and missense mutations in the prodynorphin PDYN gene appear to cause the disease. (Verbeek 2004). Prodynorphin knock-out mice are more sensitive to noxious stimuli but have normal responses to non-noxious stimuli (Wang *et al* 2001); additionally, mutant dynorphin proteins have enhanced non-opioid excitatory activities which may undermine the development of SCA23 (Wang *et al.* 2013). Nevertheless, these mice have not yet been tested for phenotypic similarity to human SCA23 (Cendelin 2014).

## Conclusions

As mentioned above, a new conception of the cerebellum including a more detailed knowledge about physiology and pathophysiology of the cerebellum as well as about the cerebellar disorders would not be possible without the above-surveyed clinical and research methods including sophisticated procedures on experimental animals. First of all, both naturally arising and also artificially by genetic engineering prepared animal models created the basis for such enormous progress in cerebellar research. Because of these advances it is indisputable that studies mainly on cerebellar mutants provided a lot of valuable information about cerebellar function, manifestation of cerebellar disorders as well as pathogenesis and therapy, though there are also some limitations that prevent a direct translation of findings to humans.

Though the basic manifestation of cerebellar dysfunction is similar in most of the mouse models of cerebellar degenerations, some particular signs differ. Moreover, some of the mutations have extra-cerebellar impacts, which are integral components of the phenotype (e.g. olfactory bulb, retina and thalamus degeneration in *pcd* mice etc.). There are, of course, species differences in anatomy, metabolism, behavior, etc. between mice and men. Spontaneous mouse mutations are usually not identical to human ones and therefore mouse diseases can only be similar to human diseases. On the other hand, transgenic mice carry human pathological alleles and thus

can be used as models for specific human diseases.

Nevertheless, even in transgenic mice the action of the mutation could significantly differ from the natural human mutation. The transgene could possibly be under the control of different promoters and its expression could differ from that seen in humans regarding intensity and cell type. The transgene usually does not replace the mouse wild type allele, if it is not knocked-out. In these cases the mouse has both pathological as well as fully expressed normal gene products. Autosomal dominant ataxias appear in humans mostly in the form of heterozygotes. However, mice can also be studied as homozygous individuals. These experiments have shown that homozygous mice often display more severe pathological phenotype than heterozygous mice suggesting that the diseases might be more accurately described as having a semi-dominant nature. Despite the limitations, cerebellar mutant mice are invaluable tools for research, when the goal is a better understanding the pathogenesis of cerebellar degenerative disorders and, hopefully, finding effective therapies for humans. Cerebellar mutant mice will continue to serve as valuable tools in preclinical studies investigating the therapeutic methods for treating human cerebellar degenerations.

Nevertheless, deep phenotypic characterizations, especially those of the new transgenic mouse models, and the elucidation of the pathogenesis and relationship of the functional disorders to the cerebellum will remain important. Verification of the conformity of the mouse models with human diseases on the morphological, functional and molecular level is also crucial for translation of experimental research to human medicine and thereby aid in its progress.

Finally, it is possible to wish our neuroscience research that it succeed, at least partially, as in the period of the J. E. Purkyně.

## Conflict of Interest

There is no conflict of interest.

## Acknowledgements

The article was supported by the Charles University Research Fund (project number P36), the Faculty of Medicine in Pilsen and by project CZ.1.05/2.1.00/03.0076 from the European Regional Development Fund. The author would like to thank to Mr. Christopher Koy, M. A., Ph.D., for language editing.

## References

- AKITA K, ARAI S: The ataxic Syrian hamster: an animal model homologous to the *pcd* mutant mouse? *Cerebellum* **8**: 202-210, 2009.
- ANDREASEN NC, PIERSON R: The role of the cerebellum in schizophrenia. *Biol Psychiatry* **64**: 81-88, 2008.
- BAKALKIN G, WATANABE H, JEZIERSKA J, DEPOORTER C, VERSCHUUREN-BEMELMANS C, BAZOV I, ARTEMENKO KA, YAKOVLEVA T, DOOIJES D, VAN DE WARRENBURG BP, ZUBAREV RA, KREMER B, KNAPP PE, HAUSER KF, WIJMENGA C, NYBERG F, SINKE RJ, VERBEEK DS: Prodynorphin mutations cause the neurodegenerative disorder spinocerebellar ataxia type 23. *Am J Hum Genet* **87**: 593-603, 2010.
- BARR ML, KIERNAN JA: The cerebellum. In: *The Human Nervous System. An Anatomical Viewpoint, 5th ed.* J. B. Lippincott Company, Philadelphia, 1988, pp 160-177.
- BECKERS MC, BAR I, HUYNH-THU T, DERNONCOURT C, BRUNIALTI AL, MONTAGUTELLI X, GUÉNET JL, GOFFINET AM: A high-resolution genetic map of mouse chromosome 5 encompassing the *reeler* (*rl*) locus. *Genomics* **23**: 685-690, 1994.
- BEREZNIUK I, FRICKER LD: A defect in cytosolic carboxypeptidase 1 (*Nna1*) causes autophagy in Purkinje cell degeneration mouse brain. *Autophagy* **6**: 558-559, 2010.
- BETTENCOURT C, SANTOS C, KAY T, VASCONCELOS J, LIMA M: Analysis of segregation patterns in Machado-Joseph disease pedigrees. *J Hum Genet* **53**: 920-923, 2008.
- BICKFORD P: Motor learning deficits in aged rats are correlated with loss of cerebellar noradrenergic function. *Brain Res* **620**: 133-138, 1993.
- BLANKS JC, SPEE C: Retinal degeneration in the *pcd/pcd* mutant mouse accumulation of spherules in the interphotoreceptor space. *Exp Eye Res* **54**: 637-644, 1992.
- BLANKS JC, MULLEN RJ, LAVAIL MM: Retinal degeneration in the *pcd* cerebellar mutant mouse. II. Electron microscopic analysis. *J Comp Neurol* **212**: 231-246, 1982.
- BLATT GJ, EISENMAN LM: A qualitative and quantitative light microscopic study of the inferior olivary complex in the adult staggerer mutant mouse. *J Neurogenet* **2**: 51-66, 1985a.
- BLATT GJ, EISENMAN LM: A qualitative and quantitative light microscopic study of the inferior olivary complex of normal, *reeler*, and *weaver* mutant mice. *J Comp Neurol* **232**: 117-128, 1985b.
- BROWN KL, AGELAN A, WOODRUFF-PAK DS: Unimpaired trace classical eyeblink conditioning in Purkinje cell degeneration (*pcd*) mutant mice. *Neurobiol Learn Mem* **93**: 303-311, 2010.
- BUCKNER RL: The cerebellum and cognitive function: 25 years of insight from anatomy and neuroimaging. *Neuron* **80**: 807-815, 2013.
- BURRIGHT EN, CLARK HB, SERVADIO A, MATILLA T, FEDDERSEN RM, YUNIS WS, DUVICK LA, ZOGHBI HY, ORR HT: *SCA1* transgenic mice: a model for neurodegeneration caused by an expanded CAG trinucleotide repeat. *Cell* **82**: 937-948, 1995.
- CADDY KW, VOŽEH F: The effect of 3-acetylpyridine on inferior olivary neuron degeneration in Lurcher mutant and wild-type mice. *Eur J Pharmacol* **330**: 139-142, 1997.
- CAJAL SR: *Histologie du système nerveux de l'homme et des vertébrés*. T. II. Ed. A Maloine, Paris, 1911, pp 1-993.
- CAMPBELL DB, HESS EJ: Chromosomal localization of the neurological mouse mutations tottering (*tg*), Purkinje cell degeneration (*pcd*), and nervous (*nr*). *Brain Res Mol Brain Res* **37**: 79-84, 1996.
- CASTON J, CHIANALE C, DELHAYE-BOUCHAUD N, MARIANI J: Role of the cerebellum in exploration behavior. *Brain Res* **808**: 232-237, 1998.
- CEMAL CK, CARROLL CJ, LAWRENCE L, LOWRIE MB, RUDDLE P, AL-MAHDAWI S, KING RH, POOK MA, HUXLEY C, CHAMBERLAIN S: YAC transgenic mice carrying pathological alleles of the *MJD1* locus exhibit a mild and slowly progressive cerebellar deficit. *Hum Mol Genet* **11**: 1075-1094, 2002.
- CENDELÍN J: From mice to men: lessons from mutant ataxic mice. *Cerebellum Ataxias* **1**: 4, 2014.
- CENDELÍN J, VOŽEH F: Lurcher mouse. In: *Handbook of the Cerebellum and Cerebellar Disorders*. MANTO M, GRUOL DL, SCHMAHMANN JD, KOIBUCHI N, ROSSI F (eds), Springer Science+Business Media, New York, 2013, pp 1499-1520.

- CENDELÍN J, KORELUSOVÁ I, VOŽEH F: The effect of repeated rotarod training on motor skills and spatial learning ability in Lurcher mutant mice. *Behav Brain Res* **189**: 65-74, 2008.
- CHAKRABARTI L, ENG J, IVANOV N, GARDEN GA, LA SPADA AR: Autophagy activation and enhanced mitophagy characterize the Purkinje cells of pcd mice prior to neuronal death. *Mol Brain* **2**: 24, 2009.
- CHANG YC, LIN CY, HSU CM, LIN HC, CHEN YH, LEE-CHEN GJ, SU MT, RO LS, CHEN CM, HSIEH-LI HM: Neuroprotective effects of granulocyte-colony stimulating factor in a novel transgenic mouse model of SCA17. *J Neurochem* **118**: 288-303, 2011.
- CHEN L, BAO S, LOCKARD JM, KIM JK, THOMPSON RF: Impaired classical eyeblink conditioning in cerebellar-lesioned and Purkinje cell degeneration (pcd) mutant mice. *J Neurosci* **16**: 2829-2838, 1996.
- CHOU AH, YEH TH, OUYANG P, CHEN YL, CHEN SY, WANG HL: Polyglutamine-expanded ataxin-3 causes cerebellar dysfunction of SCA3 transgenic mice by inducing transcriptional dysregulation. *Neurobiol Dis* **31**: 89-101, 2008.
- CHOU AH, CHEN CY, CHEN SY, CHEN WJ, CHEN YL, WENG YS, WANG HL: Polyglutamine-expanded ataxin-7 causes cerebellar dysfunction by inducing transcriptional dysregulation. *Neurochem Int* **56**: 329-339, 2010.
- CHOU AH, CHEN SY, YEH TH, WENG YH, WANG HL: HDAC inhibitor sodium butyrate reverses transcriptional downregulation and ameliorates ataxic symptoms in a transgenic mouse model of SCA3. *Neurobiol Dis* **41**: 481-488, 2011.
- CLARK HB, BURRIGHT EN, YUNIS WS, LARSON S, WILCOX C, HARTMAN B, MATILLA A, ZOGHBI HY, ORR HT: Purkinje cell expression of a mutant allele of SCA1 in transgenic mice leads to disparate effects on motor behaviors, followed by a progressive cerebellar dysfunction and histological alterations. *J Neurosci* **17**: 7385-7395, 1997.
- COSTA E, DAVIS J, PESOLD C, TUETING P, GUIDOTTI A: The heterozygote reeler mouse as a model for the development of a new generation of antipsychotics. *Curr Opin Pharmacol* **2**: 56-62, 2002.
- CUMMINGS CJ, REINSTEIN E, SUN Y, ANTALFFY B, JIANG Y, CIECHANOVER A, ORR HT, BEAUDET AL, ZOGHBI HY: Mutation of the E6-AP ubiquitin ligase reduces nuclear inclusion frequency while accelerating polyglutamine-induced pathology in SCA1 mice. *Neuron* **24**: 879-892, 1999.
- D'ARCANGELO G, MIAO GG, CHEN SC, SOARES HD, MORGAN JI, CURRAN T: A protein related to extracellular matrix proteins deleted in the mouse mutant reeler. *Nature* **374**: 719-723, 1995.
- DANBOLT NC, HOLMSETH S, SKÁR A, LEHRE KP, FURNESS DN: Glutamate uptake and transporters. In: *Excitotoxicity in Neurological Disease – New Therapeutic Challenge*. C FERRARESE, MF BEAL (eds), Kluwer Academic Publishers, Boston Dordrecht London, 2004, pp 27-49.
- DAVID G, DURR A, STEVANIN G, CANCEL G, ABBAS N, BENOMAR A, BELAL S, LEBRE, AS, ABADA-BENDIB M, GRID D, HOLMBERG M, YAHYAOU M, HENTATI F, CHKILI T, AGID Y, BRICE A: Molecular and clinical correlations in autosomal dominant cerebellar ataxia with progressive macular dystrophy (SCA7). *Hum Mol Genet* **7**: 165-170, 1998.
- DE JAGER PL, ZUO J, COOK SA, HEINTZ N: A new allele of the lurcher gene, lurcherJ. *Mamm Genome* **8**: 647-650, 1997.
- DE JAGER PL, HARVEY D, POLYDORIDES AD, ZUO J, HEINTZ N: A high-resolution genetic map of the nervous locus on mouse chromosome 8. *Genomics* **48**: 346-353, 1998.
- DRUGA R: Purkinje cells of the cerebellum. (in Czech) In: *Jan Evangelista Purkyně, Life and Work*. E. TRÁVNÍČKOVÁ (ed), Avicenum, Prague, 1986, pp 255-261.
- DUCHALA CS, SHICK HE, GARCIA J, DEWESE DM, SUN X, STEWART VJ, MACKLIN WB: The toppler mouse: a novel mutant exhibiting loss of Purkinje cells. *J Comp Neurol* **476**: 113-129, 2004.
- DUMESNIL-BOUSEZ N, SOTELO C: Early development of the Lurcher cerebellum: Purkinje cell alterations and impairment of synaptogenesis. *J Neurocytol* **21**: 506-529, 1992.
- DUSART I, GUENET JL, SOTELO C: Purkinje cell death: differences between developmental cell death and neurodegenerative death in mutant mice. *Cerebellum* **5**: 163-173, 2006.
- EBNER TJ, CHEN G: Tottering mouse. In: *Handbook of the Cerebellum and Cerebellar Disorders*. MANTO M, GRUOL DL, SCHMAHMANN JD, KOIBUCHI N, ROSSI F (eds), Springer Science+Business Media, New York, 2013, pp 1521-1540.

- ECCLES JC, ITO M, SZENTHAGOTHAI J: *The Cerebellum as a Neuronal Machine*. Springer, Berlin, 1967.
- FAN HC, HO LI, CHI CS, CHEN SJ, PENG GS, CHAN TM, LIN SZ, HARN HJ: Polyglutamine (PolyQ) diseases: genetics to treatments. *Cell Transplant* **23**: 441-458, 2014.
- FATEMI SH, ALDINGER KA, ASHWOOD P, BAUMAN ML, BLAHA CD, BLATT GJ, ET AL.: Consensus paper: pathological role of the cerebellum in autism. *Cerebellum* **11**: 777-807, 2012.
- FERNANDEZ-GONZALEZ A, LA SPADA AR, TREADAWAY J, HIGDON JC, HARRIS BS, SIDMAN RL, MORGAN JI, ZUO J: Purkinje cell degeneration (pcd) phenotypes caused by mutations in the axotomy-induced gene, *Nna1*. *Science* **295**: 1904-1906, 2002.
- FIEZ JA, PETERSEN SE, CHENEY MK, RAICHLE ME: Impaired non-motor learning and error detection associated with cerebellar damage: a single case study. *Brain* **115**: 155-178, 1992.
- FREDERIC F, CHAUTARD T, BROCHARD R, CHIANALE C, WOLLMAN E, OLIVER C, DELHAYE-BOUCHAUD N, MARIANI J: Enhanced endocrine response to novel environment stress and endotoxin in Lurcher mutant mice. *Neuroendocrinology* **66**: 341-347, 1997.
- FRIEDE RL: The relationship of body size, nerve cell size, axon length, and glial density in the cerebellum. *Proc Natl Acad Sci U S A* **49**: 187-193, 1963.
- FRIEDMAN MJ, SHAH AG, FANG ZH, WARD EG, WARREN ST, LI S, LI XJ: Polyglutamine domain modulates the TBP-TFIIB interaction: implications for its normal function and neurodegeneration. *Nat Neurosci* **10**: 1519-1528, 2007.
- FURRER SA, WALDHERR SM, MOHANACHANDRAN MS, BAUGHN TD, NGUYEN KT, SOPHER BL, DAMIAN VA, GARDEN GA, LA SPADA AR: Reduction of mutant ataxin-7 expression restores motor function and prevents cerebellar synaptic reorganization in a conditional mouse model of SCA7. *Hum Mol Genet* **22**: 890-903, 2013.
- GARDEN GA, LIBBY RT, FU YH, KINOSHITA Y, HUANG J, POSSIN DE, SMITH AC, MARTINEZ RA, FINE GC, GROTE SK, WARE CB, EINUM DD, MORRISON RS, PTACEK LJ, SOPHER BL, LA SPADA AR: Polyglutamine-expanded ataxin-7 promotes non-cell-autonomous Purkinje cell degeneration and displays proteolytic cleavage in ataxic transgenic mice. *J Neurosci* **22**: 4897-4905, 2002.
- GHETTI B, NORTON J, TRIARHOU LC: Nerve cell atrophy and loss in the inferior olivary complex of "Purkinje cell degeneration" mutant mice. *J Comp Neurol* **260**: 409-422, 1987.
- GLICKSTEIN M: The cerebellum and motor learning. *Cur Opin Neurobiol* **2**: 802-806, 1992.
- GOLDOWITZ D, CUSHING RC, LAYWELL E, D'ARCANGELO G, SHEIDON M, SWEET HO, DAVISSON M, STEINDLER D, CURRAN T: Cerebellar disorganization characteristic of reeler in scrambler mutant mice despite presence of reelin. *J Neurosci* **17**: 8767-8777, 1997.
- GOODLETT CR, HAMRE KM, WEST JR: Dissociation of spatial navigation and visual guidance performance in Purkinje cell degeneration (pcd) mutant mice. *Behav Brain Res* **47**: 129-141, 1992.
- GUASTAVINO JM, SOTELO C, DAMEZ-KINSELLE I: Hot-foot murine mutation: behavioral effects and neuroanatomical alterations. *Brain Res* **523**: 199-210, 1990.
- HAMBURGH M: Analysis of the postnatal developmental effects of "reeler", a neurological mutation in mice. A study in developmental genetics. *Dev Biol* **8**: 165-185, 1963.
- HAMILTON BA, FRANKEL WN, KERREBROCK AW, HAWKINS TL, FITZHUGH W, KUSUMI K, RUSSELL LB, MUELLER KL, VAN BERKEL V, BIREN BW, KRUGLYAK L, LANDER ES: Disruption of the nuclear hormone receptor RORalpha in staggerer mice. *Nature* **379**: 736-739, 1996.
- HECKROTH JA: Quantitative morphological analysis of the cerebellar nuclei in normal and lurcher mutant mice. I. Morphology and cell number. *J Comp Neurol* **343**: 173-182, 1994.
- HECKROTH JA, GOLDOWITZ D, EISENMAN LM: Purkinje cell reduction in the reeler mutant mouse: a quantitative immunohistochemical study. *J Comp Neurol* **279**: 546-555, 1989.
- HERCULANO-HOUZEL S: Coordinated scaling of cortical and cerebellar numbers of neurons. *Front Neuroanat* **4**: 12, 2010.
- HERRUP K, MULLEN RJ: Regional variation and absence of large neurons in the cerebellum of the staggerer mouse. *Brain Res* **172**: 1-12, 1979a.

- HERRUP K, MULLEN RJ: Staggerer chimeras: intrinsic nature of Purkinje cell defects and implications for normal cerebellar development. *Brain Res* **178**: 443-457, 1979b.
- HILBER P, CASTON J: Motor skills and motor learning in Lurcher mutant mice during aging. *Neuroscience* **102**: 615-623, 2001.
- HILBER P, LORIVEL T, DELARUE C, CASTON J: Stress and anxious-related behaviors in Lurcher mutant mice. *Brain Res* **1003**: 108-112, 2004.
- HUBENER J, WEBER JJ, RICHTER C, HONOLD L, WEISS A, MURAD F, BREUER P, WULLNER U, BELLSTEDT P, PAQUET-DURAND F, TAKANO J, SAIDO TC, RIESS O, NGUYEN HP: Calpain-mediated ataxin-3 cleavage in the molecular pathogenesis of spinocerebellar ataxia type 3 (SCA3). *Hum Mol Genet* **22**: 508-518, 2013.
- HUYNH DP, FIGUEROA K, HOANG N, PULST SM: Nuclear localization or inclusion body formation of ataxin-2 are not necessary for SCA2 pathogenesis in mouse or human. *Nat Genet* **26**: 44-50, 2000.
- ITO M: Neurophysiological aspects of the cerebellar motor control system. *Int J Neurol* **7**: 162-176, 1970.
- ITO M: Neural design of the cerebellar motor control system. *Brain Res* **40**: 81-84, 1972.
- ITO M: *The Cerebellum and Neural Control*. Raven Press, New York, 1984.
- ITO M: Cerebellar circuitry as a neuronal machine. *Prog Neurobiol* **78**: 272-303, 2006.
- JACQUELIN C, STRAZIELLE C, LALONDE R: Spontaneous alternation and spatial learning in Dab1<sup>scm</sup> (scrambler) mutant mice. *Brain Res Bull* **87**: 383-386, 2012.
- JACQUELIN C, LALONDE R, JANTZEN-OSSOLA C, STRAZIELLE C: Neurobehavioral performances and brain regional metabolism in Dab1 (scm) (scrambler) mutant mice. *Behav Brain Res* **252**: 92-100, 2013.
- KANDEL ER, SCHWARTZ JH, JESSEL TM: *Principles of Neural Science*. 4th ed. McGraw-Hill, New York, 2000.
- KANG WY, KIM SS, CHO SK, KIM S, SUH-KIM H, LEE YD: Migratory defect of mesencephalic dopaminergic neurons in developing reeler mice. *Anat Cell Biol* **43**: 241-251, 2010.
- KELP A, KOEPPEN AH, PETRASCH-PARWEZ E, CALAMINUS C, BAUER C, PORTAL E, YU-TAEGER L, PICHLER B, BAUER P, RIESS O, NGUYEN HP: A novel transgenic rat model for spinocerebellar ataxia type 17 recapitulates neuropathological changes and supplies *in vivo* imaging biomarkers. *J Neurosci* **33**: 9068-9081, 2013.
- KIM SG, UGURBIL K, STRICK PL: Activation of a cerebellar output nucleus during cognitive processing. *Science* **265**: 949-951, 1994.
- KIM N, XIAO R, CHOI H, JO H, KIM JH, UHM SJ, PARK C: Abnormal sperm development in *pcd(3J)*<sup>-/-</sup> mice: the importance of *Agtpbp1* in spermatogenesis. *Mol Cells* **31**: 39-48, 2011.
- KISHORE A, MEUNIER S, POPA T: Cerebellar influence on motor cortex plasticity: behavioral implications for Parkinson's disease. *Front Neurol* **5**: 68, 2014.
- KLEMENT IA, SKINNER PJ, KAYTOR MD, YI H, HERSCH SM, CLARK HB, ZOGHBI HY, ORR HT: Ataxin-1 nuclear localization and aggregation: role in polyglutamine-induced disease in SCA1 transgenic mice. *Cell* **95**: 41-53, 1998.
- KOEPPEN AH: The pathogenesis of spinocerebellar ataxia. *Cerebellum* **4**: 62-73, 2005.
- KONNO A, SHUVAEV AN, MIYAKE N, MIYAKE K, IIZUKA A, MATSUURA S, HUDA F, NAKAMURA K, YANAGI S, SHIMADA T, HIRAI H: Mutant ataxin-3 with an abnormally expanded polyglutamine chain disrupts dendritic development and metabotropic glutamate receptor signaling in mouse cerebellar Purkinje cells. *Cerebellum* **13**: 29-41, 2014.
- KŘÍŽKOVÁ A, VOŽEH F: Development of early motor learning and topical motor skills in a model of cerebellar degeneration. *Behav Brain Res* **150**: 65-72, 2004.
- KYUHO S, KATO N, GEMBA H: Emergence of endoplasmic reticulum stress and activated microglia in Purkinje cell degeneration mice. *Neurosci Lett* **396**: 91-96, 2006.
- LALONDE R: Pharmacotherapy in animals with hereditary cerebellar disease. In: *Recent Research Development in Neurochemistry. Vol. 1*. PANDALAI SG (ed), Research Singpost, Trivandrum, 1996, pp 55-88.
- LALONDE R, STRAZIELLE C: Motor coordination, exploration, and spatial learning in a natural mouse mutation (nervous) with Purkinje cell degeneration. *Behav Genet* **33**: 59-66, 2003.

- LALONDE R, THIFAUULT S: Absence of an association between motor coordination and spatial orientation in lurcher mutant mice. *Behav Genet* **24**: 497-501, 1994.
- LALONDE R, LAMARRE Y, SMITH AM: Does the mutant mouse lurcher have deficits in spatially oriented behaviours? *Brain Res* **455**: 24-30, 1988.
- LALONDE R, JOYAL CC, COTÉ C, BOTEZ MI: Simultaneous visual discrimination learning in lurcher mutant mice. *Brain Res* **618**: 19-22, 1993a.
- LALONDE R, JOYAL CC, GUASTAVINO JM, BOTEZ MI: Hole poking and motor coordination in lurcher mutant mice. *Physiol Behav* **54**: 41-44, 1993b.
- LALONDE R, BENSOUOLA AN, FILALI M: Rotorod sensorimotor learning in cerebellar mutant mice. *Neurosci Res* **22**: 423-426, 1995.
- LALONDE R, FILALI M, BENSOUOLA AN, LESTIENNE F: Sensorimotor learning in three cerebellar mutant mice. *Neurobiol Learn Mem* **65**: 113-120, 1996.
- LALONDE R, HAYZOUN K, SELIMI F, MARIANI J, STRAZIELLE C: Motor coordination in mice with hotfoot, Lurcher, and double mutations of the Grid2 gene encoding the delta-2 excitatory amino acid receptor. *Physiol Behav* **80**: 333-339, 2003.
- LALONDE R, HAYZOUN K, DERER M, MARIANI J, STRAZIELLE C: Neurobehavioral evaluation of Reln-rl-orl mutant mice and correlations with cytochrome oxidase activity. *Neurosci Res* **49**: 297-305, 2004.
- LALOUETTE A, GUENET JL, VRIZ S: Hotfoot mouse mutations affect the delta 2 glutamate receptor gene and are allelic to lurcher. *Genomics* **50**: 9-13, 1998.
- LANDIS SC: Ultrastructural changes in the mitochondria of cerebellar Purkinje cells of nervous mutant mice. *J Cell Biol* **57**: 782-797, 1973.
- LANDIS DM, SIDMAN RL: Electron microscopic analysis of postnatal histogenesis in the cerebellar cortex of staggerer mutant mice. *J Comp Neurol* **179**: 831-863, 1978.
- LASEK K, LENCER R, GASER C, HAGENAH J, WALTER U, WOLTERS A, KOCK N, STEINLECHNER S, NAGEL M, ZUHLKE C, NITSCHKE MF, BROCKMANN K, KLEIN C, ROLFS A, BINKOFSKI F: Morphological basis for the spectrum of clinical deficits in spinocerebellar ataxia 17 (SCA17). *Brain* **129**: 2341-2352, 2006.
- LAVAIL MM, BLANKS JC, MULLEN RJ: Retinal degeneration in the pcd cerebellar mutant mouse. I. Light microscopic and autoradiographic analysis. *J Comp Neurol* **212**: 217-230, 1982.
- LAVAIL MM, WHITE MP, GORRIN GM, YASUMURA D, PORRELLO KV, MULLEN RJ: Retinal degeneration in the nervous mutant mouse. I. Light microscopic cytopathology and changes in the interphotoreceptor matrix. *J Comp Neurol* **333**: 168-181, 1993.
- LE MAREC N, LALONDE R: Sensorimotor learning and retention during equilibrium tests in Purkinje cell degeneration mutant mice. *Brain Res* **768**: 310-316, 1997.
- LEE NS, JEONG YG: Pogo: a novel spontaneous ataxic mutant mouse. *Cerebellum* **8**: 155-162, 2009.
- LEE KY, KIM JS, KIM SH, PARK HS, JEONG YG, LEE NS, KIM DK: Altered Purkinje cell responses and calmodulin expression in the spontaneously ataxic mouse, Pogo. *Eur J Neurosci* **33**: 1493-1503, 2011.
- LEGGIO MG, TEDESCO AM, CHIRICOZZI FR, CLAUSI S, ORSINI A, MOLINARI M: Cognitive sequencing impairment in patients with focal or atrophic cerebellar damage. *Brain* **131**: 1332-1343, 2008.
- LIN X, ANTALFFY B, KANG D, ORR HT, ZOGHBI HY: Polyglutamine expansion down-regulates specific neuronal genes before pathological changes in SCA1. *Nat Neurosci* **3**: 157-163, 2000.
- LIU J, TANG TS, TU H, NELSON O, HERNDON E, HUYNH DP, PULST SM, BEZPROZVANNY I: Deranged calcium signaling and neurodegeneration in spinocerebellar ataxia type 2. *J Neurosci* **29**: 9148-9162, 2009.
- MANTO MU: The wide spectrum of spinocerebellar ataxias (SCAs). *Cerebellum* **4**: 2-6, 2005.
- MANTO M, LORIVEL T: Cognitive repercussions of hereditary cerebellar disorders. *Cortex* **47**: 81-100, 2011.
- MANTO M, MARMOLINO D: Animal models of human cerebellar ataxias: a cornerstone for the therapies of the twenty-first century. *Cerebellum* **8**: 137-154, 2009.
- MARIANI J, CREPEL F, MIKOSHIBA K, CHANGEUX JP, SOTELO C: Anatomical, physiological and biochemical studies of the cerebellum from Reeler mutant mouse. *Philos Trans R Soc Lond B Biol Sci* **281**: 1-28, 1977.

- MARICICH SM, SOHA J, TREKNER E, HERRUP K: Failed cell migration and death of purkinje cells and deep nuclear neurons in the weaver cerebellum. *J Neurosci* **17**: 3675-3683, 1997.
- MIDDLETON FA, STRICK PL: Cerebellar projections to the prefrontal cortex of the primate. *J Neurosci* **21**: 700-712, 2001.
- MIGHELI A, ATTANASIO A, LEE WH, BAYER SA, GHETTI B: Detection of apoptosis in weaver cerebellum by electron microscopic in situ end-labeling of fragmented DNA. *Neurosci Lett* **199**: 53-56, 1995.
- MIKOSHIBA K, KOHSAKA S, TAKAMATSU K, AOKI E, TSUKADA Y: Morphological and biochemical studies on the cerebral cortex from reeler mutant mice: development of cortical layers and metabolic mapping by the deoxyglucose method. *J Neurochem* **34**: 835-844, 1980.
- MULLEN RJ, LAVAIL M: Two types of retinal degeneration in cerebellar mutant mice. *Nature* **258**: 528-530, 1975.
- MULLEN RJ, EICHER EM, SIDMAN RL: Purkinje cell degeneration, a new neurological mutation in the mouse. *Proc Natl Acad Sci U S A* **73**: 208-212, 1976.
- NISHIYAMA J, YUZAKI M: Excitotoxicity and autophagy: lurcher may not be a model of "autophagic cell death". *Autophagy* **6**: 568-570, 2010.
- NORMAN DJ, FENG L, CHENG SS, GUBBAY J, CHAN E, HEINTZ N: The lurcher gene induces apoptotic death in cerebellar Purkinje cells. *Development* **121**: 1183-1193, 1995.
- O'GORMAN S, SIDMAN RL: Degeneration of thalamic neurons in "Purkinje cell degeneration" mutant mice. I. Distribution of neuron loss. *J Comp Neurol* **234**: 277-297, 1985.
- PALAY SL, CHAN-PALAY V: *Cerebellar Cortex: Cytology and Organization*. Springer-Verlag, Berlin Heidelberg New York, 1974.
- PATIL N, COX DR, BHAT D, FAHAM M, MYERS RM, PETERSON AS: A potassium channel mutation in weaver mice implicates membrane excitability in granule cell differentiation. *Nat Genet* **11**: 126-129, 1995.
- PATRYLO PR, BROWNING RA, CRANICK S: Reeler homozygous mice exhibit enhanced susceptibility to epileptiform activity. *Epilepsia* **47**: 257-266, 2006.
- PHILLIPS RJS: „Lurcher“, a new gene in linkage group XI of the house mouse. *J Genet* **57**: 35-42, 1960.
- PORAS-GARCIA E, CENDELÍN J, DOMINGUEZ-DEL-TORO E, VOŽEH F, DELGADO-GARCIA JM: Purkinje cell loss affects differentially the execution, acquisition and prepulse inhibition of skeletal and facial motor responses in Lurcher mice. *Eur J Neurosci* **21**: 979-988, 2005.
- PORAS-GARCIA E, SANCHEZ-CAMPUSANO R, MARTINEZ-VARGAS D, DOMINQUEZ-DEL-TORO E, CENDELÍN J, VOŽEH F, DELGADO-GARCIA JM: Behavioral characteristics, associative learning capabilities, and dynamic association mapping in an animal model of cerebellar degeneration. *J Neurophysiol* **104**: 346-365, 2010.
- PURKARTOVÁ Z, VOŽEH F: Cerebellar degeneration in Lurcher mice under confocal laser scanning microscope. *Microsc Res Tech* **76**: 545-551, 2013.
- REEBER SL, LOESCHEL CA, FRANKLIN A, SILLITOE RV: Establishment of topographic circuit zones in the cerebellum of scrambler mutant mice. *Front Neural Circuits* **7**: 122, 2013.
- RESIBOIS A, CUVELIER L, GOFFINET AM: Abnormalities in the cerebellum and brainstem in homozygous lurcher mice. *Neuroscience* **80**: 175-190, 1997.
- ROFFLER-TARLOV S, GRAYBIEL AM: Weaver mutation has differential effects on dopamine-containing innervation of the limbic and nonlimbic striatum. *Nature* **307**: 62-66, 1984.
- ROFFLER-TARLOV S, HERRUP K: Quantitative examination of the deep cerebellar nuclei in the staggerer mutant mouse. *Brain Res* **215**: 49-59, 1981.
- ROFFLER-TARLOV S, SIDMAN RL: Concentrations of glutamic acid in cerebellar cortex and deep nuclei of normal mice and Weaver, Staggerer and nervous mutants. *Brain Res* **142**: 269-283, 1978.
- ROKYTA R: J. E. Purkyně and his discoveries in the nervous system. (in Czech) *Živa*, Nr. 5, 227-228, 2011.
- SAEGUSA H, WAKAMORI M, MATSUDA Y, WANG J, MORI Y, ZONG S, TANABE T: Properties of human CaV2.1 channel with a spinocerebellar ataxia type 6 mutation expressed in Purkinje cells. *Mol Cell Neurosci* **34**: 261-270, 2007.



- SCHUTTER DJL: Human cerebellum in motivation and emotion. In: *Handbook of the Cerebellum and Cerebellar Disorders*. MANTO M, GRUOL DL, SCHMAHMANN JD, KOIBUCHI N, ROSSI F (eds), Springer Science+Business Media, New York, 2013, pp 1771-1782.
- SELIMI F, DOUGHTY M, DELHAYE-BOUCHAUD N, MARIANI J: Target-related and intrinsic neuronal death in Lurcher mutant mice are both mediated by caspase-3 activation. *J Neurosci* **20**: 992-1000, 2000.
- SHAKKOTTAI VG, DO CARMO COSTA M, DELL'ORCO JM, SANKARANARAYANAN A, WULFF H, PAULSON HL: Early changes in cerebellar physiology accompany motor dysfunction in the polyglutamine disease spinocerebellar ataxia type 3. *J Neurosci* **31**: 13002-13014, 2011.
- SHARMA P, MAZUMDAR B, CHATTERJEE P: Cerebellar hypermetabolism on <sup>18</sup>F-FDG PET/CT with normal MRI in a case of paraneoplastic cerebellar degeneration with negative antibodies. *Rev Esp Med Nucl Imagen Mol* **34**: 79-80, 2015.
- SCHMAHMANN JD: From movement to thought: anatomic substrates of the cerebellar contribution to cognitive processing. *Hum Brain Mapp* **4**: 174-198, 1996.
- SCHMAHMANN JD, PANDYA DN: The cerebrotocerebellar system. *Int Rev Neurobiol* **41**: 31-60, 1997.
- SCHMAHMANN JD, SHERMAN JC: Cerebellar cognitive affective syndrome. *Int Rev Neurobiol* **41**: 433-440, 1997.
- SCHMITT A, TURCK CW, PILZ PK, MALCHOW B, VON WILMSDORFF M, FALKAI P, MARTINS-DE-SOUZA D: Proteomic similarities between heterozygous reeler mice and schizophrenia. *Biol Psychiatry* **74**: e5-e10, 2013.
- SILVERI MC, LEGGIO MG, MOLINARI M: The cerebellum contributes to linguistic production. A case of agrammatic speech following a right cerebellar lesion. *Neurology* **44**: 2047-2050, 1994.
- STEVANIN G, DURR A, DAVID G, DIDIERJEAN O, CANCEL G, RIVAUD S, TOURBAH A, WARTER JM, AGID Y, BRICE A: Clinical and molecular features of spinocerebellar ataxia type 6. *Neurology* **49**: 1243-1246, 1997.
- SULTAN F, KÖNIG T, MÖCK M, THIER P: Quantitative organization of neurotransmitters in the deep cerebellar nuclei of the Lurcher mutant. *J Comp Neurol* **452**: 311-323, 2002.
- SWEET HO, BRONSON RT, JOHNSON KR, COOK SA, DAVISSON MT: Scrambler, a new neurological mutation of the mouse with abnormalities of neuronal migration. *Mamm Genome* **7**: 798-802, 1996.
- SZENTAGOTHAI J: Structure-functional considerations of the cerebellar neuron network. *Proc IEEE* **56**: 960-968, 1968.
- TARONI F, CHIAPPARINI L, MARIOTTI C: Autosomal dominant spinocerebellar ataxias and episodic ataxias. In: *Handbook of the Cerebellum and Cerebellar Disorders, 1st ed.* MANTO M, GRUOL DL, SCHMAHMANN J, KOIBUCHI N, ROSSI F (eds). Springer Science+Business Media, New York, 2013, pp 2193-2267.
- TERASHIMA T, INOUE K, INOUE Y, MIKOSHIBA K, TSUKADA Y: Distribution and morphology of corticospinal tract neurons in reeler mouse cortex by the retrograde HRP method. *J Comp Neurol* **218**: 314-326, 1983.
- TERASHIMA T, INOUE K, INOUE Y, MIKOSHIBA K, TSUKADA Y: Observations on Golgi epithelial cells and granule cells in the cerebellum of the reeler mutant mouse. *Brain Res* **350**: 103-112, 1985.
- THULLIER F, LALONDE R, COUSIN X: Neurobehavioral evaluation of lurcher mutant mice during ontogeny. *Dev Brain Res* **100**: 22-28, 1997.
- TRIARHOU LC: Biological clues on neuronal degeneration based on theoretical fits of decay patterns: towards a mathematical neuropathology. *Folia Neuropathol* **48**: 3-10, 2010.
- TRIARHOU LC, NORTON J, GHETTI B: Anterograde transsynaptic degeneration in the deep cerebellar nuclei of Purkinje cell degeneration (pcd) mutant mice. *Exp Brain Res* **66**: 577-588, 1987.
- TŮMA J, CENDELÍN J, VOŽEH F: Maternal infanticide and low maternal ability in cerebellar mutants Lurcher. *Neuro Endocrinol Lett* **34**: 618-623, 2013.
- VALENTIN G: Über den Verlauf und die letzten Enden der Nerven. *Nova Acta Acad Leop Carol* **18**: 51-240, 1836.
- VAN MIER HI, TEMPEL LW, PERTMUTTER JS, RAICHLE ME, PETERSEN SE: Changes in brain activity during motor learning measured with PET: effects of hand of performance and practice. *J Neurophysiol* **80**: 2177-2199, 1998.

- VERBEEK DS, VAN DE WARRENBURG BP, WESSELING P, PEARSON PL, KREMER HP, SINKE RJ: Mapping of the SCA23 locus involved in autosomal dominant cerebellar ataxia to chromosome region 20p13-12.3. *Brain* **127**: 2551-2557, 2004.
- VOŽEH F, CADDY KWT, MYSLIVEČEK J, MOTÁŇOVÁ A: Some characteristics of early learning in cerebellar degeneration model. *Studia Psychol* **39**: 279-281, 1997.
- VOŽEH F, CENDELÍN J, MOTÁŇOVÁ A: The development of different types of learning in cerebellar degeneration model. *Homeostasis* **39**: 248-250, 1999.
- VOŽEH F, CENDELÍN J, ŠTENGLOVÁ V, BARCAL J, ZÁHLAVA J: The development of spatial learning in a model of olivocerebellar degeneration. *Homeostasis* **41**: 64-66, 2001.
- WANG QJ, DING Y, KOHTZ S, MIZUSHIMA N, CRISTEA IM, ROUT MP, CHAIT BT, ZHONG Y, HEINTZ N, YUE Z: Induction of autophagy in axonal dystrophy and degeneration. *J Neurosci* **26**: 8057-8068, 2006.
- WANG Z, GARDELL LR, OSSIPOV MH, VANDERAH TW, BRENNAN MB, HOCHGESCHWENDER U, HRUBY VJ, MALAN TP Jr, LAI J, PORRECA F: Pronociceptive actions of dynorphin maintain chronic neuropathic pain. *J Neurosci* **21**: 1779-1786, 2001.
- WATANABE H, MIZOGUCHI H, VERBEEK DS, KUZMIN A, NYBERG F, KRISHTAL O, SAKURADA S, BAKALKIN G: Non-opioid nociceptive activity of human dynorphin mutants that cause neurodegenerative disorder spinocerebellar ataxia type 23. *Peptides* **35**: 306-310, 2012.
- WATASE K, BARRETT CF, MIYAZAKI T, ISHIGURO T, ISHIKAWA K, HU Y, UNNO T, SUN Y, KASAI S, WATANABE M, GOMEZ CM, MIZUSAWA H, TSIEN RW, ZOGHBI HY: Spinocerebellar ataxia type 6 knockin mice develop a progressive neuronal dysfunction with age-dependent accumulation of mutant CaV2.1 channels. *Proc Natl Acad Sci U S A* **105**: 11987-11992, 2008.
- WON SJ, KIM SH, XIE L, WANG Y, MAO XO, JIN K, GREENBERG DA: Reelin-deficient mice show impaired neurogenesis and increased stroke size. *Exp Neurol* **198**: 250-259, 2006.
- WULLNER U, LOSCHMANN PA, WELLER M, KLOCKGETHER T: Apoptotic cell death in the cerebellum of mutant weaver and lurcher mice. *Neurosci Lett* **200**: 109-112, 1995.
- WYSS JM, STANFIELD BB, COWAN WM: Structural abnormalities in the olfactory bulb of the Reeler mouse. *Brain Res* **188**: 566-571, 1980.
- YAMADA M, SATO T, TSUJI S, TAKAHASHI H: CAG repeat disorder models and human neuropathology: similarities and differences. *Acta Neuropathol* **115**: 71-86, 2008.
- YOO SY, PENNESI ME, WEEBER EJ, XU B, ATKINSON R, CHEN S, ARMSTRONG DL, WU SM, SWEATT JD, ZOGHBI HY: SCA7 knockin mice model human SCA7 and reveal gradual accumulation of mutant ataxin-7 in neurons and abnormalities in short-term plasticity. *Neuron* **37**: 383-401, 2003.
- ZÁHLAVA J: The cerebellum. In: *Pathological Physiology of the Nervous System*. (in Czech) J. MYSLIVEČEK (ed), Charles University, Prague, 1994, pp 150-155.
- ZANJANI SH, SELIMI F, VOGEL MW, HAEBERLE AM, BOEUF J, MARIANI J, BAILLY YJ: Survival of interneurons and parallel fiber synapses in a cerebellar cortex deprived of Purkinje cells: studies in the double mutant mouse Grid2Lc/+;Bax-/- . *J Comp Neurol* **497**: 622-635, 2006.
- ZANJANI HS, LOHOF AM, McFARLAND R, VOGEL MW, MARIANI J: Enhanced survival of wild-type and Lurcher Purkinje cells in vitro following inhibition of conventional PKCs or stress-activated MAP kinase pathways. *Cerebellum* **12**: 377-389, 2013.
- ZHUCHENKO O, BAILEY J, BONNEN P, ASHIZAWA T, STOCKTON DW, AMOS C, DOBYNS WB, SUBRAMONY SH, ZOGHBI HY, LEE CC: Autosomal dominant cerebellar ataxia (SCA6) associated with small polyglutamine expansions in the alpha 1A-voltage-dependent calcium channel. *Nat Genet* **15**: 62-69, 1997.
- ZUO J, DE JAGER PL, TAKAHASHI KJ, JIANG W, LINDEN DJ, HEINTZ N: Neurodegeneration in Lurcher mice caused by mutation of  $\delta 2$  glutamate receptor gene. *Nature* **388**: 769-773, 1997.