

## REVIEW

## Research on Experimental Hypertension in Prague (1966-2009)

Jaroslav KUNEŠ<sup>1</sup>, Josef ZICHA<sup>1</sup><sup>1</sup>Laboratory of Experimental Hypertension, Institute of Physiology, Czech Academy of Sciences, Prague, Czech Republic

Received June 6, 2024

Accepted June 14, 2024

### Summary

The study of ontogenetic aspects of water and electrolyte metabolism performed in the Institute of Physiology (Czechoslovak Academy of Sciences) led to the research on the increased susceptibility of immature rats to salt-dependent forms of hypertension since 1966. Hemodynamic studies in developing rats paved the way to the evaluation of hemodynamic mechanisms during the development of genetic hypertension in SHR. A particular attention was focused on altered renal function and kidney damage in both salt and genetic hypertension with a special respect to renin-angiotensin system. Renal damage associated with hypertension progression was in the center of interest of several research groups in Prague. The alterations in ion transport, cell calcium handling and membrane structure as well as their relationship to abnormal lipid metabolism were studied in a close cooperation with laboratories in Munich, Glasgow, Montreal and Paris. The role of NO and oxidative stress in various forms of hypertension was a subject of a joint research with our Slovak colleagues focused mainly on NO-deficient hypertension elicited by chronic L-NAME administration. Finally, we adopted a method enabling us to evaluate the balance of vasoconstrictor and vasodilator mechanisms in BP maintenance. Using this method we demonstrated sympathetic hyperactivity and relative NO deficiency in rats with either salt-dependent or genetic hypertension. At the end of the first decennium of this century we were ready to modify our traditional approach towards modern trends in the research of experimental hypertension.

**Keywords:** Salt-dependent hypertension • Genetic hypertension • Body fluids • Hemodynamics • Ion transport • Cell membrane structure and function • Renal function • Renin-angiotensin systems

### Corresponding author

Josef Zicha, Institute of Physiology, Czech Academy of Sciences, Vídeňská 1083, 142 00 Prague 4, Czech Republic. E-mail: Josef.Zicha@fgu.cas.cz

### Our teachers


Physiological changes occurring during the ontogenesis belong to the historical research topics in the Institute of Physiology (Czechoslovak Academy of Sciences) (IPHYS). Water and electrolyte metabolism, body fluid distribution, renal function and vasopressin action were studied in developing rats since 1954 [1,2]. Dr. Jiří Jelínek focused his attention to the characteristic changes of water, sodium, potassium and chloride content in the body and body fluids of laboratory rats in particular developmental periods (suckling, weaning, prepuberty) [3,4]. Later he was inspired by the paper of Guillebeau and Skelton [5] who reported that immature rats are more susceptible to the induction of salt hypertension elicited by adrenal regeneration after its enucleation. In 1966 he published two papers indicating that not only adrenal-regeneration hypertension but also deoxycorticosterone (DOCA)-salt hypertension are more severe and self-sustaining in young than in adult animals [6-8]. Jelínek's research group paid a special attention to the study of changes in body fluids, renal function and structure with a special attention to renin-angiotensin system (RAS) [9-15]. They also performed fundamental studies on salt hypertension in monkeys (*Papio hamadrys*), indicating a greater blood pressure elevation (BP) in primates exposed to high salt intake from birth as compared to those exposed to this hypertensinogenic factor during sexual maturation. On the other hand, in adult salt hypertensive primates there was enhanced pulse pressure and reduced plasma volume as compared to control animals. They also demonstrated that blood pressure was inversely related to plasma volume in salt hypertensive monkeys [16,17].

In parallel, Prof. Jiří Křeček, who investigated the antidiuretic action of vasopressin in suckling and weanling rats [2,18,19], also started to study developmental aspects of salt hypertension using a peculiar model of uninephrectomized vasopressin-deficient Brattleboro rats drinking 0.6 % saline [20]. His experiments indicated that the age-dependent salt hypertension can be elicited even in the absence of vasopressin [21] and that prepuberty seems to be a critical period for the induction of more pronounced form of salt hypertension [22]. Subsequent hemodynamic studies, which used a dye dilution technique for the estimation of cardiac output in conscious animals [23], confirmed a significant BP elevation due to the increase in peripheral resistance and a major reduction of arterial compliance in young but not in adult animals with the above form of salt hypertension [24].

### Newly formed research group

In 1983 two young investigators from the above research groups – Dr. Jaroslav Kuneš and Dr. Josef Zicha – formed an independent research team and combined their methodical experience on body fluids and hemodynamic measurements in order to study further aspects of age-dependent salt hypertension. Using homozygous and heterozygous Brattleboro rats we demonstrated that sodium retention, which is not accompanied by sufficient water retention, did not induce

of DOCA-salt hypertension. Thus, antidiuretic rather than vasopressor effects of vasopressin are important for the development of DOCA-salt hypertension [25-27]. Another important topic of our study was the age-dependent role of digoxin-like factor (DLF) in BP maintenance in young and adult rats with DOCA-salt hypertension. Using the approach of Kojima *et al.* [28] based upon the acute administration of antidigoxin antibody, we found that endogenous DLF is important for BP maintenance especially in young DOCA-salt hypertensive rats [29,30], while this was not true for adult BP-matched rats with either DOCA-salt or spontaneous hypertension [31]. These findings were esteemed by Demuth Prize (Young Investigator Award of International Society of Hypertension, Interlaken 1984). Our subsequent review in *Hypertension* [32] suggested that prepuberty might be a critical period for enhanced salt-induced DLF production (Fig. 1). The third interesting field of our research was the role of sodium and chloride in the development of DOCA-salt hypertension. At that time there was an idea that chloride might be more important than sodium in the pathogenesis of salt-dependent forms of hypertension [32-36]. Our hemodynamic studies [37-39] revealed the delayed induction of hypertension and the absence of decreased arterial compliance in DOCA-treated rats, in which the dietary NaCl was replaced by NaHCO<sub>3</sub> to avoid the excess chloride intake.



**Fig. 1.** The inverse relationship of natriuretic response to plasma volume expansion [176] and blood pressure response to high salt intake during prepuberty and puberty [21,177]. Full horizontal bars indicate the onset and duration of high salt intake in the studies revealing a positive evidence for digoxin-like natriuretic factor [178-181], whereas negative studies are depicted as broken bars [182-185]. The density of vertical lines reflects the intensity of maturation processes. Modified from our review [32].

The research of genetic hypertension at the IPHYS in Prague was started by Dr. Ivan Albrecht who received first breeding pairs of spontaneously hypertensive rats (SHR) from Prof. Yukio Yamori in 1972. His measurements of cardiac output revealed that the prepuberty is a hyperkinetic phase of hypertension development in SHR [40,41]. Furthermore, on the basis of his original pharmacological interventions he proposed the prepuberty as a critical period for the development of genetic hypertension in this rat strain [42]. In 1981 Dr. Kuneš joined the laboratory of Dr. Pavel Hamet in Montreal and they started a long-term cooperation on the research of several important aspects of genetic hypertension. At the beginning they were interested in cardiac and renal hyperplasia of newborn SHR [43-46]. Ontogenetic aspects of hypertension development in the rat were later reviewed in *Physiological Reviews* in 1999 [47].

### **Jiří Heller and Prague hypertensive rats**

In the late 80ties Dr. Jiří Heller (Institute of Clinical and Experimental Medicine, Prague) developed a new model of genetic hypertension based upon Wistar rats – Prague hypertensive (PHR) and normotensive (PNR) rats [48]. This form of hypertension had many features similar to SHR but the main advantage of this model was that PHR and PNR had no histocompatibility problems so that kidneys could be transplanted between both lines without any signs of rejection [49,50]. The cross-transplantation of PNR kidney to bilaterally nephrectomized PHR animal lowered its blood pressure, while the transplantation of PHR kidney to PNR always induced hypertension development. This was true even if the transplanted kidney originated from PHR animals treated with antihypertensive drugs (captopril or nifedipine) since weaning [49]. A later study [50] indicated that the transplanted PHR kidney is important for the development but not for the maintenance of this Prague form of genetic hypertension. Using isolated kidneys from this model, Vaněčková *et al.* [51] reported the impairment of renal sodium excretion in adult PHR animals. Renal endothelin system was considered to be a possible candidate for a "hypertensinogenic" substance produced by PHR kidney [52].

Blood pressure and proteinuria were lowered in both PHR and PNR by chronic administration of AT<sub>1</sub> receptor inhibitor losartan, which also prevented later

development of renal damage in PHR animals [53]. One of the most fascinating findings in this rat strain was the demonstration of long-term blood pressure effects elicited by a brief treatment of young PHR with antihypertensive drugs inhibiting renin-angiotensin system. The beneficial BP effects were observed in 30-week-old PHR treated with losartan or perindopril at the age 5-9 weeks. Blood pressure reduction was even enhanced if this antihypertensive intervention was repeated at the age of 15-19 weeks. In addition, this long-term BP reduction was accompanied by a substantial antiproteinuric effect [54]. Blood pressure effects of early RAS blockade in PHR were considerably greater than those that we observed in similarly treated SHR [55].

### **Recombinant inbred strains as a research tool in the research of genetic hypertension**

A major stimulus for further research of the genetics in SHR was the establishment of recombinant inbred (RI) strains derived from SHR and normotensive BN.lx rats by Prof. Vladimír Křen and Dr. Michal Pravenec [56,57]. Blood pressure in this set of RI strains was initially determined by Dr. Kuneš using a direct carotid puncture [56]. Later he verified it by radiotelemetry measurement [58]. The initial papers started a very productive international cooperations (Theodore W. Kurtz, Tim J. Aitman, Pavel Hamet) leading to numerous highly cited papers [59-64] in prestigious journals including *Nature Genetics* [64-70]. A part of the results on the role of RI strains in the progress of genetics in SHR was summarized by Pravenec *et al.* [71,72]. However, the history of this extraordinary successful international research would deserve a separate review.

### **Red cell ion transport in experimental hypertension**

At the same time Dr. Zicha received the breeding pairs of inbred salt-sensitive (SS/Jr) and salt-resistant (SR/Jr) Dahl rats [73] from Prof. John P. Rapp (Toledo, OH). In these rats we studied red cell ion transport, effect of dietary calcium on blood pressure, alterations in arterial baroreflex function, changes in particular compartments of body fluids, adrenergic vascular innervation etc. [74-78]. In 1987-1988 Dr. Zicha studied the kinetics of ouabain-sensitive Na<sup>+</sup> and K<sup>+</sup> transport in erythrocytes of young and old Dahl rats in

the laboratory of Prof. Jochen Duhm in Munich. The data obtained in Munich [79] confirmed our earlier study done in Prague [74], which disclosed the enhanced ouabain-sensitive ion transport in red blood cells but a reduced activity of  $\text{Na}^+\text{-K}^+\text{-ATPase}$  in young salt hypertensive SS/Jr rats. The explanation of these experimental findings was simple but shocking. Salt hypertension in Dahl rats was associated with two major kinetic abnormalities of  $\text{Na}^+\text{-K}^+$  pump – the combination of increased affinity for internal sodium and decreased maximal transport rate. The former alteration is responsible for the enhanced ouabain-sensitive  $\text{Na}^+$  and  $\text{K}^+$  transport (studied under low physiological concentrations of internal sodium), while the latter kinetic change caused the decreased  $\text{Na}^+\text{-K}^+\text{-ATPase}$  activity (determined under saturating  $\text{Na}^+$  concentrations) [74,79]. Our further studies on membrane ion transport in Dahl rats indicated that the altered function of  $\text{Na}^+\text{-K}^+$  pump was related to abnormalities in cholesterol metabolism rather than to a mutation in *Atp1a1* gene [80-82].

In the early 90's Dr. Hasan Karama Bin Talib from Yemen made his PhD Thesis in our lab. He was trained in the study of red cell ion transport at Prof. Alan R. Chipperfield (Dundee, UK). Thanks to his dedication to experimental work we were able to extend our ion transport studies to further hypertensive models such as DOCA-salt treated rats [83,84], hereditary hypertriglyceridemic rats [85] or SHR and RI strains [86]. Our papers indicated that enhanced inward sodium leak and/or augmented  $\text{Na}^+\text{-K}^+\text{-cotransport}$  are responsible for

the increased red cell  $\text{Na}^+$  content, which is compensated by the acceleration of ouabain-sensitive  $\text{Na}^+$  extrusion by the  $\text{Na}^+\text{-K}^+$  pump. This was in line with our cooperative study with Dr. Sergei N. Orlov [87] which revealed a higher  $\text{Na}^+\text{-K}^+\text{-ATPase}$  activity and enhanced passive  $\text{K}^+$  permeability in erythrocytes of SHR compared to WKY or BN.lx rats. The findings on the red cell  $\text{Na}^+$  and  $\text{K}^+$  transport in experimental hypertension were later summarized in our reviews [88,89].


### New cooperations, new possibilities, new challenges

After the Velvet revolution the close cooperation of Dr. Zicha and Dr. Kuneš continued not only in traditional but also in newly added fields. They became to be the Editors of the journal *Physiological Research* (formerly *Physiologia bohemoslovaca*) published by IPHYS since 1954. Dr. Kuneš returned from his second stay in Montreal where the attention was focused on numerous genetic determinants of high BP in SHR - HSP70 [90-92], HSP27 [93], stress genes [94], major histocompatibility complex [95] etc. In 1993 Dr. Kuneš accepted a surprising idea of Dr. Hamet to apply for the organization of the congress of the International Society of Hypertension in Prague 2002. Thus, we spent further 10 years not only with scientific research but also with the preparation of this major event which brought to Prague almost 8000 participants of the first Joint ISH/ESH Meeting (Fig. 2).



J Zicha, K Arakawa, P Dominiak, J Kunes

**Fig. 2.** Dr. Kuneš and Dr. Zicha with the President of ISH Prof. Kikuo Arakawa.



**Fig. 3.** The relationship of plasma triglycerides to membrane microviscosity (left panels) and cell calcium handling (right panels) in platelets of Wistar rats (green symbols) and HTG rats (red symbols). The data were published in our papers [110, 111].

A new model of Prague hereditary hypertriglyceridemic (HTG) rats was developed by Vrána and Kazdová [96] and a moderate hypertension was disclosed in these animals by our lab [97]. HTG rats were utilized to evaluate the relationship of abnormalities in BP, lipid and glucose metabolism or ion transport [85,97,98] with particular genetic determinants [99-101]. Further studies evaluated the role of *Cd36* and *Igf2* genes in this model of hypertension [102,103]. The findings obtained in HTG rats were summarized in two review papers [104,105].

In 1991 we established a long-term cooperation with Dr. Marie-Aude Devynck (Hospital Necker, Paris) on the analysis of alterations in membrane structure and function, cell calcium handling, intracellular pH regulation and platelet aggregation in several hypertensive models such as Sabra, Dahl, Lyon and HTG rats. Our first joint study, which was focused on the effects of sodium on membrane fluidity in platelets of Wistar rats, indicated that the changes of intracellular rather than extracellular sodium are responsible for the alterations of platelet membrane microviscosity in the membrane outer leaflet (TMA-DPH anisotropy) but not in the membrane lipid core (DPH anisotropy). Sodium depletion increased TMA-DPH anisotropy and sodium repletion lowered it [106]. The next study in Lyon hypertensive (LH) rats demonstrated a positive correlation of TMA-DPH anisotropy with the intracellular calcium ( $Ca^{2+}_i$ ) in both platelets and

erythrocytes, while DPH anisotropy correlated negatively with blood pressure and  $Ca^{2+}_i$  [107]. DPH anisotropy but not TMA-DPH anisotropy was reduced in erythrocytes of Sabra and Dahl rats prone to develop salt hypertension in which DPH anisotropy also correlated negatively with blood pressure [108]. Platelets of LH rats were characterized by substantially elevated basal  $Ca^{2+}_i$  values, higher  $Ca^{2+}_i$  levels after thrombin stimulation, and enhanced initial rate of thrombin-induced  $Mn^{2+}$  entry through the receptor-operated  $Ca^{2+}$  channels. Plasma triglycerides but not cholesterol seemed to be related to platelet calcium handling [109].

Platelet or erythrocyte  $Ca^{2+}_i$  values were similar in HTG and Wistar rats, the same was true for  $Ca^{2+}$  influx into erythrocytes. On the other hand,  $Ca^{2+}_i$  response to thrombin stimulation and  $Mn^{2+}$  entry through the receptor-operated  $Ca^{2+}$  channels were reduced in platelets of HTG rats [110]. Plasma triglycerides correlated positively with platelet TMA-DPH anisotropy and negatively with DPH anisotropy. These relationships were present in both HTG and Wistar rats but the slopes of these relationships were considerably smaller in HTG than in Wistar rats (Fig. 3). In addition, platelet  $Ca^{2+}_i$  correlated positively with TMA-DPH anisotropy and negatively with DPH anisotropy, but the slopes of these relationships were almost identical in both rat strains [111,112]. HTG rats were also characterized by platelet hypoaggregability. The initial rate of platelet aggregation

was dependent on plasma triglycerides and the slope of this relationship was smaller in HTG than in normotensive control rats [113]. Furthermore, we tried to increase circulating triglycerides by drinking of fructose solution or to lower them by gemfibrozil treatment. Chronic reduction of plasma triglycerides was associated with increased DPH anisotropy, while chronic increase of plasma triglycerides was accompanied by decreased DPH anisotropy [114].

Our experiments also revealed different platelet calcium handling in rats with salt-dependent hypertension (Sabra and Dahl rats) and those with genetic hypertension (Lyon rats). In the former rat strains there was a highly significant correlation of platelet  $Ca^{2+}_i$  with pulse pressure but not with diastolic blood pressure, whereas platelet  $Ca^{2+}_i$  in Lyon rats correlated with diastolic blood pressure but not with pulse pressure [115,116]. Finally, we also paid attention to the cytosolic pH ( $pH_i$ ) and  $Ca^{2+}_i$  in platelets of Dahl and Sabra rats susceptible to salt hypertension development. Although there were no strain or salt-dependent differences in platelet  $Ca^{2+}_i$ , both strains had lower cytosolic  $pH_i$  [116,117]. Basal platelet  $pH_i$  of Dahl rats correlated positively with plasma triglycerides and plasma cholesterol and the changes in microviscosity of the outer membrane leaflet might be involved in  $pH_i$  regulation [117,118]. The above findings were summarized in a review on the abnormalities of membrane function and lipid metabolism in hypertension, which was published in *American Journal of Hypertension* in 1999 [112].

## Center for Cardiovascular Research

In 1999 the *Center for Cardiovascular Research* was established to combine the effort and expertise of leading Czech cardiovascular investigators. On this platform we have met an excellent newly formed research team of Dr. Luděk Červenka (Institute of Clinical and Experimental Medicine, Prague). He started his study in experimental nephrology and hypertension under the supervision of Dr. Jiří Heller [119,120]. In 1998 Dr. Červenka joined the laboratory of Dr. L. Gabriel Navar in New Orleans. There he studied the participation of renin-angiotensin system in the pathogenesis of two-kidney, one-clip Goldblatt hypertension with a special focus on renal functions [121-124]. He also paid the attention to the salt-sensitive hypertensive mice, in which bradykinin  $B_2$  receptor was inactivated, showing that high-salt diet and angiotensin II infusion induces the increase in blood pressure [125,126]. Moreover, genetic

inactivation of  $B_2$  receptor led to the worsening of 2K-1C Goldblatt hypertension [127] (Červenka 2003). After the return to Prague, Dr. Červenka succeeded to establish a breeding colony of Ren-2 transgenic rats (harboring mouse renin gene), the breeding pairs being provided by Prof. Detlev Ganten and Dr. Michael Bader (Berlin). Thanks to a long-term fruitful cooperation with Prof. Herbert J. Kramer (Bonn) the experimental research performed in Ren-2 transgenic rats covered not only the contribution of particular RAS components such as angiotensin II receptor subtype  $AT_{1A}$  [128,129] or angiotensin 1-7 [130,131] but also the role of neuronal NO synthase [132,133] and oxidative stress [134-136] as well as the influence of salt intake [137,138] or anesthesia [139]. Importantly, they demonstrated substantial differences in plasma and renal angiotensin II concentrations depending on whether the animals were anesthetized or conscious. A considerable attention was paid to endothelin system. They demonstrated that non-selective ET receptor blockade reduced proteinuria and attenuated cardiac hypertrophy in homozygous TGR [140]. Later, they focused on selective  $ET_A$  receptor blockade both in heterozygous and homozygous TGR [141-143]. Another important topic of his research were CYP-450-dependent oxygenase products – epoxyeicosatrienoic acids (EETs) and 20-hydroxyeicosatetraenoic acid (20-HETE). They found higher urinary 20-HETE and lower EETs excretion in TGR as compared with normotensive HanSD, suggesting that the imbalance between pro-hypertensive and anti-hypertensive CYP-450 products contribute to hypertension in TGR [144]. The inhibition of 20-HETE formation and EETs degradation led to BP decrease [145,146]. Dr. Červenka also acquired CYP1a1-Ren-2 transgenic rats in which hypertension is inducible by xenobiotic indole-3-carbinole. Severe hypertension develops already two days following its administration and is accompanied by substantial body weight loss and cardiac hypertrophy [147]. Moreover, impaired renal autoregulation precedes the development of hypertension in this experimental model [148].

Since hypertension is largely caused by the enhanced constriction of small resistance arteries together with the attenuation of vasodilator mechanisms, we directed our research to both *in vitro* and *in vivo* abnormalities of vascular tone in various form of experimental hypertension. We also paid our attention to the accelerated growth of vascular smooth muscle cells (VSMC) isolated from SHR aorta and to the respective


sex differences in this strain. We demonstrated a shorter doubling time in VSMC from male SHR compared to those from female animals [149,150]. VSMC isolated from the aorta of male SHR proliferate more rapidly than those obtained from female SHR. Angiotensin II stimulation of VSMC growth was more pronounced in cells isolated from the aorta of male SHR compared to cells from female SHR [151]. Furthermore, we observed that the augmented  $[Ca^{2+}]_i$  response to angiotensin II in male compared to female aortic VSMC was dependent on  $Ca^{2+}$  influx [152].

### The balance of vasoconstrictor and vasodilator mechanisms

In 1987 we started to evaluate the contribution of various pressor systems to BP maintenance in DOCA-salt hypertensive rats [153]. Simultaneously, we were inspired by the papers of Dr. Haralambos Gavras and Dr. Bernard Waeber [154-157] who performed the acute blockade of particular vasoactive systems in conscious rats. Therefore, we adapted the experimental protocol described by Minami *et al.* [158] for the estimation of the contribution of three principal vasoactive systems (renin-angiotensin, sympathetic and nitric oxide) to BP maintenance. A sequential blockade of these three systems in conscious cannulated animals was used for this purpose. This gave us the opportunity to study the role of sympathetic nervous system (SNS) which seemed to be enhanced in various salt-dependent forms of hypertension [76,159,160]. Our first studies, which were performed in Dahl and HTG rats, indicated the sympathetic hyperactivity and relative NO deficiency in both forms of hypertension [161,162]. A comparison of BP response to the acute administration of tempol (superoxide dismutase mimetic) in young and adult Dahl rats suggested a greater involvement of reactive oxygen species in young salt hypertensive animals [163].

At that time Dr. Kuneš became a director of IPHYS and we established a valuable cooperation with Dr. Olga Pecháňová from the Institute of Normal and Pathological Physiology (Slovak Academy of Sciences, Bratislava). She introduced to us a model of NO-deficient hypertension elicited by chronic administration of non-specific NO synthase inhibitor L-NAME [164-166] and we started to examine this model using our techniques. One of our first joint papers [167] revealed the importance of sympathetic hyperactivity in this form of hypertension. We also reported that a considerable part of vasodilation persisting in L-NAME-treated rats can be abolished by the acute administration of the inhibitors of inducible NO synthase. Our further studies indicated a similarity of L-NAME-induced hypertension in immature and adult rats [169] and a possibility to attenuate the development of this NO-deficient form of hypertension by chronic N-acetylcysteine (NAC) administration [170]. Furthermore, we demonstrated that chronic NAC treatment augmented NO-dependent vasodilatation, whereas chronic captopril treatment reduced sympathetic vasoconstriction in rats with L-NAME-induced hypertension [171]. The importance of sympathetic hyperactivity in this form of hypertension was confirmed in our further study indicating the attenuated development of L-NAME-induced hypertension in rats pretreated with pertussis toxin inactivating  $G_i$  protein [172].

Within the frame of this cooperation we also studied the effects of chronic NAC administration on the development of genetic hypertension in young SHR (preventive study) [173] and on the maintenance of hypertension in adult SHR with established hypertension (therapeutic study) [174]. Finally, Dr. Kuneš upgraded the original concept that hypertension is a result of the interaction between genetic and environmental factors [175] by considering the significant role of epigenetic inheritance [176] (Fig. 4).



**Fig. 4.** Epigenetic and gene interactions with environmental factors during blood pressure ontogeny and hypertension development. E<sub>1</sub> - E<sub>n</sub> represent environmental stimuli affecting expression of genetic information (G<sub>1</sub> - G<sub>n</sub>) occurring in particular critical periods (developmental windows). Modified from our review [47].

## Future perspectives

Another major profit from the collaboration with Dr. Pecháňová was the facilitation of the long-term work of several young Slovak colleagues in our lab – Drs Ludovít Paulis, Silvia Líšková, Mária Pintérová, Michal Behuliak and Michal Bencze. Together with the return of Dr. Ivana Vaněčková from the Institute of Clinical and Experimental Medicine (Prague) in 2010, this team was prepared for a further development of the Laboratory of Experimental Hypertension in IPHYS. The original direction of our research was modified towards new hypertensive models (Ren-2 transgenic rats) and new mechanisms in blood pressure control and hypertension development (role of endothelin, interaction of RAS and SNS, central and peripheral effects of angiotensin II, mechanisms of sympathetic hyperactivity, contribution of

Ca<sup>2+</sup> influx and Ca<sup>2+</sup> sensitization, wire myography for the examination of conduit and resistance arteries). Thus, we were ready for a new chapter in the history of the Laboratory of Experimental Hypertension in IPHYS.

## Conflict of Interest

There is no conflict of interest.

## Acknowledgements

This study was supported by institutional support from the Institute of Physiology, Czech Academy of Sciences, the grant Nr. RVO 67985823 and by the project National Institute for Research of Metabolic and Cardiovascular Diseases (Programme EXCELES, ID Project No. LX22NPO5104) – Funded by the European Union – Next Generation EU.

## References

- Čapek K, Jelínek J. The development of the control of water metabolism. I. The excretion of urine in young rats. *Physiol Bohemoslov* 1956;5:91-96.
- Křeček J, Křečková J. The development of the regulation of water metabolism. III. The relation between water and milk intake in infant rats. *Physiol Bohemoslov* 1957;6: 26-34.
- Jelínek J. The development of the regulation of water metabolism .5. Changes in content of water, potassium, sodium and chloride in body and body fluids of the rat during development. *Physiol Bohemoslov* 1961;10: 249-258.
- Jelínek J. The development of the regulation of water metabolism .6. Changes in the volume of cellular and extracellular fluid in the body of rats during development. *Physiol Bohemoslov* 1961;10: 259-266.
- Guillebeau J, Skelton FR. The influence of age on the development of adrenal-regeneration hypertension. *Endocrinology* 1956;59:201-212. <https://doi.org/10.1210/endo-59-2-201>
- Jelínek J, Albrecht I, Musilová H. The age factor in experimental hypertension: hypertension due to adrenal regeneration. *Physiol Bohemoslov* 1966;15:424-433.
- Jelínek J. Salt intake and sexual differences in sensitivity of rats to adrenal-regeneration hypertension. *Physiol Bohemoslov* 1967;16:389-399.
- Musilová H, Jelínek J, Albrecht I. The age of factor in experimental hypertension of the DCA type in rats. *Physiol Bohemoslov* 1966;15:525-531.
- Jelínek J. Blood volume in rats with DOCA-hypertension elicited at different ages. *Physiol Bohemoslov* 1972;21:581-588.
- Karen P, Deyl Z, Jelínek J. Influence of age and treatment with sodium chloride and desoxycorticosterone acetate on collagenous stroma of compensatory growing kidneys in the rat. *Mech Ageing Dev* 1974;3:157-163. [https://doi.org/10.1016/0047-6374\(74\)90012-8](https://doi.org/10.1016/0047-6374(74)90012-8)
- Kuneš J, Karen P, Čapek K, Jelínek J. The influence of salt intake on glomerular count in compensatory kidney hypertrophy in rats of different ages. *Experientia* 1976;32:887-888. <https://doi.org/10.1007/BF02003748>
- Pohlová I, Kuneš J, Jelínek J. The influence of age on the activity of the renin-angiotensin system in rats with adrenal-regeneration hypertension. *Physiol Bohemoslov* 1983;32:438-448.
- Kuneš J, Jelínek J. Influence of age on saline hypertension in subtotal nephrectomized rats. *Physiol Bohemoslov* 1984;33:123-128.
- Hausmann J, Jelínek J, Deyl Z. Changes in glomerular basement membrane collagen caused by DOCA-NaCl treatment in rats. *Nephron* 1985;39:250-254. <https://doi.org/10.1159/000183380>



15. Vaněčková I, Kuneš J, Jelínek J. Function of the isolated perfused kidneys from young or adult rats with post-DOCA-salt hypertension. *Clin Exp Hypertens* 1999;21:407-421. <https://doi.org/10.3109/10641969909068673>
16. Cherchovich GM, Čapek K, Jefremova Z, Pohlová I, Jelínek J. High salt intake and blood pressure in lower primates (*Papio hamadryas*). *J Appl Physiol* 1976;40:601-604. <https://doi.org/10.1152/jappl.1976.40.4.601>
17. Kuneš J, Čapek K, Jelínek J, Cherkovich GM. Extracellular fluid distribution in salt-hypertensive monkeys. *Experientia* 1978;34:753-754. <https://doi.org/10.1007/BF01947304>
18. Dlouhá H, Křeček J, Krečková J. Water diuresis and the effect of vasopressin in infant rats. *Physiol Bohemoslov* 1963;12:443-452.
19. Dlouhá H, Kraus M, Křeček J, Pliška V. Sensitivity of rats to vasopressin in the weaning period. *Physiol Bohemoslov* 1965;14:217-224.
20. Hall CE, Ayachi S, Hall O. Spontaneous hypertension in rats with hereditary hypothalamic diabetes insipidus (Brattleboro strain). *Tex Rep Biol Med* 1973;31:471-487.
21. Dlouhá H, Křeček J, Zicha J. Hypertension in rats with hereditary diabetes insipidus. The role of age. *Pflugers Arch*. 1977;369:177-182. <https://doi.org/10.1007/BF00591574>
22. Dlouhá H, Křeček J, Zicha J. Effect of age on hypertensive stimuli and the development of hypertension in Brattleboro rats. *Clin Sci (Lond)* 1979;57:273-275. <https://doi.org/10.1042/cs0570273>
23. Albrecht I, Hallbäck M, Julius S, Lundgren Y, Stage L, Weiss L, Folkow B. Arterial pressure, cardiac output and systemic resistance before and after pithing in normotensive and spontaneously hypertensive rats. *Acta Physiol Scand*. 1975;94:378-385. <https://doi.org/10.1111/j.1748-1716.1975.tb05897.x>
24. Zicha J, Křeček J, Dlouhá H. Age-dependent salt hypertension in Brattleboro rats: a hemodynamic analysis. *Ann N Y Acad Sci* 1982;394:330-342. <https://doi.org/10.1111/j.1749-6632.1982.tb37444.x>
25. Zicha J, Kuneš J, Jelínek J. Age-dependent DOCA-salt hypertension in Brattleboro rats: the role of vasopressin. *Physiol Bohemoslov* 1987;36:33-42.
26. Kuneš J, Nedvídek J, Zicha J. Vasopressin and water distribution in rats with DOCA-salt hypertension. *J Hypertens* 1989;7 (Suppl 6):S204-S205. <https://doi.org/10.1097/00004872-198900076-00098>
27. Zicha J, Kuneš J, Lébl M, Pohlová I, Slaninová J, Jelínek J. Antidiuretic and pressor actions of vasopressin in age-dependent DOCA-salt hypertension. *Am J Physiol* 1989;256:R138-R145. <https://doi.org/10.1152/ajpregu.1989.256.1.R138>
28. Kojima I, Yoshihara S, Ogata E. Involvement of endogenous digitalis-like substance in genesis of deoxycorticosterone-salt hypertension. *Life Sci* 1982 May 24;30:1775-1781. [https://doi.org/10.1016/0024-3205\(82\)90313-7](https://doi.org/10.1016/0024-3205(82)90313-7)
29. Zicha J, Štolba P, Pohlová I, Kuneš J, Jelínek J. A different role of digoxin-like factor in the maintenance of elevated blood pressure in rats treated with DOCA and saline in youth or only in adulthood. *J Hypertens Suppl*. 1984;2 (Suppl 3):S481-S483.
30. Zicha J, Kuneš J, Štolba P. Endogenous digoxin-like factor contributes to the elevation of systemic resistance in rats exposed to high salt intake from prepuberty. *J Hypertens* 1985;3 (Suppl 3):S17-S19.
31. Kuneš J, Štolba P, Pohlová I, Jelínek J, Zicha J. The importance of endogenous digoxin-like factors in rats with various forms of experimental hypertension. *Clin Exp Hypertens A* 1985;7(5-6):707-720. <https://doi.org/10.3109/10641968509077222>
32. Zicha J, Kuneš J, Jelínek J. Experimental hypertension in young and adult animals. *Hypertension* 1986;8:1096-1104. <https://doi.org/10.1161/01.HYP.8.12.1096>
33. Kurtz TW, Morris RC Jr. Dietary chloride as a determinant of "sodium-dependent" hypertension. *Science* 1983;222:1139-1141. <https://doi.org/10.1126/science.6648527>
34. Kurtz TW, Morris RC Jr. Hypertension in the recently weaned Dahl salt-sensitive rat despite a diet deficient in sodium chloride. *Science* 1985;230:808-810. <https://doi.org/10.1126/science.4059913>
35. Whitescarver SA, Ott CE, Jackson BA, Guthrie GP Jr, Kotchen TA. Salt-sensitive hypertension: contribution of chloride. *Science* 1984;223:1430-1432. <https://doi.org/10.1126/science.6322303>
36. Whitescarver SA, Holtzclaw BJ, Downs JH, Ott CE, Sowers JR, Kotchen TA. Effect of dietary chloride on salt-sensitive and renin-dependent hypertension. *Hypertension* 1986;8:56-61. <https://doi.org/10.1161/01.HYP.8.6.552>, <https://doi.org/10.1161/01.HYP.8.1.56>

37. Zicha J, Kuneš J, Jelínek J, Leont'eva GR, Govyrin VA. The importance of sodium and chloride ions for the development of DOCA-NaCl hypertension: a haemodynamic study. *Physiol Bohemoslov* 1986;35:308-312.
38. Zicha J, Kuneš J. Haemodynamic changes induced by short- and long-term sodium chloride or sodium bicarbonate intake in deoxycorticosterone-treated rats. *Acta Physiol Scand* 1994;151:217-223. <https://doi.org/10.1111/j.1748-1716.1994.tb09740.x>
39. Kuneš J, Zicha J, Jelínek J. The role of chloride in deoxycorticosterone hypertension: selective sodium loading by diet or drinking fluid. *Physiol Res* 2004;53:149-154. <https://doi.org/10.33549/physiolres.930000.53.149>
40. Albrecht I. The hemodynamics of early stages of spontaneous hypertension in rats. Part I: Male study. *Jpn Circ J* 1974;38:985-990. <https://doi.org/10.1253/jcj.38.985>
41. Albrecht I. The hemodynamics of early stages of spontaneous hypertension in rats. Part II: Female study. *Jpn Circ J* 1974;38:991-996. <https://doi.org/10.1253/jcj.38.991>
42. Albrecht I. Critical period for the development of spontaneous hypertension in rats. *Mech Ageing Dev* 1974;3:75-79. [https://doi.org/10.1016/0047-6374\(74\)90007-4](https://doi.org/10.1016/0047-6374(74)90007-4)
43. Pang SC, Long C, Poirier M, Tremblay J, Kuneš J, Vincent M, Sassard J, Duzzi L, Bianchi G, Ledingham J, et al. Cardiac and renal hyperplasia in newborn genetically hypertensive rats. *J Hypertens* 1986;4 (Suppl 3):S119-S122.
44. Kuneš J, Pang SC, Cantin M, Genest J, Hamet P. Cardiac and renal hyperplasia in newborn spontaneously hypertensive rats. *Clin Sci (Lond)* 1987;72:271-275. <https://doi.org/10.1042/cs0720271>
45. Dobešová Z, Zicha J, Kuneš J. Newborn organ weight and spontaneous hypertension: recombinant inbred strain study. *Clin Exp Hypertens* 1997;19:403-415. <https://doi.org/10.3109/10641969709084504>
46. Hamet P, Pausova Z, Dumas P, Sun YL, Tremblay J, Pravenec M, Kuneš J, Křenová D, Křen V. Newborn and adult recombinant inbred strains: a tool to search for genetic determinants of target organ damage in hypertension. *Kidney Int* 1998;53:1488-1492. <https://doi.org/10.1046/j.1523-1755.1998.00938.x>
47. Zicha J, Kuneš J. Ontogenetic aspects of hypertension development: analysis in the rat. *Physiol Rev* 1999;79:1227-1282. <https://doi.org/10.1152/physrev.1999.79.4.1227>
48. Heller J, Hellerová S, Dobešová Z, Kuneš J, Zicha J. The Prague Hypertensive Rat: a new model of genetic hypertension. *Clin Exp Hypertens* 1993;15:807-818. <https://doi.org/10.3109/10641969309041643>
49. Heller J, Schubert G, Havlíčková J, Thureau K. The role of the kidney in the development of hypertension: a transplantation study in the Prague hypertensive rat. *Pflugers Arch* 1993;425:208-212. <https://doi.org/10.1007/BF00374168>
50. Heller J, Havlíčková J, Hellerová S, Schubert G, Thureau K. A transient role of the kidney in the maintenance of hypertension. *Kidney Int* 1996; Suppl 55:S163-S165.
51. Vaněčková I, Heller J, Thureau K. Possible contribution of impaired sodium excretion to the development and maintenance of hypertension: a study of the isolated kidneys of the Prague hypertensive rat. *Pflugers Arch* 1997;434:587-591. <https://doi.org/10.1007/s004240050440>
52. Vogel V, Bäcker A, Heller J, Kramer HJ. The renal endothelin system in the Prague hypertensive rat, a new model of spontaneous hypertension. *Clin Sci (Lond)* 1999;97:919-918. <https://doi.org/10.1042/cs0970091>
53. Heller J, Kramer HJ, Červenka L, Hellerová S. Losartan protects the rat kidney from ischemic injury. *Kidney Int* 1996; Suppl 55:S113-S114.
54. Heller J, Hellerová S. Long-term effect on blood pressure of early brief treatment by different antihypertensive agents: a study in the Prague hypertensive rat. *Kidney Blood Press Res* 1998;21:445-451. <https://doi.org/10.1159/000025898>
55. Zicha J, Dobešová Z, Kuneš J. Late blood pressure reduction in SHR subjected to transient captopril treatment in youth: possible mechanisms. *Physiol Res* 2008;57:495-498. <https://doi.org/10.33549/physiolres.931615>
56. Pravenec M, Klír P, Křen V, Zicha J, Kuneš J. An analysis of spontaneous hypertension in spontaneously hypertensive rats by means of new recombinant inbred strains. *J Hypertens* 1989;7:217-221. <https://doi.org/10.1097/00004872-198903000-00008>
57. Kuneš J, Křen V, Klír P, Zicha J, Pravenec M. Genetic determination of heart and kidney weights studied using a set of recombinant inbred strains: the relationship to blood pressure. *J Hypertens* 1990;8:1091-1095. <https://doi.org/10.1097/00004872-199012000-00004>

58. Kuneš J, Dobešová Z, Musilová A, Zídek V, Vorlíček J, Pravenec M, Křen V, Zicha J. Hemodynamic characterization of recombinant inbred strains: twenty years later. *Hypertens Res* 2008;31:1659-1668. <https://doi.org/10.1291/hypres.31.1659>
59. Pravenec M, Křen V, Kuneš J, Scicli AG, Carretero OA, Simonet L, Kurtz TW. Cosegregation of blood pressure with a kallikrein gene family polymorphism. *Hypertension* 1991;17:242-246. <https://doi.org/10.1161/01.HYP.17.2.242>
60. Pravenec M, Simonet L, Křen V, Kuneš J, Levan G, Szpirer J, Szpirer C, Kurtz T. The rat renin gene: assignment to chromosome 13 and linkage to the regulation of blood pressure. *Genomics* 1991;9:466-472. [https://doi.org/10.1016/0888-7543\(91\)90412-8](https://doi.org/10.1016/0888-7543(91)90412-8)
61. Pravenec M, Gauguier D, Schott JJ, Buard J, Křen V, Bílá V, Szpirer C, Szpirer J, Wang JM, Huang H, et al. Mapping of quantitative trait loci for blood pressure and cardiac mass in the rat by genome scanning of recombinant inbred strains. *J Clin Invest* 1995;96:1973-1978. <https://doi.org/10.1172/JCI118244>
62. Bottger A, van Lith HA, Křen V, Křenová D, Bílá V, Vorlíček J, Zídek V, Musilová A, Zdobinská M, Wang JM, van Zutphen BF, Kurtz TW, Pravenec M. Quantitative trait loci influencing cholesterol and phospholipid phenotypes map to chromosomes that contain genes regulating blood pressure in the spontaneously hypertensive rat. *J Clin Invest* 1996;98:856-862. <https://doi.org/10.1172/JCI118858>
63. Pravenec M, Gauguier D, Schott JJ, Buard J, Křen V, Bílá V, Szpirer C, Szpirer J, Wang JM, Huang H, St Lezin E, Spence MA, Flodman P, Printz M, Lathrop GM, Vergnaud G, Kurtz TW. A genetic linkage map of the rat derived from recombinant inbred strains. *Mamm Genome* 1996;7:117-127. <https://doi.org/10.1007/s003359900031>
64. Křen V, Pravenec M, Lu S, Křenová D, Wang JM, Wang N, Merriouns T, Wong A, St Lezin E, Lau D, Szpirer C, Szpirer J, Kurtz TW. Genetic isolation of a region of chromosome 8 that exerts major effects on blood pressure and cardiac mass in the spontaneously hypertensive rat. *J Clin Invest* 1997;99:577-581. <https://doi.org/10.1172/JCI119198>
65. Aitman TJ, Gotoda T, Evans AL, Imrie H, Heath KE, Trembling PM, Truman H, Wallace CA, Rahman A, Doré C, Flint J, Křen V, Zídek V, Kurtz TW, Pravenec M, Scott J. Quantitative trait loci for cellular defects in glucose and fatty acid metabolism in hypertensive rats. *Nat Genet* 1997;16:197-201. <https://doi.org/10.1038/ng0697-197>
66. Aitman TJ, Glazier AM, Wallace CA, Cooper LD, Norsworthy PJ, Wahid FN, Al-Majali KM, Trembling PM, Mann CJ, Shoulders CC, Graf D, St Lezin E, Kurtz TW, Křen V, Pravenec M, Ibrahimi A, Abumrad NA, Stanton LW, Scott J. Identification of Cd36 (Fat) as an insulin-resistance gene causing defective fatty acid and glucose metabolism in hypertensive rats. *Nat Genet* 1999;21:76-83. <https://doi.org/10.1038/5013>
67. Pravenec M, Landa V, Zídek V, Musilová A, Křen V, Kazdová L, Aitman TJ, Glazier AM, Ibrahimi A, Abumrad NA, Qi N, Wang JM, St Lezin EM, Kurtz TW. Transgenic rescue of defective Cd36 ameliorates insulin resistance in spontaneously hypertensive rats. *Nat Genet* 2001;27:156-158. <https://doi.org/10.1038/84777>
68. Hübner N, Wallace CA, Zimdahl H, Petretto E, Schulz H, Maciver F, Mueller M, Hummel O, Monti J, Zídek V, Musilová A, Křen V, Causton H, Game L, Born G, Schmidt S, Müller A, Cook SA, Kurtz TW, Whittaker J,
69. Pravenec M, Aitman TJ. Integrated transcriptional profiling and linkage analysis for identification of genes underlying disease. *Nat Genet* 2005;37:243-253. <https://doi.org/10.1038/ng1522>
70. Guryev V, Saar K, Adamovic T, Verheul M, van Heesch SA, Cook S, Pravenec M, Aitman T, Jacob H, Shull JD, Hubner N, Cuppen E. Distribution and functional impact of DNA copy number variation in the rat. *Nat Genet* 2008;40:538-545. <https://doi.org/10.1038/ng.141>
71. Pravenec M, Churchill PC, Churchill MC, Viklický O, Kazdová L, Aitman TJ, Petretto E, Hubner N, Wallace CA, Zimdahl H, Zídek V, Landa V, Dunbar J, Bidani A, Griffin K, Qi N, Maxová M, Křen V, Mlejnek P, Wang J, Kurtz TW. Identification of renal Cd36 as a determinant of blood pressure and risk for hypertension. *Nat Genet* 2008;40:952-954. <https://doi.org/10.1038/ng.164>
72. Pravenec M, Zídek V, Landa V, Šimáková M, Mlejnek P, Kazdová L, Bílá V, Křenová D, Křen V. Genetic analysis of "metabolic syndrome" in the spontaneously hypertensive rat. *Physiol Res* 2004;53 (Suppl 1):S15-S22. <https://doi.org/10.33549/physiolres.930000.53.S15>

73. Pravenec M, Křen V, Landa V, Mlejnek P, Musilová A, Šilhavý J, Šimáková M, Zidek V. Recent progress in the genetics of spontaneously hypertensive rats. *Physiol Res* 2014;63 (Suppl 1):S1-S8. <https://doi.org/10.33549/physiolres.932622>
74. Rapp JP, Dene H. Development and characteristics of inbred strains of Dahl salt-sensitive and salt-resistant rats. *Hypertension* 1985;7:340-349. [https://doi.org/10.1161/01.HYP.7.3.Pt\\_1.340](https://doi.org/10.1161/01.HYP.7.3.Pt_1.340), <https://doi.org/10.1161/01.HYP.7.3.340>
75. Zicha J, Byšková E, Kuneš J, Pohlová I, Jelínek J. Sodium pump activity in young and adult salt hypertensive Dahl rats. *Klin Wochenschr.* 1987;65 (Suppl 8):76-81.
76. Kuneš J, Zicha J, Hamet P. Effects of dietary calcium on the development of salt hypertension in young and adult Dahl rats. *J Hypertens* 1988;6 (Suppl 4):S225-S227. <https://doi.org/10.1097/00004872-198812040-00067>
77. Kuneš J, Leontjeva GR, Byšková E, Pohlová I, Govyrin VA, Zicha J. Adrenergic innervation of blood vessels in Dahl rats with salt hypertension. *Clin Exp Hypertens A* 1991;13:1343-1355. <https://doi.org/10.3109/10641969109048797>
78. Dobešová Z, Kuneš J, Zicha J. Body fluid alterations and organ hypertrophy in age-dependent salt hypertension of Dahl rats. *Physiol Res* 1995;44:377-387.
79. Nedvídek J, Zicha J. Baroreflex control of heart rate in young and adult salt hypertensive inbred Dahl rats. *Physiol Res* 2000;49:323-330.
80. Zicha J, Duhm J. Kinetics of Na<sup>+</sup> and K<sup>+</sup> transport in red blood cells of Dahl rats. Effects of age and salt. *Hypertension* 1990;15:612-627. <https://doi.org/10.1161/01.HYP.15.6.612>
81. Zicha J, Negrin CD, Dobešová Z, Carr F, Vokurková M, McBride MW, Kuneš J, Dominiczak AF. Altered Na<sup>+</sup>-K<sup>+</sup> pump activity and plasma lipids in salt-hypertensive Dahl rats: relationship to Atp1a1 gene. *Physiol Genomics* 2001;6:99-104. <https://doi.org/10.1152/physiolgenomics.2001.6.2.99>
82. Vokurková M, Dobešová Z, Kuneš J, Zicha J. Membrane ion transport in erythrocytes of salt hypertensive Dahl rats and their F2 hybrids: the importance of cholesterol. *Hypertens Res* 2003;26:397-404. <https://doi.org/10.1291/hypres.26.397>
83. Vokurková M, Nováková O, Dobešová Z, Kuneš J, Zicha J. Relationships between membrane lipids and ion transport in red blood cells of Dahl rats. *Life Sci* 2005;77:1452-1464. <https://doi.org/10.1016/j.lfs.2005.03.014>
84. Bin Talib HK, Zicha J. Red cell sodium in DOCA-salt hypertension: a Brattleboro study. *Life Sci* 1992;50:1021-1030. [https://doi.org/10.1016/0024-3205\(92\)90097-9](https://doi.org/10.1016/0024-3205(92)90097-9)
85. Bin Talib HK, Kuneš J, Zicha J. Complete dissociation of DOCA-salt hypertension and red cell ion transport alterations. *Life Sci* 1992;51:PL243-PL248. [https://doi.org/10.1016/0024-3205\(92\)90044-P](https://doi.org/10.1016/0024-3205(92)90044-P)
86. Kuneš J, Bin Talib HK, Dobešová Z, Vrána A, Zicha J. Erythrocyte ion transport alterations in hypertriglyceridaemic rats. *Clin Sci (Lond)* 1994;86:11-13. <https://doi.org/10.1042/cs0860011>
87. Bin Talib HK, Dobešová Z, Klír P, Křen V, Kuneš J, Pravenec M, Zicha J. Association of red blood cell sodium leak with blood pressure in recombinant inbred strains. *Hypertension* 1992;20:575-582. <https://doi.org/10.1161/01.HYP.20.4.575>
88. Orlov SN, Petrunyaka VV, Pokudin NI, Kotelevtsev YV, Postnov YV, Kunes J, Zicha J. Cation transport and adenosine triphosphatase activity in rat erythrocytes: a comparison of spontaneously hypertensive rats with the normotensive Brown-Norway strain. *J Hypertens* 1991;9:977-982. <https://doi.org/10.1097/00004872-199110000-00012>
89. Zicha J. Red cell ion transport abnormalities in experimental hypertension. *Fundam Clin Pharmacol.* 1993;7(3-4):129-41. <https://doi.org/10.1111/j.1472-8206.1993.tb00227.x>
90. Zicha J, Kuneš J. Membrane defects and the development of hypertension. In: *Handbook of Hypertension*, Vol. 19, Development of Hypertensive Phenotype: Basic and Clinical Studies, R. McCarty, D.A. Blizard, R.L. Chevalier (eds), Elsevier, Amsterdam, 1999, pp 213-251.
91. Hamet P, Tremblay J, Malo D, Kuneš J, Hashimoto T. Genetic hypertension is characterized by the abnormal expression of a gene localized in major histocompatibility complex HSP70. *Transplant Proc* 1990;22:2566-2567.
92. Hamet P, Kong D, Pravenec M, Kuneš J, Křen V, Klír P, Sun YL, Tremblay J. Restriction fragment length polymorphism of hsp70 gene, localized in the RT1 complex, is associated with hypertension in spontaneously hypertensive rats. *Hypertension* 1992;19:611-614. <https://doi.org/10.1161/01.HYP.19.6.611.a>

93. Kuneš J, Poirier M, Tremblay J, Hamet P. Expression of hsp70 gene in lymphocytes from normotensive and hypertensive humans. *Acta Physiol Scand* 1992;146:307-311. <https://doi.org/10.1111/j.1748-1716.1992.tb09424.x>
94. Hamet P, Kaiser MA, Sun Y, Pagé V, Vincent M, Křen V, Pravenec M, Kuneš J, Tremblay J, Samani NJ. HSP27 locus cosegregates with left ventricular mass independently of blood pressure. *Hypertension* 1996;28:1112-1117.
95. Hamet P, Sun YL, Malo D, Kong D, Křen V, Pravenec M, Kuneš J, Dumas P, Richard L, Gagnon F, et al. Genes of stress in experimental hypertension. *Clin Exp Pharmacol Physiol*. 1994;21:907-911. <https://doi.org/10.1111/j.1440-1681.1994.tb02464.x>
96. Kuneš J, Zicha J. Association of salt sensitivity in rats with genes of the major histocompatibility complex. *Hypertension* 1994;24:645-647. <https://doi.org/10.1161/01.HYP.24.6.645>
97. Vrána A, Kazdová L. The hereditary hypertriglyceridemic nonobese rat: an experimental model of human hypertriglyceridemia. *Transplant Proc* 1990;22:2579.
98. Štolba P, Dobešová Z, Hušek P, Opltová H, Zicha J, Vrána A, Kuneš J. The hypertriglyceridemic rat as a genetic model of hypertension and diabetes. *Life Sci* 1992;51:733-740. [https://doi.org/10.1016/0024-3205\(92\)90482-5](https://doi.org/10.1016/0024-3205(92)90482-5)
99. Zicha J, Dobešová Z, Kuneš J. Plasma triglycerides and red cell ion transport alterations in genetically hypertensive rats. *Hypertension* 1997;30:636-640. <https://doi.org/10.1161/01.HYP.30.3.636>
100. Ueno T, Tremblay J, Kuneš J, Zicha J, Dobešová Z, Pausova Z, Deng AY, Sun YL, Jacob HJ, Hamet P. Resolving the composite trait of hypertension into its pharmacogenetic determinants by acute pharmacological modulation of blood pressure regulatory systems. *J Mol Med (Berl)* 2003;81:51-60. <https://doi.org/10.1007/s00109-002-0394-7>
101. Ueno T, Tremblay J, Kuneš J, Zicha J, Dobešová Z, Pausova Z, Deng AY, Sun Y, Jacob HJ, Hamet P. Gender-specific genetic determinants of blood pressure and organ weight: pharmacogenetic approach. *Physiol Res* 2003;52:689-700. <https://doi.org/10.33549/physiolres.930000.52.689>
102. Ueno T, Tremblay J, Kuneš J, Zicha J, Dobešová Z, Pausova Z, Deng AY, Sun YL, Jacob HJ, Hamet P. Rat model of familial combined hyperlipidemia as a result of comparative mapping. *Physiol Genomics* 2004;17:38-47. <https://doi.org/10.1152/physiolgenomics.00043.2003>
103. Kadlecová M, Čejka J, Zicha J, Kuneš J. Does Cd36 gene play a key role in disturbed glucose and fatty acid metabolism in Prague hypertensive hypertriglyceridemic rats? *Physiol Res* 2004;53:265-271. <https://doi.org/10.33549/physiolres.930000.53.265>
104. Kadlecová M, Dobešová Z, Zicha J, Kuneš J. Abnormal Igf2 gene in Prague hereditary hypertriglyceridemic rats: its relation to blood pressure and plasma lipids. *Mol Cell Biochem* 2008;314:37-43. <https://doi.org/10.1007/s11010-008-9762-0>
105. Klimeš I, Vrána A, Kuneš J, Šeböková E, Dobešová Z, Štolba P, Zicha J. Hereditary hypertriglyceridemic rat: a new animal model of metabolic alterations in hypertension. *Blood Press* 1995;4:137-142. <https://doi.org/10.3109/08037059509077585>
106. Zicha J, Pecháňová O, Čačányiová S, Cebová M, Kristek F, Török J, Šimko F, Dobešová Z, Kuneš J. Hereditary hypertriglyceridemic rat: a suitable model of cardiovascular disease and metabolic syndrome? *Physiol Res* 2006;55 (Suppl 1):S49-S63. <https://doi.org/10.33549/physiolres.930000.55.S1.49>
107. Zicha J, Kuneš J, Le Quan Sang KH, Devynck MA. Regulation of the dynamic properties of platelet plasma membrane by intracellular sodium ions. *Life Sci* 1993;52:1559-1565. [https://doi.org/10.1016/0024-3205\(93\)90056-9](https://doi.org/10.1016/0024-3205(93)90056-9)
108. Le Quan Sang KH, Kuneš J, Zicha J, Vincent M, Sassard J, Devynck MA. Platelet and erythrocyte membrane microviscosity in Lyon hypertensive rats. *Am J Hypertens* 1994;7:276-281. <https://doi.org/10.1093/ajh/7.3.276>
109. Kuneš J, Zicha J, Devynck MA. Erythrocyte membrane microviscosity and blood pressure in rats with salt-induced and spontaneous hypertension. *J Hypertens* 1994;12:229-234. <https://doi.org/10.1097/00004872-199403000-00002>
110. Zicha J, Pernollet MG, Kuneš J, Lacour B, Vincent M, Sassard J, Devynck MA. Alterations of cytosolic calcium in platelets and erythrocytes of Lyon hypertensive rats. *Am J Hypertens* 1995;8:842-849. [https://doi.org/10.1016/0895-7061\(95\)00121-5](https://doi.org/10.1016/0895-7061(95)00121-5)
111. Zicha J, Kuneš J, David-Dufilho M, Pernollet MG, Devynck MA. Cell calcium handling and intracellular pH regulation in hereditary hypertriglyceridemic rats: reduced platelet response to thrombin stimulation. *Life Sci* 1996;59:803-813. [https://doi.org/10.1016/0024-3205\(96\)00371-2](https://doi.org/10.1016/0024-3205(96)00371-2)

112. Devynck MA, Kuneš J, Le Quan Sang KH, Zicha J. Membrane microviscosity and plasma triacylglycerols in the rat. *Clin Sci (Lond)* 1998;94:79-85. <https://doi.org/10.1042/cs0940079>
113. Zicha J, Kuneš J, Devynck MA. Abnormalities of membrane function and lipid metabolism in hypertension: a review. *Am J Hypertens* 1999;12:315-331. [https://doi.org/10.1016/S0895-7061\(98\)00178-2](https://doi.org/10.1016/S0895-7061(98)00178-2)
114. Kuneš J, Mazeaud MM, Devynck MA, Zicha J. Platelet hypoaggregability in hereditary hypertriglyceridemic rats: relation to plasma triglycerides. *Thromb Res* 1997;88:347-353. [https://doi.org/10.1016/S0049-3848\(97\)00264-8](https://doi.org/10.1016/S0049-3848(97)00264-8)
115. Kuneš J, Devynck MA, Zicha J. Chronic changes in plasma triglyceride levels do modify platelet membrane microviscosity in rats. *Life Sci* 2000;67:959-967. [https://doi.org/10.1016/S0024-3205\(00\)00691-3](https://doi.org/10.1016/S0024-3205(00)00691-3)
116. Zicha J, Kuneš J, Devynck MA. Platelet calcium handling is different in rats with salt-dependent and spontaneous forms of genetic hypertension. *Am J Hypertens* 1996;9:812-818. [https://doi.org/10.1016/0895-7061\(96\)00101-X](https://doi.org/10.1016/0895-7061(96)00101-X)
117. Zicha J, Kuneš J, Ben-Ishay D, Devynck MA. Abnormal regulation of cytosolic calcium and pH in platelets of Sabra rats in early phases of salt hypertension development. *Can J Physiol Pharmacol* 1996;74:1222-1228. <https://doi.org/10.1139/y96-134>
118. Zicha J, David-Dufilho M, Kuneš J, Pernollet MG, Devynck MA. Cytosolic pH and calcium in Dahl salt-sensitive and salt-resistant rats: the relationship to plasma lipids. *J Hypertens* 1997;15:1715-1721. <https://doi.org/10.1097/00004872-199715120-00078>
119. Zicha J, Sang KH, Kuneš J, Devynck MA. Membrane microviscosity, blood pressure and cytosolic pH in Dahl rats: the influence of plasma lipids. *J Hypertens* 1999;17:785-792. <https://doi.org/10.1097/00004872-199917060-00010>
120. Červenka L, Heller J. A comparison of the effect of feeding a low-protein diet and of pharmacological intervention on the course of ablation nephropathy in the rat. *Ren Fail* 1995;17:27-35. <https://doi.org/10.3109/08860229509036372>
121. Červenka L, Heller J. Comparison of the effects of a low-protein diet with the effects of a converting enzyme inhibitor on the progression of renal insufficiency in hypertensive rats. *Ren Fail* 1996;18:173-180. <https://doi.org/10.3109/08860229609052787>
122. Červenka L, Wang CT, Navar LG. Effects of acute AT1 receptor blockade by candesartan on arterial pressure and renal function in rats. *Am J Physiol* 1998;274:F940-F945. <https://doi.org/10.1152/ajprenal.1998.274.5.F940>
123. Červenka L, Wang CT, Mitchell KD, Navar LG. Proximal tubular angiotensin II levels and renal functional responses to AT1 receptor blockade in nonclipped kidneys of Goldblatt hypertensive rats. *Hypertension* 1999;33:102-107. <https://doi.org/10.1161/01.HYP.33.1.102>
124. Červenka L, Navar LG. Renal responses of the nonclipped kidney of two-kidney/one-clip Goldblatt hypertensive rats to type 1 angiotensin II receptor blockade with candesartan. *J Am Soc Nephrol* 1999;10 (Suppl 11):S197-S201.
125. Červenka L, Horáček V, Vaněčková I, Hubáček JA, Oliverio MI, Coffman TM, Navar LG. Essential role of AT1A receptor in the development of 2K1C hypertension. *Hypertension* 2002;40:735-741. <https://doi.org/10.1161/01.HYP.0000036452.28493.74>
126. Červenka L, Harrison-Bernard LM, Dipp S, Primrose G, Imig JD, El-Dahr SS. Early onset salt-sensitive hypertension in bradykinin B2 receptor null mice. *Hypertension* 1999;34:176-180. <https://doi.org/10.1161/01.HYP.34.2.176>
127. Červenka L, Malý J, Karasová L, Šimová M, Vítko Š, Hellerová S, Heller J, El-Dahr SS. Angiotensin II-induced hypertension in bradykinin B2 receptor knockout mice. *Hypertension* 2001;37:967-973. <https://doi.org/10.1161/01.HYP.37.3.967>
128. Červenka L, Vaněčková I, Malý J, Horáček V, El-Dahr SS. Genetic inactivation of the B2 receptor in mice worsens two-kidney, one-clip hypertension: role of NO and the AT2 receptor. *J Hypertens* 2003;21:1531-1538. <https://doi.org/10.1097/00004872-200308000-00018>
129. Kopkan L, Kramer HJ, Husková Z, Vaňourková Z, Bäcker A, Bader M, Ganten D, Červenka L. Plasma and kidney angiotensin II levels and renal functional responses to AT1 receptor blockade in hypertensive Ren-2 transgenic rats. *J Hypertens* 2004;22:819-825. <https://doi.org/10.1097/00004872-200404000-00026>

130. Červenka L, Vaněčková I, Husková Z, Vaňourková Z, Erbanová M, Thumová M, Škaroupková P, Opočenský M, Malý J, Čertíková Chábová V, Tesař V, Bürgelová M, Viklický O, Teplan V, Želízko M, Kramer HJ, Navar LG. Pivotal role of angiotensin II receptor subtype 1A in the development of two-kidney, one-clip hypertension: study in angiotensin II receptor subtype 1A knockout mice. *J Hypertens*. 2008 Jul;26(7):1379-89. <https://doi.org/10.1097/HJH.0b013e3282fe6eaa>
131. Bürgelová M, Kramer HJ, Teplan V, Thumová M, Červenka L. Effects of angiotensin-(1-7) blockade on renal function in rats with enhanced intrarenal Ang II activity. *Kidney Int* 2005;67:1453-1461. <https://doi.org/10.1111/j.1523-1755.2005.00222.x>
132. Bürgelová M, Vaňourková Z, Thumová M, Dvořák P, Opočenský M, Kramer HJ, Želízko M, Malý J, Bader M, Červenka L. Impairment of the angiotensin-converting enzyme 2-angiotensin-(1-7)-Mas axis contributes to the acceleration of two-kidney, one-clip Goldblatt hypertension. *J Hypertens* 2009;27:1988-2000. <https://doi.org/10.1097/HJH.0b013e32832f0d06>
133. Červenka L, Kramer HJ, Malý J, Heller J. Role of nNOS in regulation of renal function in angiotensin II-induced hypertension. *Hypertension* 2001;38:280-285. <https://doi.org/10.1161/01.HYP.38.2.280>
134. Červenka L, Kramer HJ, Malý J, Vaněčková I, Bäcker A, Bokemeyer D, Bader M, Ganten D, Mitchell KD. Role of nNOS in regulation of renal function in hypertensive Ren-2 transgenic rats. *Physiol Res* 2002;51:571-580. <https://doi.org/10.33549/physiolres.930316>
135. Vaněčková I, Kramer HJ, Novotná J, Kazdová L, Opočenský M, Bader M, Ganten D, Červenka L. Roles of nitric oxide and oxidative stress in the regulation of blood pressure and renal function in prehypertensive Ren-2 transgenic rats. *Kidney Blood Press Res* 2005;28:117-126. <https://doi.org/10.1159/000084649>
136. Kopkan L, Husková Z, Vaňourková Z, Thumová M, Škaroupková P, Červenka L, Majid DS. Superoxide and its interaction with nitric oxide modulates renal function in prehypertensive Ren-2 transgenic rats. *J Hypertens* 2007;25:2257-2265. <https://doi.org/10.1097/HJH.0b013e3282efb195>
137. Kopkan L, Červenka L. Renal interactions of renin-angiotensin system, nitric oxide and superoxide anion: implications in the pathophysiology of salt-sensitivity and hypertension. *Physiol Res* 2009;58 (Suppl 2):S55-S68. <https://doi.org/10.33549/physiolres.931917>
138. Husková Z, Kramer HJ, Vaňourková Z, Červenka L. Effects of changes in sodium balance on plasma and kidney angiotensin II levels in anesthetized and conscious Ren-2 transgenic rats. *J Hypertens* 2006;24:517-527. <https://doi.org/10.1097/01.hjh.0000209988.51606.c7>
139. Husková Z, Kramer H, Vaňourková Z, Thumová M, Malý J, Opočenský M, Škaroupková P, Kolský A, Vernerová Z, Červenka L. Effects of dietary salt load and salt depletion on the course of hypertension and angiotensin II levels in male and female heterozygous Ren-2 transgenic rats. *Kidney Blood Press Res* 2007;30:45-55. <https://doi.org/10.1159/000099028>
140. Husková Z, Kramer HJ, Thumová M, Vaňourková Z, Bürgelová M, Teplan V, Malý J, Červenka L. Effects of anesthesia on plasma and kidney ANG II levels in normotensive and ANG II-dependent hypertensive rats. *Kidney Blood Press Res* 2006;29:74-83. <https://doi.org/10.1159/000092981>
141. Dvořák P, Kramer HJ, Bäcker A, Malý J, Kopkan L, Vaněčková I, Vernerová Z, Opočenský M, Tesař V, Bader M, Ganten D, Janda J, Červenka L. Blockade of endothelin receptors attenuates end-organ damage in homozygous hypertensive ren-2 transgenic rats. *Kidney Blood Press Res* 2004;27:248-258. <https://doi.org/10.1159/000080052>
142. Vaněčková I, Kramer HJ, Bäcker A, Vernerová Z, Opočenský M, Červenka L. Early endothelin-A receptor blockade decreases blood pressure and ameliorates end-organ damage in homozygous Ren-2 rats. *Hypertension* 2005;46:969-974. <https://doi.org/10.1161/01.HYP.0000173426.06832.b5>
143. Opočenský M, Kramer HJ, Bäcker A, Vernerová Z, Eis V, Červenka L, Čertíková Chábová V, Tesař V, Vaněčková I. Late-onset endothelin-A receptor blockade reduces podocyte injury in homozygous Ren-2 rats despite severe hypertension. *Hypertension* 2006;48:965-971. <https://doi.org/10.1161/01.HYP.0000245117.57524.d6>
144. Vernerová Z, Kramer HJ, Bäcker A, Červenka L, Opočenský M, Husková Z, Vaňourková Z, Eis V, Chábová VC, Tesař V, Malý J, Vaněčková I. Late-onset endothelin receptor blockade in hypertensive heterozygous REN-2 transgenic rats. *Vascul Pharmacol* 2008;48:165-173. <https://doi.org/10.1016/j.vph.2008.01.009>

145. Čertíková Chábová V, Kramer HJ, Vaněčková I, Thumová M, Škaroupková P, Tesař V, Falck JR, Imig JD, Červenka L. The roles of intrarenal 20-hydroxyeicosatetraenoic and epoxyeicosatrienoic acids in the regulation of renal function in hypertensive Ren-2 transgenic rats. *Kidney Blood Press Res* 2007;30:335-346. <https://doi.org/10.1159/000107710>
146. Čertíková Chábová V, Kramer HJ, Vaněčková I, Vernerová Z, Eis V, Tesař V, Škaroupková P, Thumová M, Schejbalová S, Husková Z, Vaňourková Z, Kolský A, Imig JD, Červenka L. Effects of chronic cytochrome P-450 inhibition on the course of hypertension and end-organ damage in Ren-2 transgenic rats. *Vascul Pharmacol* 2007;47:145-159. <https://doi.org/10.1016/j.vph.2007.05.005>
147. Čertíková Chábová V, Walkowska A, Kompanowska-Jeziarska E, Sadowski J, Kujal P, Vernerová Z, Vaňourková Z, Kopkan L, Kramer HJ, Falck JR, Imig JD, Hammock BD, Vaněčková I, Červenka L. Combined inhibition of 20-hydroxyeicosatetraenoic acid formation and of epoxyeicosatrienoic acids degradation attenuates hypertension and hypertension-induced end-organ damage in Ren-2 transgenic rats. *Clin Sci (Lond)* 2010;118:617-632. <https://doi.org/10.1042/CS20090459>
148. Vaňourková Z, Kramer HJ, Husková Z, Vaněčková I, Opočenský M, Čertíková Chábová V, Tesař V, Škaroupková P, Thumová M, Dohnalová M, Mullins JJ, Červenka L. AT1 receptor blockade is superior to conventional triple therapy in protecting against end-organ damage in Cyp11 $\alpha$ -Ren-2 transgenic rats with inducible hypertension. *J Hypertens* 2006;24:2465-2472. <https://doi.org/10.1097/01.hjh.0000251909.00923.22>
149. Erbanová M, Thumová M, Husková Z, Vaněčková I, Vaňourková Z, Mullins JJ, Kramer HJ, Bürgelová M, Rakušan D, Červenka L. Impairment of the autoregulation of renal hemodynamics and of the pressure-natriuresis relationship precedes the development of hypertension in Cyp11 $\alpha$ -Ren-2 transgenic rats. *J Hypertens* 2009;27:575-586. <https://doi.org/10.1097/HJH.0b013e32831cbd5a>
150. Bačáková L, Kuneš J. Sex-dependent differences in growth of vascular smooth muscles cells from spontaneously hypertensive rats. *Physiol Res* 1995;44:127-130.
151. Bačáková L, Kuneš J. Gender differences in growth of vascular smooth muscle cells isolated from hypertensive and normotensive rats. *Clin Exp Hypertens*. 2000 Jan;22(1):33-44. <https://doi.org/10.1081/CEH-100100060>
152. Loukotová J, Bačáková L, Zicha J, Kuneš J. The influence of angiotensin II on sex-dependent proliferation of aortic VSMC isolated from SHR. *Physiol Res* 1998;47:501-505.
153. Loukotová J, Kuneš J, Zicha J. Gender-dependent difference in cell calcium handling in VSMC isolated from SHR: the effect of angiotensin II. *J Hypertens* 2002;20:2213-2219. <https://doi.org/10.1097/00004872-200211000-00021>
154. Zicha J, Kuneš J, Lébl M, Pohlová I, Jelínek J. Haemodynamics and the participation of pressor systems in young and adult rats with age-dependent DOCA-salt hypertension. *Physiol Bohemoslov* 1987;36:89-92.
155. Hatzinikolaou P, Gavras H, Brunner HR, Gavras I. Role of vasopressin, catecholamines, and plasma volume in hypertonic saline-induced hypertension. *Am J Physiol* 1981 Jun;240:H827-H831. <https://doi.org/10.1152/ajpheart.1981.240.6.H827>
156. Gavras H, Hatzinikolaou P, North WG, Bresnahan M, Gavras I. Interaction of the sympathetic nervous system with vasopressin and renin in the maintenance of blood pressure. *Hypertension* 1982;4:400-405. <https://doi.org/10.1161/01.HYP.4.3.400>
157. Burnier M, Biollaz J, Brunner DB, Gavras H, Brunner HR. Alpha and beta adrenoceptor blockade in normotensive and deoxycorticosterone (DOC)-hypertensive rats; plasma vasopressin and vasopressin pressor effect. *J Pharmacol Exp Ther* 1983;224:222-227.
158. Waeber B, Nussberger J, Brunner HR. Alpha 1-adrenoceptor blockade in renal hypertensive rats with low and high renin levels. *Am J Physiol* 1984;246:H573-H578. <https://doi.org/10.1152/ajpheart.1984.246.4.H573>
159. Minami N, Imai Y, Hashimoto J, Abe K. Contribution of vascular nitric oxide to basal blood pressure in conscious spontaneously hypertensive rats and normotensive Wistar Kyoto rats. *Clin Sci (Lond)* 1995;89:177-182. <https://doi.org/10.1042/cs0890177>
160. Kazda S, Pohlová I, Bíbr B, Kočková J. Norepinephrine content of tissues in Doca-hypertensive rats. *Am J Physiol* 1969;216:1472-1475. <https://doi.org/10.1152/ajplegacy.1969.216.6.1472>



161. Křeček J, Doležel S, Dlouhá H, Zicha J. Sympathetic nervous system and age-dependent salt hypertension in Brattleboro rats. *Clin Exp Hypertens A* 1987;9:2075-2093. <https://doi.org/10.1080/07300077.1987.11978716>, <https://doi.org/10.3109/10641968709159076>
162. Zicha J, Dobešová Z, Kuneš J. Relative deficiency of nitric oxide-dependent vasodilation in salt-hypertensive Dahl rats: the possible role of superoxide anions. *J Hypertens* 2001;19:247-254. <https://doi.org/10.1097/00004872-200102000-00011>
163. Kuneš J, Dobešová Z, Zicha J. Altered balance of main vasopressor and vasodepressor systems in rats with genetic hypertension and hypertriglyceridaemia. *Clin Sci (Lond)* 2002;102:269-277. <https://doi.org/10.1042/cs1020269>
164. Dobešová Z, Kuneš J, Zicha J. The altered balance between sympathetic nervous system and nitric oxide in salt hypertensive Dahl rats: ontogenetic and F2 hybrid studies. *J Hypertens* 2002;20:945-955. <https://doi.org/10.1097/00004872-200205000-00030>
165. Pecháňová O, Bernátová I, Pelouch V, Šimko F. Protein remodelling of the heart in NO-deficient hypertension: the effect of captopril. *J Mol Cell Cardiol* 1997;29:3365-3374. <https://doi.org/10.1006/jmcc.1997.0566>
166. Bernátová I, Pecháňová O, Pelouch V, Šimko F. Regression of chronic L-NAME-treatment-induced left ventricular hypertrophy: effect of captopril. *J Mol Cell Cardiol* 2000;32:177-185. <https://doi.org/10.1006/jmcc.1999.1071>
167. Bernátová I, Pecháňová O, Babál P, Kyselá S, Štvrtina S, Andriantsitohaina R. Wine polyphenols improve cardiovascular remodeling and vascular function in NO-deficient hypertension. *Am J Physiol Heart Circ Physiol* 2002;282:H942-H948. <https://doi.org/10.1152/ajpheart.00724.2001>
168. Pecháňová O, Dobešová Z, Čejka J, Kuneš J, Zicha J. Vasoactive systems in L-NAME hypertension: the role of inducible nitric oxide synthase. *J Hypertens* 2004;22:167-173. <https://doi.org/10.1097/00004872-200401000-00026>
169. Zicha J, Pecháňová O, Dobešová Z, Kuneš J. Hypertensive response to chronic NG-nitro-L-arginine methyl ester (L-NAME) treatment is similar in immature and adult Wistar rats. *Clin Sci (Lond)* 2003;105:483-489. <https://doi.org/10.1042/CS20030078>
170. Rauchová H, Pecháňová O, Kuneš J, Vokurková M, Dobešová Z, Zicha J. Chronic N-acetylcysteine administration prevents development of hypertension in N<sup>ω</sup>-nitro-L-arginine methyl ester-treated rats: the role of reactive oxygen species. *Hypertens Res* 2005;28:475-482. <https://doi.org/10.1291/hypres.28.475>
171. Zicha J, Dobešová Z, Kuneš J. Antihypertensive mechanisms of chronic captopril or N-acetylcysteine treatment in L-NAME hypertensive rats. *Hypertens Res* 2006;29:1021-1027. <https://doi.org/10.1291/hypres.29.1021>
172. Zicha J, Kuneš J, Vranková S, Jendeková L, Dobešová Z, Pintéřová M, Pecháňová O. Influence of pertussis toxin pretreatment on the development of L-NAME-induced hypertension. *Physiol Res* 2009;58:751-755. <https://doi.org/10.33549/physiolres.931898>
173. Pecháňová O, Zicha J, Kojšová S, Dobešová Z, Jendeková L, Kuneš J. Effect of chronic N-acetylcysteine treatment on the development of spontaneous hypertension. *Clin Sci (Lond)*. 2006;110:235-242. <https://doi.org/10.1042/CS20050227>
174. Pecháňová O, Zicha J, Paulis L, Zenebe W, Dobešová Z, Kojšová S, Jendeková L, Sládková M, Dvořáková I, Šimko F, Kuneš J. The effect of N-acetylcysteine and melatonin in adult spontaneously hypertensive rats with established hypertension. *Eur J Pharmacol*. 2007;561:129-136. <https://doi.org/10.1016/j.ejphar.2007.01.035>
175. Kuneš J, Zicha J. Developmental windows and environment as important factors in the expression of genetic information: a cardiovascular physiologist's view. *Clin Sci (Lond)* 2006;111:295-305. <https://doi.org/10.1042/CS20050271>
176. Kuneš J, Zicha J. The interaction of genetic and environmental factors in the etiology of hypertension. *Physiol Res* 2009;58 (Suppl 2):S33-S42. <https://doi.org/10.33549/physiolres.931913>
177. Bengel HH, Solomon S. Development of renal response to blood volume expansion in the rat. *Am J Physiol* 1974;227:364-368. <https://doi.org/10.1152/ajplegacy.1974.227.2.364>
178. Brownie AC, Bernardis LL, Niwa T, Kamura S, Skelton FR. The influence of age and sex on the development of adrenal regeneration hypertension. *Lab Invest* 1966;15:1342-1356.

- 
179. Dahl LK, Knudsen KD, Heine M, Leitel G. Effects of chronic excess salt ingestion. Genetic influence on the development of salt hypertension in parabiotic rats: evidence for a humoral factor. *J Exp Med* 1967;126:687-699. <https://doi.org/10.1084/jem.126.4.687>
  180. Self LE, Battarbee HD, Gaar KA Jr, Meneely GR. A vasopressor potentiator for norepinephrine in hypertensive rats. *Proc Soc Exp Biol Med* 1976;153:7-12. <https://doi.org/10.3181/00379727-153-39470>
  181. Kojima I. Circulating digitalis-like substance is increased in DOCA-salt hypertension. *Biochem Biophys Res Commun* 1984;122:129-136. [https://doi.org/10.1016/0006-291X\(84\)90449-2](https://doi.org/10.1016/0006-291X(84)90449-2)
  182. Alaghband-Zadeh J, Fenton S, Hancock K, Millett J, de Wardener HE. Evidence that the hypothalamus may be a source of a circulating Na<sup>+</sup>-K<sup>+</sup>-ATPase inhibitor. *J Endocrinol* 1983;98:221-226. <https://doi.org/10.1677/joe.0.0980221>
  183. Castaneda-Hernandez G, Godfraind T. Effect of high sodium intake on tissue distribution of endogenous digitalis-like material in the rat. *Clin Sci (Lond)* 1984;66:225-228. <https://doi.org/10.1042/cs0660225>
  184. Abel PW, Trapani A, Matsuki N, Ingram MJ, Ingram FD, Hermsmeyer K. Unaltered membrane properties of arterial muscle in Dahl strain genetic hypertension. *Am J Physiol* 1981;241:H224-H227. <https://doi.org/10.1152/ajpheart.1981.241.2.H224>
  185. Overbeck HW, Ku DD, Rapp JP. Sodium pump activity in arteries of Dahl salt-sensitive rats. *Hypertension* 1981;3:306-312. <https://doi.org/10.1161/01.HYP.3.3.306>
  186. Pamnani MB, Clough DL, Huot SJ, Haddy FJ. Vascular Na<sup>+</sup>-K<sup>+</sup> pump activity in Dahl S and R rats. *Proc Soc Exp Biol Med* 1980;165:440-444. <https://doi.org/10.3181/00379727-165-41001>
-