

Blockade of α 2-Adrenergic Receptors in the Caudal Raphe Region Enhances the Renal Sympathetic Nerve Activity Response to Acute Intermittent Hypercapnia in Rats

Katarina MADIRAZZA¹, Renata PECOTIC¹, Ivana PAVLINAC DODIG¹, Maja VALIC¹, Zoran DOGAS¹

¹Department of Neuroscience, University of Split School of Medicine, Split, Croatia

Received April 30, 2021

Accepted November 11, 2021

Epub Ahead of Print January 19, 2022

Summary

The study investigated the role of α 2-adrenergic receptors of the caudal raphe region in the sympathetic and cardiovascular responses to the acute intermittent hypercapnia (AIHc). Urethane-anesthetized, vagotomized, mechanically ventilated Sprague-Dawley rats (n=38) were exposed to the AIHc protocol (5×3 min, 15 % CO₂+50 % O₂) in hyperoxic background (50 % O₂). α 2-adrenergic receptor antagonist-yohimbine was applied intravenously (1 mg/kg, n=9) or microinjected into the caudal raphe region (2 mM, n=12) prior to exposure to AIHc. Control groups of animals received saline intravenously (n=7) or into the caudal raphe region (n=10) prior to exposure to AIHc. Renal sympathetic nerve activity (RSNA), mean arterial pressure (MAP) and heart rate (HR) were monitored before exposure to the AIHc protocol (T0), during five hypercapnic episodes (THc1-5) and at 15 min following the end of the last hypercapnic episode (T15). Following intravenous administration of yohimbine, RSNA was significantly greater during THc1-5 and at T15 than in the control group (P<0.05). When yohimbine was microinjected into the caudal raphe region, AIHc elicited greater increases in RSNA during THc1-5 when compared to the controls (THc1: 138.0±4.0 % vs. 123.7±4.8 %, P=0.032; THc2: 137.1±5.0 % vs. 124.1±4.5 %, P=0.071; THc3: 143.1±6.4 % vs. 122.0±4.8 %, P=0.020; THc4: 146.1±6.2 % vs. 120.7±5.7 %, P=0.007 and THc5: 143.2±7.7 % vs. 119.2±7.2 %, P=0.038). During THc1-5, significant decreases in HR from T0 were observed in all groups, while changes in MAP were observed in the group that received yohimbine intravenously. These findings suggest that blockade of the α 2-adrenergic receptors in the caudal raphe region might have an important role in sympathetic responses to AIHc.

Key words

Acute intermittent hypercapnia • Sympathetic nervous activity • α 2-adrenergic receptors • Raphe nuclei • Yohimbine

Corresponding author

R. Pecotic, Department of Neuroscience, University of Split School of Medicine, Soltanska 2, 21000 Split, Croatia. E-mail: renata.pecotic@mefst.hr

Introduction

Hypercapnia, which is a hallmark of many respiratory disorders, has been shown to have a substantial effect on various organ systems [1]. It is known to stimulate both peripheral and central chemoreceptors [2,3], consequently driving hyperventilation, raised blood pressure and elevated sympathetic nerve activity [1,4]. Hypercapnia-induced ventilatory response has been well described previously, such that the elevation of the arterial partial pressure of carbon dioxide (PaCO₂) leads to a strong activation of the areas controlling the respiratory activity, and the response seems to be largely dependent on the medullary raphe neurons [5-7]. Moreover, when administered intermittently, hypercapnia is documented to cause a long-term depression of breathing modulated by both, serotonergic and adrenergic receptors [8-12].

Although a large pool of evidence shows single episode or sustained hypercapnia potently stimulate the sympathetic nervous system [1,3,13-15], the effects of intermittent hypercapnia on sympathetic activity remain

unclear. Recently, our laboratory demonstrated that the acute intermittent hypercapnia (AIHc) causes substantial activation of the sympathetic nervous system measured through the renal sympathetic nerve activity (RSNA), which is dependent on the background oxygen level [16]. Evaluating different protocols, we found that severe acute intermittent hyperoxic hypercapnia (15 % CO₂+50 % O₂) modestly activates the RSNA, but the magnitude of the response is significantly lower than in a normoxic background, suggesting the involvement of a central mechanism.

Central neural pathways controlling the cardiorespiratory outflow are complex and to this date not fully worked out [17]. Undoubtedly, central chemoreception evokes increases in sympathetic nervous activity by direct action on cardiovascular centers or mediated by the central respiratory network [17,18]. Thus far, different regions have been proposed to act as central CO₂ chemoreceptors [14,15,18-20], among which locus coeruleus (LC) and raphe nuclei have been reported to be extremely responsive to elevations in PaCO₂ [6,19-22].

Projections of the noradrenergic groups participate in increases in breathing and sympathetic nerve activity caused by hypercapnia or reduced pH in noradrenergic brainstem regions [20,21]. Dense projections from LC to the caudal raphe region exist, where they can induce, either, excitatory effects *via* α 1-adrenergic receptors or inhibitory effects *via* α 2-adrenergic receptors [23]. Nuclei of the caudal raphe region (nucl. raphe magnus, obscurus and pallidus) send extensive descending projections to the spinal cord, innervating the dorsal and ventral horns and the intermediolateral cell column (IML) [24]. The caudal raphe region, comprised of mainly serotonergic neurons, has been implicated in many neurophysiological functions including the modulation of pain, respiration, motor activity and various autonomic functions [24-26]. It has been proposed that the caudal raphe nuclei are involved in sympathetic control by means of direct projections to the IML, innervating the preganglionic sympathetic neurons [23,26-28], but also to other regions controlling the sympathetic activity, such as the rostral ventrolateral medulla (RVLM) [29].

In light of this, the aim of the present study was to determine the role of α 2-adrenergic receptors of the caudal raphe region in the sympathetic and cardiovascular responses to the AIHc. The effects of blockade of α 2-adrenergic receptors in the caudal raphe region on renal sympathetic nerve, arterial pressure and heart rate

responses to acute intermittent hypercapnia were examined.

Material and Methods

All experimental procedures were designed in accordance with the European Guidelines on Laboratory Animal Care and were approved by the Ethical Committee for Biomedical Research of the University of Split School of Medicine (Split, Croatia) and the National Ethics Committee of the Veterinary Directorate, Ministry of Agriculture, Republic of Croatia. The animals used in this study were bred and maintained in controlled environment rooms (22-24 °C; 55-70 % relative humidity; 12 h-light: 12 h-dark cycle) with access to food and water *ad libitum* at the University of Split School of Medicine Animal Facility.

General procedures

Male Sprague-Dawley rats (n=38; body weight 280-350 g) were anesthetized by an intraperitoneal injection of urethane (20 % urethane in 0.9 % saline, dose 1.2 g/kg). The adequacy of anesthesia was determined by the hind paw withdrawal and corneal reflex loss. Additional doses of urethane (0.2 g/kg i.v.) were given, if required, in order to maintain a deep anesthesia before exposing the animals to the experimental protocol. Upon achieving the adequate level of anesthesia, femoral arteries and veins were catheterized for arterial blood pressure monitoring, blood gas sampling and drug/fluid administration, respectively. The trachea was cannulated and vagus nerves cut bilaterally at cervical level (C3-5) to facilitate the mechanical ventilation. The animals were ventilated using a 50:50 oxygen-nitrogen mixture throughout the experiment by means of a small animal ventilator (SAR 830-P, CWE, Ardmore, PA, USA). Physiological levels of pH and PaCO₂ were maintained by adjusting the frequency and/or inspiratory time parameters on the ventilator in accordance with the blood gas analysis results (RAPIDPoint 500; Siemens Healthcare Limited, Surrey, UK). Since the animals were maintained on a hyperoxic mixture, the arterial partial pressure of oxygen (PaO₂) was in the range of 250-360 mm Hg. To ensure the overall stability of the preparation, the fluids were supplemented continuously (0.6 ml·h⁻¹·kg⁻¹) and the animals' body temperature was maintained by a heating pad (FST, Heidelberg, Germany) throughout the experiment. The animals were then positioned in a stereotaxic frame (Lab Standard,

Stoelting, Wood Dale, IL, USA) in a prone position and the left renal nerve was exposed using a retroperitoneal approach. The nerve was placed on a bipolar silver wire electrode and the surgical field was covered with silicone gel to secure the electrode, provide electrical noise isolation and prevent desiccation.

Recording parameters

The signal from the recording electrode was amplified (SuperZ, System 1000, CWE Inc., Ardmore, USA), filtered (300 Hz-10 kHz, bandpass filter), full wave rectified and integrated (MA-1000 PowerLab Moving Averager module for System 1000; 50 ms time constant). Chart 5.4.2. for Windows software (ADInstruments, Bella Vista, Australia) was used to obtain the RSNA electrograms, arterial blood pressure tracings (Memscap blood pressure transducer, Skoppum, Norway) and record the signals simultaneously in high resolution at a sampling rate of 20 kHz.

Raphe nuclei identification and microinjection technique

A partial occipital craniotomy was performed and the dura reflected to expose the dorsal surface of the brain stem and the obex. Relative to the obex, the initial coordinates for identification of the caudal raphe region were: +0.2 mm anteroposterior, 0.0 mm mediolateral and -2.5 mm dorsoventral and were determined based on our previous study [11] relying on the rat brain atlas of Paxinos and Watson [30]. A four-barrel glass micropipette (external tip diameter: 30-50 μ m) was filled with the vehicle (0.9 % saline), selective glutamate agonist D,L-homocysteic acid (DLH; 10 mM, Sigma-Aldrich, St Louis, MO, USA), selective α 2-adrenergic receptor antagonist yohimbine (2 mM, Sigma-Aldrich) and diluted India ink solution. The tip of the micropipette was lowered into the position targeting the caudal raphe region according to the initial coordinates. Following the placement of the micropipette, DLH (10 mM in 0.9 % saline; 20 \pm 5 nl, Sigma-Aldrich) was microinjected and RSNA and blood pressure changes monitored. If no response was observed, the coordinates were finely adjusted until an increase in RSNA and a pressor response of >20 mm Hg was evoked [29]. After the transient effects of DLH dissipated, the micropipette was flushed with the vehicle to ensure no glutamate agonist remained in the pipette. Microinjections were performed by a pressure ejection system using a large plastic syringe connected to the barrels by polyethylene tubing. The application of positive pressure allowed for the solution ejection until the final volume reached 20 \pm 5 nl. The

volume ejected was controlled by monitoring the fluid meniscus under a monocular microscope with a finely graduated eyepiece.

Experimental design

The animals were allowed a 30-min stabilization period before the onset of recordings and the exposure to the experimental protocol. Then, the animals were subjected to the AIHc protocol, which consisted of 5 exposures to a hypercapnic gas mixture interspersed by 3 min recovery periods. The hypercapnic gas mixture applied was 15 % CO₂+50 % O₂ in N₂ and the duration of each episode was 3 min. During the recovery periods the animals were ventilated with a hyperoxic mixture of 50 % O₂ in N₂. Two sets of experiments were performed: first, involving the systemic blockade of the α 2-adrenergic receptors and, second, involving the central blockade of the α 2-adrenergic receptors with their respective control groups. In the first set, the yohimbine group (n=9) received a bolus injection of yohimbine (1 mg/kg, 0.6 ml, Sigma-Aldrich) intravenously before exposure to the AIHc protocol. In order to comply with the ethical standards for animal experimentation, one set of data from our previous study was used as a control group (n=7) [16]. In this group, the animals were given the same volume of 0.9 % saline in a bolus before exposure to the AIHc protocol under identical conditions as applied in this study. In the second set of experiments, the yohimbine group (n=12) received a microinjection of α 2-adrenergic receptor antagonist yohimbine (2 mM in 0.9 % saline, 20 \pm 5 nl, Sigma-Aldrich) into the same caudal raphe site previously mapped using DLH and then exposed to the experimental protocol. In the control group (n=10), the same volume of 0.9 % saline was microinjected into the caudal raphe region before the animals were subjected to the AIHc protocol. Before commencing the experimental protocol, an arterial blood sample (0.2 ml) was taken and arterial blood gas values measured. Another control arterial blood sample was taken 15 min following the end of the last hypercapnic episode. No blood samples were taken during the hypercapnic exposures to minimize the overall circulating volume loss and its possible effects on blood pressure and sympathetic activity. The experimental design is shown in Figure 1.

Histological processing

After the completion of each experiment, diluted India ink (20 \pm 5 nl) was deposited at the microinjection site and the animal was perfused transcardially with




Fig. 1. Schematic representation of the surgical preparation of the experimental animals (**A**) and the experimental design with the protocols applied in four experimental groups (**B**). A group of animals received α_2 -adrenergic receptor antagonist-yohimbine intravenously (YOHIMBINE_{i.v.}) and a control group received saline intravenously (CONTROL_{i.v.}) prior to exposure to the acute intermittent hypercapnia (AIHc) protocol. Subsequently, a group of animals received a microinjection of yohimbine into the caudal raphe region (YOHIMBINE_{raphe}) whereas a control group received a saline microinjection into the caudal raphe region (CONTROL_{raphe}) prior to exposure to the AIHc protocol. RSNA: renal sympathetic nerve activity; BP: blood pressure; ABS: arterial blood status; FiO_2 : fraction of inspired oxygen; T_i : inspiratory time; Freq: ventilator frequency; T0: baseline conditions immediately preceding the first hypercapnic episode; THc1-5: five hypercapnic episodes; T15: 15 min following the end of the last hypercapnic episode.

saline, followed by Zamboni's fixative (4 % formaldehyde and 15 % picric acid in 0.1 M PBS (phosphate buffered saline)). The brainstem was then removed and stored in Zamboni's fixative for 24 h at 4 °C. Following fixation, the brainstem was washed with 0.1 M PBS and cut in 50 μ m coronal sections using a vibrating microtome (Vibratome Series 1000, Pelco 101; Vibratome, St. Louis, MO, USA). The injection sites were verified using a conventional microscope and determined in respect to a reference section from the atlas of Paxinos and Watson [30].

Data analysis

Recorded variables were measured at 7 predetermined time points: immediately before the onset of the first hypercapnic episode (T0), during each of five hypercapnic episodes (THc1-THc5) and at 15 min following the end of the last hypercapnic episode (T15) using LabChart 8.1.13. for Windows software (ADInstruments). At each experimental time point, 20-second intervals were used to analyze RSNA, MAP and HR. The integrated signal was used to quantify the RSNA by measuring the area under the curve (i.e. calculating the integral from the minimum) in arbitrary units. To allow for comparison within and across groups, RSNA values were reported as a percentage of the

baseline activity for each experiment. MAP was derived from the blood pressure signal and expressed in mm Hg. HR was calculated from the arterial blood pressure waveform using average cycling rate function and expressed in beats per minute. All data analyses were conducted in MedCalc statistical package, version 19.1.2 (MedCalc Software, Mariakerke, Belgium). The normality of the data distribution was verified using Shapiro-Wilk's test of normality for all studied variables. Two-way repeated measures ANOVA with a *post hoc* Bonferroni correction was used for multiple comparisons within groups for all experimental variables. One-way ANOVA with Student-Newman-Keuls test was used for pairwise comparisons of corresponding experimental data points between groups. Data are reported as mean \pm standard error of the mean (SEM). Statistical significance was set at $P < 0.05$.

Results

Systemic blockade of α_2 -adrenergic receptors

Intravenous application of yohimbine evoked significant increases in RSNA during all five hypercapnic episodes when compared to the baseline (THc1: 169.2 ± 10.0 %, $P = 0.009$; THc2: 178.9 ± 11.0 %, $P = 0.008$; THc3: 180.7 ± 13.2 %, $P = 0.023$; THc4: 179.0 ± 12.7 %, $P = 0.023$; THc5: 179.0 ± 12.7 %, $P = 0.023$).

$P=0.019$; THc5: $173.2\pm 11.5\%$, $P=0.018$; $F=30.81$, $df=6$, $P<0.001$, ANOVA, Fig. 2), whereas in the control group a significant increase was observed only during the first hypercapnic episode (THc1: $126.0\pm 5.2\%$, $P=0.036$; $F=5.82$, $df=5$, $P=0.014$, ANOVA, Fig. 2). A significant interaction between hypercapnic episodes and treatment was found ($F=10.84$, $df=6$, $P=0.001$, ANOVA). Pair-wise comparisons revealed that RSNA activation was greater during all hypercapnic episodes (THc1-5) and at T15 in the yohimbine group than in the control group (Fig. 2). At baseline, MAP was significantly lower in the yohimbine

group in comparison to the control group (54.9 ± 1.9 vs. 94.3 ± 7.2 mm Hg, $F=35.57$, $df=1$, $P<0.001$, ANOVA), but no significant differences in HR were observed. During the hypercapnic episodes, significant decreases in HR were observed in both studied groups. In the control group, AIHc did not evoke significant changes in MAP, while in the yohimbine group MAP was significantly higher at THc4, THc5 and T15 when compared to the baseline (71.1 ± 3.7 mm Hg, $P=0.018$; 73.4 ± 3.3 mm Hg, $P=0.004$; 67.3 ± 3.3 mm Hg, $P=0.028$ vs. 54.9 ± 1.9 mm Hg, respectively; $F=11.31$, $df=6$, $P<0.001$, ANOVA, Table 1).




Fig. 2. (A) Time-course of changes in the renal sympathetic nerve activity (RSNA) during exposure to the acute intermittent hypercapnia (AIHc) protocol. RSNA response to AIHc was greater during all five hypercapnic episodes in a group that received yohimbine intravenously (YOHIMBINE_{i.v.}, $n=8$, dark blue, \blacklozenge) than in a control group that received saline intravenously (CONTROL_{i.v.}, $n=7$, orange, \bullet). Moreover, when yohimbine was microinjected into the caudal raphe region (YOHIMBINE_{raphe}, $n=12$, blue, \blacksquare), the RSNA response to AIHc was greater during all five hypercapnic episodes than in a control group that received saline into the same region (CONTROL_{raphe}, $n=10$, red, \blacktriangle) (*significantly different from corresponding baseline value; \dagger significantly different from corresponding T15; #significantly different from respective control group; ANOVA, $P<0.05$). **(B)** Tracings of representative experiments in four experimental groups, from top to bottom: an experiment involving the systemic blockade of the α_2 -adrenergic receptors (YOHIMBINE_{i.v.}) and a respective control experiment (CONTROL_{i.v.}) followed by an experiment with the blockade of the α_2 -adrenergic receptors in the caudal raphe region (YOHIMBINE_{raphe}) and a respective control experiment (CONTROL_{raphe}) each showing arterial blood pressure (BP; mm Hg, red), integrated renal sympathetic nerve activity (IRSNA; arbitrary units, a.u., blue) and raw renal sympathetic nerve activity (RSNA; arbitrary units, a.u., green) at seven experimental time points. Scale bar represents 20 s. T0: baseline conditions immediately preceding the first hypercapnic episode; THc1-5: five hypercapnic episodes; T15: 15 min following the end of the last hypercapnic episode.

Blockade of α_2 -adrenergic receptors in the caudal raphe region

Control group. A significant increase in RSNA was observed during the first three hypercapnic episodes when compared to the baseline ($F=5.54$, $df=6$, $P=0.014$, ANOVA; Fig. 2). There were no significant changes in RSNA at 15 min following the end of the last hypercapnic episode when compared to the baseline (T15: $103.1\pm 9.8\%$ baseline, $F=5.54$, $df=6$, $P=1.000$, ANOVA; Fig. 2). No significant changes in MAP were

observed during or following the AIHc protocol ($F=1.37$, $df=6$, $P=0.280$, ANOVA; Table 1). HR was significantly lower when compared to the baseline during all five hypercapnic episodes ($F=143.85$, $df=6$, $P<0.001$, ANOVA; Table 1). There were no significant changes in pH and PaCO₂ at 15 min following the last hypercapnic episode, while PaO₂ was significantly higher at T15 when compared to the baseline (299.5 ± 5.3 mm Hg vs. 286.6 ± 3.9 mm Hg, $F=18.83$, $df=1$, $P=0.002$, ANOVA; Table 2).

Table 1. Mean arterial blood pressure (MAP, mm Hg) and heart rate (HR, beats/min) in experimental groups at all experimental time points.

Experimental groups		T0	THc1	THc2	THc3	THc4	THc5	T15
<i>Systemic α_2-adrenergic receptor modulation</i>								
Control (n=7)	MAP	94.3±7.2	95.7±4.3	100.1±5.32	99.9±5.7	98.9±5.5	98.7±5.4	99.0±7.0
	HR	371.4±8.0	337.0±7.7* [†]	340.1±7.3* [†]	338.4±7.5* [†]	338.0±7.3* [†]	336.9±7.3* [†]	377.0±8.5
Yohimbine (n=9)	MAP	54.9±1.9 [#]	64.5±4.0	67.6±4.3	71.9±5.3	71.1±3.7*	73.4±3.3*	67.3±3.3*
	HR	375.1±6.0	348.1±5.7* [†]	348.1±5.9	348.2±5.8* [†]	344.0±6.3* [†]	345.4±5.1* [†]	376.2±4.3
<i>Central α_2-adrenergic receptor modulation</i>								
Control (n=10)	MAP	89.4±4.5	86.5±3.5	89.1±3.4	91.0±3.8	90.7±3.8	90.7±4.4	86.0±4.3
	HR	331.3±7.96	283.5±8.06* [†]	281.6±6.53* [†]	281.0±8.22* [†]	279.6±7.56* [†]	277.5±6.93* [†]	327.2±6.96
Yohimbine (n=12)	MAP	82.0±3.4	82.1±3.6	85.0±3.9	88.3±3.8 [†]	87.9±3.5 [†]	88.1±3.7 [†]	75.8±3.5
	HR	344.5±8.18	304.1±9.38* [†]	306.4±9.66* [†]	306.1±8.34* [†]	304.6±8.71* [†]	302.1±8.63* [†]	339.4±8.33

Data are presented as mean ± SEM. T0: baseline value immediately before the first hypercapnia; THc1: first hypercapnia; THc2: second hypercapnia; THc3: third hypercapnia; THc4: fourth hypercapnia; THc5: fifth hypercapnia; T15: 15 min following the end of the last hypercapnic episode. (* significantly different from respective baseline, [†] significantly different from respective T15, [#] significantly different from respective control group; P<0.05).

Table 2. Arterial blood gas analysis results in four experimental groups showing pH values and partial pressures of carbon dioxide (PaCO₂, mm Hg) and oxygen (PaO₂, mm Hg) at two experimental time points.

Experimental groups	T0			T15		
	pH	PaCO ₂	PaO ₂	pH	PaCO ₂	PaO ₂
<i>Systemic α_2-adrenergic receptor modulation</i>						
Control (n=7)	7.300±0.005	46.2±1.3	281.5±8.5	7.279±0.012	49.3±2.3	292.5±7.3
Yohimbine (n=9)	7.314±0.021	44.0±2.4	316.6±8.7	7.255±0.023	47.4±3.1	319.2±7.3
<i>Central α_2-adrenergic receptor modulation</i>						
Control (n=10)	7.271±0.015	42.8±1.5	286.6±3.9	7.262±0.018	44.8±2.8	299.5±5.3*
Yohimbine (n=12)	7.288±0.016	44.3±1.5	312.3±8.8	7.293±0.020	43.7±2.8	313.1±12.9

Data are presented as mean ± SEM. T0: baseline value immediately before the first hypercapnic episode; T15: 15 min following the end of the last hypercapnic episode. (* significantly different from respective baseline, P<0.05).

Yohimbine group. A significant increase in RSNA was observed during all five hypercapnic episodes when compared to the baseline. At T15, RSNA was significantly lower when compared to all five hypercapnic episodes, but not compared to the baseline (F=27.40, df=6, P<0.001, ANOVA; Fig. 2). MAP was lower at T15 when compared to last three hypercapnic episodes, but not compared to the baseline (F=7.26, df=6, P=0.001, ANOVA; Table 1). During THc1-5, HR was significantly lower when compared to the baseline and T15 (F=78.57, df=6, P<0.001, ANOVA; Table 1). No differences in HR were observed between baseline and T15 (344.5±8.2 vs. 339.4±8.3 beats/min, F=78.57, df=6, P=1.000, ANOVA; Table 1). There were no significant changes in pH, PaCO₂ or PaO₂ at T15 when compared to

the baseline (Table 2).

Control vs. yohimbine group. A significant interaction between hypercapnic episodes and treatment was found (F=4.44, df=6, P=0.012, ANOVA). AIHc elicited a greater increase in RSNA during all five hypercapnic episodes in the group that received a yohimbine microinjection into the caudal raphe region when compared to the control group (THc1: 138.0±4.0 % vs. 123.7±4.8 %, F=5.32, df=1, P=0.032; THc2: 137.1±5.0 % vs. 124.1±4.5 %, F=3.63, df=1, P=0.071; THc3: 143.1±6.4 % vs. 122.0±4.8 %, F=6.39, df=1, P=0.020; THc4: 146.1±6.2 % vs. 120.7±5.7 %, F=8.85, df=1, P=0.007 and THc5: 143.2±7.7 % vs. 119.2±7.2 % baseline, F=4.97, df=1, P=0.038, ANOVA, Fig. 2). At T15 no differences in RSNA were observed between




Fig. 3. Photomicrograph of a coronal section of the brainstem showing the microinjection site in the caudal raphe region marked by diluted India ink dye (blue) and indicated by the arrow. Microinjection coordinates were: +0.2 mm anteroposterior, 0.0 mm mediolateral and -2.5 mm dorsoventral relative to the obex. Scale bar represents 500 μ m.

yohimbine and control groups (T15: $103.1 \pm 9.8\%$ vs. $94.6 \pm 7.3\%$ baseline, $F=0.49$, $df=1$, $P=0.490$, ANOVA, Fig. 2).

Histological verification of the microinjection sites. Microinjection sites into the caudal raphe region were verified histologically, as shown in Figure 3.

Discussion

This study demonstrated that the blockade of α 2-adrenergic receptors by yohimbine enhanced the renal sympathetic nerve response to the acute intermittent hypercapnia in urethane-anesthetized rats. Moreover, when yohimbine was microinjected into the caudal raphe region, more pronounced increases in RSNA were evoked than in the control group, indicating that the caudal raphe region might have an important role in the regulation of sympathetic outflow during exposure to the AIHc.

The present finding that systemic blockade of α 2-adrenergic receptors profoundly affects the RSNA response to AIHc is not unexpected as α 2-adrenergic receptors are known to generally mediate

a sympathoinhibitory role and are involved in blood pressure homeostasis [31,32]. Following intravenous administration, yohimbine (α 2-adrenergic receptor antagonist) acts *via* receptors located throughout the peripheral vasculature and enters the brain rapidly where it binds to α 2-adrenergic receptors, widely distributed at multiple sites involved in sympathetic control [31,33,34]. Therefore, blocking the α 2-adrenergic receptors systemically does not allow for drawing clear conclusions about the mechanism and the precise site of action of yohimbine.

Hypercapnia is known to potently activate distinct brainstem neuronal groups and might evoke different responses as a consequence of acute or chronic hypercapnic exposures [10,11,15,35]. The conventional setting for subjecting animals to AIHc consists of cyclically changing the CO_2 fraction in different background gas mixtures and as such, the model yielded dose-dependent responses related to hypercapnia severity [8-12]. Previous studies have shown that the acute exposure to a severe hypercapnic stimulus (15% CO_2) lasting 30 min to 24 h results in changes of noradrenaline, dopamine and serotonin concentrations at various CNS regions [36,37]. Many previous studies provided

evidence that hypercapnia, by means of peripheral and central chemoreception, leads to increases in respiratory and sympathetic activity [1,3,14,15]. The retrotrapezoid nucleus (RTN) is widely recognized as the principal site of the central CO₂ chemoreception [19], essential in respiratory response to hypercapnia [15]. However, the increase in sympathetic activity produced by hypercapnia is partially dependent on the activity of the RTN neurons [15], along with various medullary regions with chemoreception properties, including the raphe neurons [27,38-40]. Thus, we aimed to investigate the role of the caudal raphe region in the sympathetic response to severe AIHc.

The caudal raphe region has been recognized as a source of input to numerous medullary and spinal sites, where it can exert an influence on many different functions, including cardiovascular control and autonomic activity [6,39]. It has been established that the caudal raphe nuclei send extensive direct projections to the sympathetic preganglionic neurons of the IML cell column [24,26,39]. Moreover, the raphe nuclei may indirectly contribute to the overall sympathetic output by means of projections to other brainstem regions involved in sympathetic control such as the RVLM [29] and RTN [15,41]. Additionally, the caudal raphe region has reciprocal connections with the RVLM [24,29] and the LC [21,23], regions that play a central role in the regulation of the autonomic activity at rest and in stressful conditions. Altogether, these anatomical projections suggest that the regulation of the arterial pressure, heart rate and sympathetic activity may be mediated by the caudal raphe region.

Previous studies proposed that hypercapnia-induced release of noradrenaline acting on α 2-adrenergic receptors of the caudal raphe region leads to its inhibition and lowers the release of serotonin henceforth affecting the descending pathways [9,11,42-44]. Thus, the increased effect of AIHc on RSNA following α 2-adrenergic receptor blockade in the caudal raphe region in our study might be a consequence of disinhibition of the caudal raphe region by the LC noradrenergic neurons. Based on the findings of this study, we might speculate that disinhibition of the caudal raphe region probably led to the increased direct input to the IML, but also *via* other regions that modulate the sympathetic activity such as the RVLM and RTN [29,41]. This speculation is supported by the results of similar studies that found that disinhibition of the raphe pallidus leads to increases in RSNA and splanchnic SNA [38,45].

One might speculate whether anesthesia might affect the sympathetic nerve discharge. However, urethane anesthesia has been shown to have minimal effects on cardiorespiratory and sympathetic activity and is commonly used in experiments requiring the preservation of the reflex response loops as the results closely resemble those observed in conscious animals [15,45,46]. Another consideration is the possible recruitment of the serotonergic signaling in relatively robust exposures to severe hypercapnic stimuli (15 % CO₂), which may activate the wake promoting pathways [5]. Considering this experimental design is used as a model of obstructive sleep apnea (OSA), we find this stimulus to be appropriate to evoke similar effects to those seen during the airway obstruction episodes in OSA, which, among other consequences, lead to hypercapnia and arousal. Yohimbine has been reported to have different affinities toward α 2-adrenergic receptor subtypes and is also known to bind to α 1-adrenergic receptors [47]. Since both α 1- and α 2-adrenergic receptors are present in the caudal raphe region [34], it is possible that the sympathoexcitatory effect observed in this study may be due to stimulation of the α 1-adrenergic receptors. Moreover, yohimbine is known to evoke responses *via* other receptors, including serotonin 5-HT_{1A} receptors, at which it can act as a partial agonist [48,49]. Since microinjections of yohimbine were performed into the caudal raphe region, it may be possible that some yohimbine bound to the 5-HT_{1A} receptors and influenced the observed effects. However, at concentration applied and considering the 80-fold higher affinity for α 2-adrenergic than for 5HT_{1A} receptors in the rat [49], it is unlikely that yohimbine produced notable effects *via* 5HT_{1A} receptors in this study. Finally, the primary stimulus for the central chemoreceptor response to changes in PaCO₂ has not been identified up to now and it has been argued whether it is solely related to the changes in CO₂ or it combines several other physiological parameters such as extracellular/intracellular pH, bicarbonate concentration or a combination of these stimuli [19,50]. In this study, we used a paradigm centered around changing the inspiratory CO₂ fraction, which is likely to have produced hypercapnic acidosis. However, it is difficult to define the primary chemoreceptor stimulus eliciting the observed changes in the renal sympathetic nerve response using this type of *in vivo* experiment.

The results of the current study indicate that the chemical blockade of the α 2-adrenergic receptors in the

caudal raphe region enhanced the renal sympathetic nerve response to the acute intermittent hypercapnia in anesthetized rats. Thus, we conclude that the neurons in the caudal raphe region make a significant contribution to the renal sympathoexcitatory response evoked by severe acute intermittent hypercapnia.

Conflict of Interest

There is no conflict of interest.

Acknowledgements

Funding for this study was provided by the Croatian science foundation project “Translational research on neuroplasticity of breathing and effect of intermittent hypoxia in anesthesia and sleep” granted to Prof. Zoran Dogas (grant number IP-11-2013-5935). The authors are grateful for Mrs. Jelena Baricevic’s continuous technical support in the laboratory. Furthermore, the authors would like to thank Dr. Linda Lusic Kalcina for statistics assistance and Prof. Zoran Valic for the advice on RSNA analysis methodology.

References

1. Makeham JM, Goodchild AK, Costin NS, Pilowsky PM. Hypercapnia selectively attenuates the somato-sympathetic reflex. *Respir Physiol Neurobiol* 2004;140:133-143. <https://doi.org/10.1016/j.resp.2003.11.003>
2. Dogas Z, Stuth EA, Hopp FA, McCrimmon DR, Zuperku EJ. NMDA receptor-mediated transmission of carotid body chemoreceptor input to expiratory bulbospinal neurones in dogs. *J Physiol* 1995;487:639-651. <https://doi.org/10.1113/jphysiol.1995.sp020906>
3. Oikawa S, Hirakawa H, Kusakabe T, Nakashima Y, Hayashida Y. Autonomic cardiovascular responses to hypercapnia in conscious rats: the roles of the chemo- and baroreceptors. *Auton Neurosci* 2005;117:105-114. <https://doi.org/10.1016/j.autneu.2004.11.009>
4. Takakura AC, Moreira TS. Contribution of excitatory amino acid receptors of the retrotrapezoid nucleus to the sympathetic chemoreflex in rats. *Exp Physiol* 2011;96:989-999. <https://doi.org/10.1113/expphysiol.2011.058842>
5. Guyenet PG. Regulation of breathing and autonomic outflows by chemoreceptors. *Compr Physiol* 2014;4:1511-1562. <https://doi.org/10.1002/cphy.c140004>
6. Richerson GB. Serotonergic neurons as carbon dioxide sensors that maintain pH homeostasis. *Nat Rev Neurosci* 2004;5:449-461. <https://doi.org/10.1038/nrn1409>
7. Andrzejewski K, Budzinska K, Kaczynska K. Effect of 6-OHDA on hypercapnic ventilatory response in the rat model of Parkinson's disease. *Physiol Res* 2019;68:285-293. <https://doi.org/10.33549/physiolres.933949>
8. Baker TL, Fuller DD, Zabka AG, Mitchell GS. Respiratory plasticity: Differential actions of continuous and episodic hypoxia and hypercapnia. *Respir Physiol* 2001;129:25-35. [https://doi.org/10.1016/s0034-5687\(01\)00280-8](https://doi.org/10.1016/s0034-5687(01)00280-8)
9. Kinkead R, Bach KB, Johnson SM, Hodgeman BA, Mitchell GS. Plasticity in respiratory motor control: intermittent hypoxia and hypercapnia activate opposing serotonergic and noradrenergic modulatory systems. *Comp Biochem Physiol A Mol Integr Physiol* 2001;130:207-218. [https://doi.org/10.1016/s1095-6433\(01\)00393-2](https://doi.org/10.1016/s1095-6433(01)00393-2)
10. Stipica I, Pavlinac Dodig I, Pecotic R, Dogas Z, Valic Z, Valic M. Periodicity during hypercapnic and hypoxic stimulus is crucial in distinct aspects of phrenic nerve plasticity. *Physiol Res* 2016;65:133-143. <https://doi.org/10.33549/physiolres.933012>
11. Stipica Safic I, Pecotic R, Pavlinac Dodig I, Dogas Z, Valic Z, Valic M. Phrenic long-term depression evoked by intermittent hypercapnia is modulated by serotonergic and adrenergic receptors in raphe nuclei. *J Neurophysiol* 2018;120:321-329. <https://doi.org/10.1152/jn.00776.2017>
12. Valic M, Pecotic R, Pavlinac Dodig I, Valic Z, Stipica I, Dogas Z. Intermittent hypercapnia-induced phrenic long-term depression is revealed after serotonin receptor blockade with methysergide in anaesthetized rats. *Exp Physiol* 2016;101:319-331. <https://doi.org/10.1113/EP085161>
13. Huang J, Lusina S, Xie T, Ji E, Xiang S, Liu Y, Weiss JW. Sympathetic response to chemostimulation in conscious rats exposed to chronic intermittent hypoxia. *Respir Physiol Neurobiol* 2009;166:102-106. <https://doi.org/10.1016/j.resp.2009.02.010>
14. Moreira TS, Takakura AC, Colombari E, Guyenet PG. Central chemoreceptors and sympathetic vasomotor outflow. *J Physiol* 2006;577:369-386. <https://doi.org/10.1113/jphysiol.2006.115600>

15. Takakura AC, Colombari E, Menani JV, Moreira TS. Ventrolateral medulla mechanisms involved in cardiorespiratory responses to central chemoreceptor activation in rats. *Am J Physiol Regul Integr Comp Physiol* 2011;300:R501-R510. <https://doi.org/10.1152/ajpregu.00220.2010>
16. Madirazza K, Pecotic R, Pavlinac Dodig I, Valic M, Dogas Z. Hyperoxia blunts renal sympathetic nerve activity response to acute intermittent hypercapnia in rats. *J Physiol Pharmacol* 2019;70:737-746. <https://doi.org/10.26402/jpp.2019.5.09>
17. Barnett WH, Abdala AP, Paton JF, Rybak IA, Zoccal DB, Molkov YI. Chemoreception and neuroplasticity in respiratory circuits. *Exp Neurol* 2017;287:153-164. <https://doi.org/10.1016/j.expneurol.2016.05.036>
18. Guyenet PG, Stornetta RL, Abbott SB, Depuy SD, Fortuna MG, Kanbar R. Central CO₂ chemoreception and integrated neural mechanisms of cardiovascular and respiratory control. *J Appl Physiol* (1985) 2010;108:995-1002. <https://doi.org/10.1152/jappphysiol.00712.2009>
19. Guyenet PG, Stornetta RL, Souza G, Abbott SBG, Shi Y, Bayliss DA. The retrotrapezoid nucleus: Central chemoreceptor and regulator of breathing automaticity. *Trends Neurosci* 2019;42:807-824. <https://doi.org/10.1016/j.tins.2019.09.002>
20. Moreira TS, Takakura AC, Damasceno RS, Falquetto B, Totola LT, Sobrinho CR, Ragioto DT, Zolezi FP. Central chemoreceptors and neural mechanisms of cardiorespiratory control. *Braz J Med Biol Res* 2011;44:883-889. <https://doi.org/10.1590/s0100-879x2011007500094>
21. Biancardi V, Bicego KC, Almeida MC, Gargaglioni LH. Locus coeruleus noradrenergic neurons and CO₂ drive to breathing. *Pflugers Arch* 2008;455:1119-1128. <https://doi.org/10.1007/s00424-007-0338-8>
22. Elam M, Yao T, Thoren P, Svensson TH. Hypercapnia and hypoxia: chemoreceptor-mediated control of locus coeruleus neurons and splanchnic, sympathetic nerves. *Brain Res* 1981;222:373-381. [https://doi.org/10.1016/0006-8993\(81\)91040-4](https://doi.org/10.1016/0006-8993(81)91040-4)
23. Samuels ER, Szabadi E. Functional neuroanatomy of the noradrenergic locus coeruleus: its roles in the regulation of arousal and autonomic function part II: physiological and pharmacological manipulations and pathological alterations of locus coeruleus activity in humans. *Curr Neuropharmacol* 2008;6:254-285. <https://doi.org/10.2174/157015908785777193>
24. Jacobs BL, Martin-Cora FJ, Fornal CA. Activity of medullary serotonergic neurons in freely moving animals. *Brain Res Brain Res Rev* 2002;40:45-52. [https://doi.org/10.1016/s0165-0173\(02\)00187-x](https://doi.org/10.1016/s0165-0173(02)00187-x)
25. Allen GV, Cechetto DF. Serotonergic and nonserotonergic neurons in the medullary raphe system have axon collateral projections to autonomic and somatic cell groups in the medulla and spinal cord. *J Comp Neurol* 1994;350:357-366. <https://doi.org/10.1002/cne.903500303>
26. Loewy AD, Neil JJ. The role of descending monoaminergic systems in central control of blood pressure. *Fed Proc* 1981;40:2778-2785.
27. Morrison SF. Raphe pallidus excites a unique class of sympathetic preganglionic neurons. *Am J Physiol* 1993;265:R82-R89. <https://doi.org/10.1152/ajpregu.1993.265.1.R82>
28. Pilowsky PM, Lung MS, Spirovski D, McMullan S. Differential regulation of the central neural cardiorespiratory system by metabotropic neurotransmitters. *Philos Trans R Soc Lond B Biol Sci* 2009;364:2537-2552. <https://doi.org/10.1098/rstb.2009.0092>
29. Bago M, Marson L, Dean C. Serotonergic projections to the rostroventrolateral medulla from midbrain and raphe nuclei. *Brain Res* 2002;945:249-258. [https://doi.org/10.1016/s0006-8993\(02\)02811-1](https://doi.org/10.1016/s0006-8993(02)02811-1)
30. Paxinos G, Watson C. *The Rat Brain in Stereotaxic Coordinates, 4th Edition*. Academic Press, San Diego, 1998, 256 p.
31. Philipp M, Brede M, Hein L. Physiological significance of alpha(2)-adrenergic receptor subtype diversity: one receptor is not enough. *Am J Physiol Regul Integr Comp Physiol* 2002;283:R287-R295. <https://doi.org/10.1152/ajpregu.00123.2002>
32. Lin YH, Liu YP, Lin YC, Lee PL, Tung CS. Cooling-evoked hemodynamic perturbations facilitate sympathetic activity with subsequent myogenic vascular oscillations via alpha2-adrenergic receptors. *Physiol Res* 2017;66:449-457. <https://doi.org/10.33549/physiolres.933385>

33. Hubbard JW, Pfister SL, Biediger AM, Herzig TC, Keeton TK. The pharmacokinetic properties of yohimbine in the conscious rat. *Naunyn Schmiedebergs Arch Pharmacol* 1988;337:583-587. <https://doi.org/10.1007/BF00182736>
 34. Unnerstall JR, Kopajtic TA, Kuhar MJ. Distribution of alpha 2 agonist binding sites in the rat and human central nervous system: analysis of some functional, anatomic correlates of the pharmacologic effects of clonidine and related adrenergic agents. *Brain Res* 1984;319:69-101. [https://doi.org/10.1016/0165-0173\(84\)90030-4](https://doi.org/10.1016/0165-0173(84)90030-4)
 35. Burgraff NJ, Neumueller SE, Buchholz KJ, LeClaire J, Hodges MR, Pan L, Forster HV. Brainstem serotonergic, catecholaminergic, and inflammatory adaptations during chronic hypercapnia in goats. *FASEB J* 2019;33:14491-14505. <https://doi.org/10.1096/fj.201901288RR>
 36. Garcia de Yébenes Prous J, Carlsson A, Mena Gomez MA. The effect of CO₂ on monoamine metabolism in rat brain. *Naunyn Schmiedebergs Arch Pharmacol* 1977;301:11-15. <https://doi.org/10.1007/BF00501258>
 37. Stone EA. Rapid adaptation of the stimulatory effect of CO₂ on brain norepinephrine metabolism. *Naunyn Schmiedebergs Arch Pharmacol* 1983;324:313-315. <https://doi.org/10.1007/BF00502629>
 38. Horiuchi J, McAllen RM, Allen AM, Killinger S, Fontes MA, Dampney RA. Descending vasomotor pathways from the dorsomedial hypothalamic nucleus: role of medullary raphe and RVLM. *Am J Physiol Regul Integr Comp Physiol* 2004;287:R824-R832. <https://doi.org/10.1152/ajpregu.00221.2004>
 39. Lovick TA. The medullary raphe nuclei: a system for integration and gain control in autonomic and somatomotor responsiveness? *Exp Physiol* 1997;82:31-41. <https://doi.org/10.1113/expphysiol.1997.sp004013>
 40. Nattie E, Li A. Central chemoreceptors: Locations and functions. *Compr Physiol* 2012;2:221-254. <https://doi.org/10.1002/cphy.c100083>
 41. Leirao IP, Colombari DSA, da Silva GSF, Zoccal DB. Lesion of serotonergic afferents to the retrotrapezoid nucleus impairs the tachypneic response to hypercapnia in unanesthetized animals. *Neuroscience* 2021;452:63-77. <https://doi.org/10.1016/j.neuroscience.2020.11.005>
 42. Bach KB, Mitchell GS. Hypercapnia-induced long-term depression of respiratory activity requires alpha2-adrenergic receptors. *J Appl Physiol* (1985) 1998;84:2099-2105. <https://doi.org/10.1152/jappl.1998.84.6.2099>
 43. Dobbins EG, Feldman JL. Brainstem network controlling descending drive to phrenic motoneurons in rat. *J Comp Neurol* 1994;347:64-86. <https://doi.org/10.1002/cne.903470106>
 44. Haddjeri N, Blier P, de Montigny C. Effect of the alpha-2 adrenoceptor antagonist mirtazapine on the 5-hydroxytryptamine system in the rat brain. *J Pharmacol Exp Ther* 1996;277:861-871.
 45. Morrison SF. RVLM and raphe differentially regulate sympathetic outflows to splanchnic and brown adipose tissue. *Am J Physiol* 1999;276:R962-R973. <https://doi.org/10.1152/ajpregu.1999.276.4.R962>
 46. Maggi CA, Meli A. Suitability of urethane anesthesia for physiopharmacological investigations in various systems. Part 1: General considerations. *Experientia* 1986;42:109-114. <https://doi.org/10.1007/bf01952426>
 47. Goldberg MR, Robertson D. Yohimbine: a pharmacological probe for study of the alpha 2-adrenoreceptor. *Pharmacol Rev* 1983;35:143-180.
 48. Winter JC, Rabin RA. Yohimbine as a serotonergic agent: evidence from receptor binding and drug discrimination. *J Pharmacol Exp Ther* 1992;263:682-689.
 49. Zaretsky DV, Zaretskaia MV, DiMicco JA, Rusyniak DE. Yohimbine is a 5-HT1A agonist in rats in doses exceeding 1 mg/kg. *Neurosci Lett* 2015;606:215-219. <https://doi.org/10.1016/j.neulet.2015.09.008>
 50. Wang W, Bradley SR, Richerson GB. Quantification of the response of rat medullary raphe neurones to independent changes in pH(o) and P(CO₂). *J Physiol* 2002;540:951-970. <https://doi.org/10.1113/jphysiol.2001.013443>
-